

GROTHENDIECK TOPOLOGIES AND IDEAL CLOSURE OPERATIONS

HOLGER BRENNER

ABSTRACT. We relate closure operations for ideals and for submodules to non-flat Grothendieck topologies. We show how a Grothendieck topology on an affine scheme induces a closure operation in a natural way, and how to construct for a given closure operation fulfilling certain properties a Grothendieck topology which induces this operation. In this way we relate the radical to the surjective topology and the constructible topology, the integral closure to the submersive topology, to the proper topology and to Voevodsky's h -topology, the Frobenius closure to the Frobenius topology and the plus closure to the finite topology. The topologies which are induced by a Zariski filter yield the closure operations which are studied under the name of hereditary torsion theories. The Grothendieck topologies enrich the corresponding closure operation by providing cohomology theories, rings of global sections, concepts of exactness and of stalks.

Mathematical Subject Classification (2000): primary: 18F10; secondary: 13A10; 13A15; 13A35; 13B02; 13B22; 13C99; 13D30; 14A15; 14F05; 14F20; 18A99; 18B25; 18E15; 18F10; 54B40

Keywords: Grothendieck topology, sheaves, integral closure, Frobenius, plus closure, tight closure, radical, submersion, torsion theory.

INTRODUCTION

Hilbert's Nullstellensatz states that for a polynomial $f \in \mathbb{C}[X_1, \dots, X_m]$ which vanishes on the common zero set $V(f_1, \dots, f_n) \subseteq \mathbb{C}^m$ of a set of polynomials there exists $k \in \mathbb{N}$ such that $f^k \in (f_1, \dots, f_n) = I$, or, in other words, that f belongs to the *radical* $\text{rad}(I)$ of I . This theorem is probably the first instance in history where an ideal is replaced by a bigger ideal which reflects better certain properties of the ideal in the given context. The *integral closure* $\bar{I}$ of an ideal in a commutative ring is the biggest ideal with the property that it has the same Hilbert-Samuel multiplicity as the given ideal. It was first studied by Zariski in the context of equisingularity problems. Northcott and Rees [51] enriched this notion introducing the theory of reductions and analytic spread.

In the last 20 years, the *tight closure* I^* , as defined by Hochster and Huneke in positive characteristic ([35], [38]), has proved useful in homological algebra, commutative algebra, invariant theory, singularity theory and algebraic geometry. Tight closure comes along with *plus closure* (conjecturally the same), *Frobenius closure*, *regular closure*, *diamond closure* and *solid closure*. It can be extended to rings containing a field of characteristic 0 by reduction to positive characteristic and there have been several attempts to obtain closure operations with similar features also in mixed characteristic ([33], [6], [30], [31], [32]). These constructions are related to the search for *big Cohen-Macaulay modules* and *algebras*. Other closure operations include the *weak subintegral closure*, the *blow-up*

closure, Ratliff's $\triangle$ -*closure*, the *2,3-closure* and the *continuous closure* ([67], [13], [55], [52], [7])

These closure operations have more or less natural extensions to submodules. Also, a coherent subsheaf of a locally free sheaf is often replaced by its *reflexive hull* or by its *saturation* in the sense that the quotient is torsion-free. Moreover, some constructions in commutative algebra can be thought of as a closure of 0 inside an R -module M , like its *torsion* or the *submodule of sections with support* inside a given closed subset of $\text{Spec } R$, as it is used in the definition of local cohomology. The idea of torsion has been systematically studied under the name of *hereditary torsion theories* in the categorical context, but in particular for modules over a ring ([3], [16]).

On the other hand, *Grothendieck topologies*, in particular the *étale topology* for schemes, were introduced to give for varieties over a finite field a suitable replacement for the complex topology, in order to define algebraically a cohomology theory with similar properties as the singular cohomology in algebraic topology ([1], [24], [49]). This development culminated in Deligne's proof of the Weil conjectures.

The concept of a Grothendieck topology axiomatizes the idea of an *open set* and of *coverings*. However, the open sets need neither be subsets nor is a covering something which can be tested by containment of points. Instead, coverings are given as a collection of morphisms $U_i \rightarrow U$, $i \in I$, fulfilling certain structural conditions, and points need not to exist at all. With this notion of coverings at hand, one can talk about *local* and *global* properties and one can define *presheaves*, *sheaves*, *sheafification*, *exactness* and *cohomology*.

The basic idea of this (and subsequent) paper is to describe closure operations for ideals and submodules over a commutative ring R as a *sheafification procedure* in a suitable Grothendieck topology over $X = \text{Spec } R$. More generally, for a morphism between *ringed sites* (i.e. categories with a Grothendieck topology and a sheaf of rings) we can talk about *extension* and *contraction* of modules and hence of a closure operation. For R -modules $N \subseteq M$, a Grothendieck topology on X denoted by X_{top} yields sheafified $\mathcal{O}_{\text{top}}$ -modules $N_{\text{top}} \rightarrow M_{\text{top}}$ on X_{top} and an image sheaf $N^{\text{top}} \subseteq M_{\text{top}}$. Then we declare for $s \in M$ that $s \in N^c$ if and only if $\theta(s) \in \Gamma(X_{\text{top}}, N^{\text{top}})$, where $\theta : M \rightarrow \Gamma(X_{\text{top}}, M_{\text{top}})$ is the sheafification homomorphism. This means, in a certain sense, that $s \in N^c$ holds if “ $s \in N$ ” holds *locally* in the Grothendieck topology, and this means that it holds for a covering. In particular, $s \in 0^c$, if $s = 0$ holds locally in the topology.

The classical Grothendieck topologies, like the Zariski, the flat and the étale topology (this is also true for the Nisnevich topology [50], which is between the Zariski and the étale topology, and for the *primitive topology* of M. Walker [69]) do not yield in this way anything new, i.e. $N^c = N$. This is due to the fact that the assignment $V \mapsto \Gamma(V, j^*N)$ for $j : V \rightarrow X = \text{Spec } R$ flat (denoted by $W(M)$ in [24, Example VII.2c]) and in [49]), is already a sheaf in the flat topology [49, Proposition I.2.18 and Corollary II.1.6], and since a faithfully flat homomorphism $R \rightarrow S$ is *pure*, which means that $M \rightarrow M \otimes_R S$ is injective for every R -module M . If $\varphi : R \rightarrow S$ is however not pure, then the inclusion $I \subset \varphi^{-1}(IS)$ might be strict, and if we declare φ to be a covering, then $\varphi(f) \in IS$ says that f belongs locally to the ideal I , hence to I^c .

Lets explain more concretely how a Grothendieck topology induces a closure operation and what properties it has. Lets restrict for simplicity in this introduction to the *affine single-handed* case. This means that we have a category of R -algebras, closed under composition and tensor product, and certain R -algebra homomorphisms are declared to

be covers (so all coverings consist of just one morphism). In this situation the closure operation given by sheafification in the topology boils down to saying that for $f \in R$ and an ideal $I \subseteq R$ the containment $f \in I^c$ holds if and only if there exists a cover $\varphi : R \rightarrow S$ such that $\varphi(f) \in IS$. More generally, for R -submodules $N \subseteq M$ and $s \in M$, the containment $s \in N^c$ holds if and only if there exists such a cover such that $s \otimes 1$ is in the image of $N \otimes_R S \rightarrow M \otimes_R S$. It follows immediately that $N \subseteq N^c = (N^c)^c$ holds and that the closure operation is *order preserving*, that is for $N \subseteq N'$ in M one has $N^c \subseteq (N')^c$.

A great deal of this project is in finding the right concept of covers for a given closure operation and to study properties of these morphisms instead (or rather as a way) of studying the closure operation. For some closure operations it is immediately clear which ring homomorphisms should be considered to be a cover, but not for all. We go through the closure operations which we will deal with in this and in subsequent papers ([4], [5], [7], [8]) and we describe the natural covers for them.

Let's look first at the *Frobenius closure* in positive characteristic p , which is defined by $f \in I^F$ if and only if there exists a power of the Frobenius $\Phi^e : R \rightarrow R$, $f \mapsto f^{p^e}$, such that $f^{p^e} \in \Phi^e(I) =: I^{[p^e]}$. In this case one wants to allow exactly the Φ^e to be covers to define the *Frobenius topology*. However, in order to get a theory with good properties regarding base change and also to have only covers of finite type, it is better to define a homomorphism $R \rightarrow S$ of finite type to be a cover in the Frobenius topology if there exists a factorization $\Phi^e : R \rightarrow S \rightarrow R$ for some e such that $\text{Spec } R \rightarrow \text{Spec } S$ is surjective.

The *plus closure* of an ideal I in a domain R consists of all elements $f \in R$ such that there exists a finite extension of domains $R \subseteq S$ with $f \in IS$. Here the appropriate concept of a cover, which works also for rings with zero divisors, is to impose that $R \rightarrow S$ is finite and with surjective morphism $\text{Spec } S \rightarrow \text{Spec } R$. This gives the *finite topology*.

What is the appropriate concept of a cover for the *radical* of an ideal (and what is the radical of a submodule)? If we take Hilbert's Nullstellensatz as our guide, then f belongs to $\text{rad}(I)$ if and only if $V(f) \supseteq V(I)$ in $\text{Spec } R$. This means for all prime ideals $I \subseteq \mathfrak{p}$ that also $f \in \mathfrak{p}$ holds, or that all homomorphisms $R \rightarrow \kappa(\mathfrak{p})$ which map I to 0 also map f to 0. This property is then true for all homomorphisms from R to a field. Which ring homomorphisms $R \rightarrow S$ behave nicely with respect to homomorphism to fields? There are several possibilities, but we will use the fact that $\text{Spec } S \rightarrow \text{Spec } R$ is surjective if and only if for all $R \rightarrow K$ we have $S \otimes_R K \neq 0$ (or $\text{Spec}(S \otimes_R K) \neq \emptyset$).

Declaring *spec-surjective morphisms* to be covers is in fact a way to obtain the radical via a Grothendieck topology, the *surjective topology*. For, if $R \rightarrow S$ is spec-surjective and $f \in IS$, then for all homomorphisms to a field $\psi : R \rightarrow K$ it follows by the faithful flatness of $K \rightarrow S \otimes_R K$ ($\neq 0$) that $\psi(f) \in IK$. Hence if $IK = 0$, then also $\psi(f) = 0$.

For the other direction we have to construct for $f \in \text{rad}(I)$ a spec-surjective homomorphism $\varphi : R \rightarrow S$ with $\varphi(f) \in IS$. For this we use *forcing algebras*, as introduced by Hochster [33] in connection with solid closure in an (at the end not successful) attempt to construct a closure operation in all characteristics with tight closure properties [33]. Fix ideal generators $I = (f_1, \dots, f_n)$. The forcing algebra for these data is $B = R[T_1, \dots, T_n]/(f_1 T_1 + \dots + f_n T_n + f)$. In B , $f \in IB$ holds, and every R -algebra S with $f \in IS$ factors (non uniquely) through B . It is clear by this property that forcing algebras should play an important role in the study of closure operations. In the context of the radical, $f \in \text{rad}(I)$ holds if and only if the corresponding forcing algebra induces a spec-surjective morphism.

There are several equivalent characterizations of the *integral closure* of an ideal. If R is noetherian, then $f \in \bar{I}$ if and only if $f \in IV$ for all ring homomorphisms $R \rightarrow V$ to discrete valuation domains. Hence a forcing algebra $\text{Spec } B \rightarrow \text{Spec } R$ should be declared to be a cover if and only if for all $\text{Spec } V \rightarrow \text{Spec } R$ there exists a lifting $\text{Spec } V \rightarrow \text{Spec } B$. For forcing algebras, but not in general, this path-lifting property is equivalent to say that for every $R \rightarrow V$ the homomorphism $V \rightarrow B \otimes_R V$ has a V -module section or that it is pure. There exists even a topological characterization for an V -algebra T to be pure, namely that over the prime ideals in $\text{Spec } V$, i.e. $0 \subset \mathfrak{m}_V$, there exists a pair of prime ideals $\mathfrak{q} \subset \mathfrak{p}$ in $\text{Spec } T$ mapping to 0 and $\mathfrak{m}_V$ respectively.

This topological observation ties in purity after base change to discrete valuation domains with submersions. A *submersion* $\varphi : \text{Spec } S \rightarrow \text{Spec } R$ is a surjective mapping such that the Zariski topology of $\text{Spec } R$ is the image (or quotient) topology under the morphism, that is, a subset $W \subseteq \text{Spec } R$ is open if and only if its preimage $\varphi^{-1}(W)$ is open. This definition can be found in SGA 1 (see [23, Définition IX.2.1] and [21, Définition 3.10.1]) and also in [68, Definition 3.1.1] under the name of *topological epimorphism*. A morphism $\varphi : \text{Spec } S \rightarrow \text{Spec } R$ (R noetherian) is *universally submersive* if and only if for all $R \rightarrow V$ to discrete valuation domains $V \rightarrow S \otimes_R V$ is pure ([23, Remarque IX.2.6], [53, Théorème 37]). Hence the *submersive topology*, in which the universal submersions are covers, gives a Grothendieck topology which induces the integral closure. Surprisingly, this topology is very closely related to Voevodsky's *h-topology*, which arose as a new approach to understand the algebraic topology of varieties and which was the first systematic study of a non-flat Grothendieck topology ([68], [64]). We will study submersions and integral closure in a separate paper [4].

Recall that for a ring of positive characteristic p the *tight closure* of an ideal I is by definition $I^* = \{f \in R : \text{there exists } z \in R^\circ, z f^q \in I^{[q]} \text{ for all } q = p^e\}$, where R° denotes the complement of all minimal primes. Tight closure agrees in positive characteristic (if R has a completely stable test element) with *solid closure* [33, Theorem 8.6]. This means that the containment $f \in I^*$ for an $\mathfrak{m}$ -primary ideal I , where $\mathfrak{m}$ is a maximal ideal of height d , is characterized by the property that the local cohomology module $H_{\mathfrak{m}}^d(B)$ is not zero, where B is the forcing algebra [33, Corollary 2.4]. This property makes sense in every (even mixed) characteristic, leading to *solid closure* in general and to the *solid topology*. It is however known that for rings containing a field of characteristic 0 solid closure does not exhibit all features one expects from a tight closure type theory, in particular, ideals in a regular ring are not always solidly closed [60]. A variant of solid closure, called *parasolid closure*, introduced in [6], is defined characteristic free by the property that all cohomology classes $1/g_1 \cdots g_d \in H_{\mathfrak{m}}^d(R)$, where $g_1, \dots, g_d$ is a system of parameters in R , do not vanish in $H_{\mathfrak{m}}^d(B)$. We will continue the search for the *tight topology* along these lines in [5].

Another source for closure operations is given by *torsion theories*. Torsion theories have been studied in various categorical contexts; for the category of R -modules over a commutative ring it means that we have two classes $\mathcal{T}$ and $\mathcal{F}$ of modules ('*torsion modules*' and '*torsion-free modules*' in an abstract sense) such that $\mathcal{T}$ is closed under quotients and $\mathcal{F}$ is closed under submodules and such that for every R -module M there exists a short exact sequence $0 \rightarrow T \rightarrow M \rightarrow F \rightarrow 0$ ($T \in \mathcal{T}$, $F \in \mathcal{F}$). In particular then, every module has a *biggest torsion submodule*, and this can be considered as the closure of 0. A torsion theory is called *hereditary* if the torsion class is also closed under submodules.

A hereditary torsion theory is basically given by its so-called *Gabriel filter*, which consists of all ideals $J \subseteq R$ such that R/J is torsion. In the noetherian case, this Gabriel filter is stable under radical, so it is essentially a (Zariski-)topological filter. Such a *Zariski filter* gives rise to a (non-affine) Grothendieck topology, where the Zariski open subset $V \subseteq \operatorname{Spec} R$ is said to be a cover if V belongs to this filter. So here the covers are flat, but they are not surjective. This is quite a natural construction, since the relevant information lies often already in certain open subsets, like e.g. the divisor class group depends only on points of codimension one. The induced closure of a submodule N consists of all $s \in M$ such that the restriction $s|_V \in N$ for some open subset in the filter. For $N = 0$ and one fixed $U = D(\mathfrak{a})$ this also equals $\Gamma_{V(\mathfrak{a})}(M) = \Gamma_{\mathfrak{a}}(M)$, the *module of sections with support in $V(\mathfrak{a})$* , as used in the definition of local cohomology. In general, a typical feature for hereditary torsion theories is that the closure operation $0 \mapsto 0^c$ is left exact.

So for a lot of closure operations we can describe a Grothendieck topology which induces it, but one can also give an exact characterization of the closures for which one can construct a Grothendieck topology. This construction uses again forcing algebras. Assume that $f \in (f_1, \dots, f_n)^c$ in a given closure operation c . Then it is natural to declare the corresponding forcing algebra B to be a cover. It may then happen that $g \in JB$, where J is another ideal, so g will belong to the induced closure of J in every topology where $R \rightarrow B$ is a cover. So in order that this construction does not produce too many elements in the closure we need that $g \in J^c$. We express this requirement by saying that *c-admissible forcing algebras* must be *c-pure*. This property characterizes exactly the closure operations which come from a Grothendieck topology. It implies that the closure is *module persistent*, i.e. for a module homomorphism $\varphi : M \rightarrow M'$ one has $\varphi(N^c) \subseteq \varphi(N)^c$ and that it is *independent of the presentation*, i.e. if $\varphi : M \rightarrow M/N$, then $N^c = \varphi^{-1}(0^c)$.

An important feature of closure operations are *persistence properties* with respect to ring changes, which should be reflected by the Grothendieck topologies. A closure operation is said to be persistent with respect to a class of ring homomorphisms if for $\varphi : R \rightarrow S$ and $I \subseteq R$ one has $\varphi(I^c) \subseteq \varphi(I)^c$. Most closure operations mentioned above (with the exception of torsion theories) are persistent with respect to all ring homomorphisms; for tight closure this is a non-trivial fact. If we have a Grothendieck topology on a category of rings, then a ring homomorphism $R \rightarrow S$ might or might not respect the Grothendieck topology, i.e. induces a *site morphism* $(\operatorname{Spec} S)_{\text{top}} \rightarrow (\operatorname{Spec} R)_{\text{top}}$ or not. If it does, then the closure operation is persistent with respect to this homomorphism.

After having explained how a Grothendieck topology induces a closure operation we shall look now at some other features which a Grothendieck topology provides. Some of these features have occurred in the study of closure operations in some disguise, and Grothendieck topologies give a conceptual view on these.

The containment in the Frobenius closure can also be expressed by saying that $f \in IR^\infty$, where R^∞ is the *perfect closure* of R . The containment in the plus closure is equivalent with $f \in IR^+$, where R^+ is the integral closure of R in an algebraic closure of its quotient field $Q(R)$ (R a domain), called the *absolute integral closure*. The study of tight closure is closely related to the search for *big Cohen-Macaulay modules* or *algebras*. The common feature here is that there exists one “big” algebra which encompasses in a certain sense all relevant covers, like in topology the universal covering space encompasses all unramified covers. We systemize this observation with the concept of an *absolute filter* in a Grothendieck topology, at which we can evaluate a (pre-)sheaf to get an *absolute stalk*. This is similar

to the étale topology of a local ring, where a suitable colimit over all étale covers yields the *strict Henselization* of the ring. An absolute stalk should carry essentially the whole information of the Grothendieck topology and of the closure operation, but its construction and its explicit description might be difficult, and its functorial properties are weak in general.

A Grothendieck topology provides a concept of *exactness* for a complex of R -modules $N \rightarrow M \rightarrow L$, namely the exactness of the complex of sheaves $N_{\text{top}} \rightarrow M_{\text{top}} \rightarrow L_{\text{top}}$ on the underlying category of the Grothendieck topology. This notion depends not only on the concept of a covering, but also on the size of the category. Quite often we will work with a *small site*, where (roughly spoken) only objects occur which are relevant for some covering of the final object $\text{Spec } R$. This exactness on the small site is still a strong condition and we will also consider weaker notions which are closely related to the property that the kernel in the complex of modules is contained in the closure of the image. This property has been studied in tight closure theory under the name of *phantom homology* [38, Chapter 10] and for the integral closure (for the Koszul complex) by D. Katz [41]. In the surjective topology, the sheaf exactness is equivalent to the property that $N \otimes_R K \rightarrow M \otimes_R K \rightarrow L \otimes_R K$ is exact for every ring homomorphism $R \rightarrow K$ to a field.

A Grothendieck topology on $X = \text{Spec } R$ gives also for an R -module M via sheafification a *global module of sections*, namely $\Gamma(X_{\text{top}}, M_{\text{top}})$ (with a homomorphism from M to it), and in particular a *global ring of sections* $\Gamma(X_{\text{top}}, \mathcal{O}_{\text{top}})$. An element in this ring is represented by a compatible element $f \in S$, where $R \rightarrow S$ is a cover and where compatible means that $f \otimes 1 - 1 \otimes f \in S \otimes_R S$ is 0 in the topology, i.e. in some cover $S \otimes_R S \rightarrow T$. The assignment $R \mapsto \Gamma(X_{\text{top}}, \mathcal{O}_{\text{top}})$ is functorial for ring homomorphisms $S \rightarrow S$ which respect the Grothendieck topology.

For the Frobenius topology this construction yields the perfect closure, since for $f \in {}^e R$ the element $f \otimes 1 - 1 \otimes f$ is nilpotent. In this case the ring of global sections equals the absolute stalk. In the surjective topology (in characteristic 0), a global function is essentially given by an affine *constructible partition* $X = V_1 \uplus \dots \uplus V_k$, where V_i is an intersection of a closed and an open subset, and functions $f_i \in \Gamma(V_i, \mathcal{O}_{V_i})$ (an absolute stalk is given as the product of the algebraic closures of all residue class fields).

For other topologies the ring of global sections is closer to the ring itself, e.g. for the plus closure and in the submersive topology (in characteristic 0), it is the so-called *seminormalization*. In particular, $f \in I^c$ does not mean that $f \in I\Gamma(X_{\text{top}}, \mathcal{O}_{\text{top}})$, which is a global and much stronger condition.

As the module of global sections is the 0th cohomology, we arrive at *sheaf cohomology* in general. Cohomology can be defined quite generally via injective resolutions, and at least the first cohomology might be computed by *Čech cohomology*. The cohomology of a coherent $\mathcal{O}_X$ -module over an affine scheme might be non-trivial, contrary to the flat, étale or Zariski topology. A typical short exact sequence arises from ideal generators $I = (f_1, \dots, f_n)$ and gives $0 \rightarrow \text{Syz} \rightarrow \mathcal{O}_{\text{top}}^n \xrightarrow{f_1, \dots, f_n} I^{\text{top}} \rightarrow 0$, where $I^{\text{top}} \subseteq \mathcal{O}_{\text{top}}$ is the image sheaf of I_{top} inside $\mathcal{O}_{\text{top}}$ and where Syz is defined as the kernel of the surjection (not as Syz_{top}). The corresponding long cohomology sequence is $0 \rightarrow \Gamma(X_{\text{top}}, \text{Syz}) \rightarrow \oplus_n \Gamma(X_{\text{top}}, \mathcal{O}_{\text{top}}) \xrightarrow{f_1, \dots, f_n} \Gamma(X_{\text{top}}, I^{\text{top}}) \rightarrow H^1(X_{\text{top}}, \text{Syz}) \rightarrow$

$\oplus_n H^1(X_{\text{top}}, \mathcal{O}_{\text{top}})$. In the case of the Frobenius topology and of the surjective topology we can prove that $H^1(X_{\text{top}}, \mathcal{O}_{\text{top}}) = 0$ and also that $H^1(X_{\text{top}}, \text{Syz}) = 0$, and so all $f \in I^c$ come from a ‘global solution’ $f \in \Gamma(X_{\text{top}}, \mathcal{O}_{\text{top}})$.

In cases where $\Gamma(X_{\text{top}}, \mathcal{O}_{\text{top}}) = R$ we have $\Gamma(X_{\text{top}}, I^{\text{top}}) = I^c \subseteq R$ and the image from the left is $\Gamma(X_{\text{top}}, \mathcal{O}_{\text{top}}) = I$, so $I^c/I \subseteq H^1(X_{\text{top}}, \text{Syz})$, which is a bijection provided that $H^1(X_{\text{top}}, \mathcal{O}_{\text{top}}) = 0$. We expect this to hold in the submersive topology for a normal domain in characteristic 0. The relation between the first cohomology of the sheaf of syzygies and the closure quotient I^c/I extends to a relation between the naive sequence $0 \rightarrow I^c/I \rightarrow R/I \rightarrow R/I^c \rightarrow 0$ and the long exact sequence $\dots \rightarrow H^1(X_{\text{top}}, \text{Syz}) \rightarrow H^1(U_{\text{top}}, \text{Syz}) \rightarrow H^2_{X_{\text{top}}-U_{\text{top}}}(\text{Syz}) \rightarrow \dots$ relating local and global cohomology, where $U = D(I)$.

If a closure operation or a Grothendieck topology is defined for a class of rings (usually the class of all noetherian rings), then it is natural to ask for which rings the Grothendieck topology is *pure* and the closure is trivial. Purity means here that the sheafification does not annihilate anything, and for a single-handed Grothendieck topology it just means that every cover is pure. A stronger condition is that every cover allows a module section or even that every cover allows a ring section, like in the étale topology the algebraically simply connected varieties are characterized by the property that every étale cover has a section. The class of rings (or singularities) which are pure with respect to a certain Grothendieck topology might be quite significant, which is in particular true for tight closure, where the so-called *F-regular* rings are closely related to rational singularities.

We believe that Grothendieck topologies provide a natural framework to study closure operations. It gives a conceptual view on certain features which have been loosely accompanying closure operations ever since (like exactness or absolute covers). Once the ‘right’ Grothendieck topology for a given closure operation for ideals is found we can immediately extend the closure to submodules.

The focus on morphisms coming along with studying the appropriate coverings brings in a geometric flavor to questions on closure of ideals. For example, the complex-topological characterization of submersions yields at once the complex-analytic characterization of the integral closure of submodules. Also, statements on closures have often natural generalizations to morphisms, in such a way that the original statement is regained by looking at the morphisms given by forcing algebras.

Grothendieck topologies provide also a new way to construct closure operations. Here the most challenging problem is to give in mixed characteristic an operation with similar properties as tight closure. This would solve most remaining homological conjectures. Less ambitious is the task to give a tight closure like operation in equal characteristic zero without reduction to positive characteristic. This was essentially accomplished in [6], but a complex-analytic characterization extending the relation between tight closure test ideals and multiplier ideals remains to be done.

We give a brief summary of the content and the structure of this paper.

We start in part 1 with the concepts of Grothendieck topologies, sheaves (1.1), sheafification (1.2) and of morphisms of (ringed) sites (1.3). A morphism of (ringed) sites is the natural setting in which it makes sense to talk about extensions and contractions of (sub-)modules, which gives a closure operation (1.4). This approach has the advantage of being a relative setting, so that we can sheafify and get the closures in several steps. Later on our site morphisms will be between a scheme with its Zariski topology (or even

its trivial topology, where only isomorphisms are coverings) and another larger (or finer) topology. In (1.5) we discuss such Grothendieck topologies on a scheme and how a module defines a (pre-)sheaf. In (1.6) we discuss modules of global sections and provide some lemmata which will later be of use in computations for several Grothendieck topologies.

Starting with (1.7) we look at filters in a Grothendieck topology and the stalks they lead to. Basically, a filter is a cofiltered system (or diagram) in the topology. An easy way to produce filters is by a fixation (Construction 1.7.2). In some sense, a filter, and in particular the irreducible ones, are a replacement for points (1.8). A filter which consists only of covers and has the property that for every cover it has a factorization is called *absolute* (1.9). They carry in some sense the whole information of the Grothendieck topology. In (1.10) we discuss several notions of exactness and in (1.11) we discuss cohomology, Čech cohomology and local cohomology in a site. It might be a good idea to read initially only (1.4) of the first part and go back if needed.

The second part relates Grothendieck topologies on a scheme, in particular on an affine scheme, with closure operations. In (2.1) we start with the basics about forcing algebras. In (2.2) we introduce *admissible closure operations* which will turn out to be the ones for which one can construct a Grothendieck topology. We also introduce certain finiteness conditions for a closure operation. In (2.3) we give examples of admissible closure operations and also of non-admissible closures like the Ratliff-Rush closure. In (2.4) we describe, on the basis of (1.4), how a Grothendieck topology on a scheme induces an admissible closure operation. In (2.5) we characterize the pure topologies and in (2.6) we construct for a given admissible closure a Grothendieck topology which induces it. In (2.7) we relate the notion of exactness with properties of the closure, and in (2.8) we deal with the sheaf of syzygies and how the closure operation is reflected in the cohomology of this sheaf.

The following parts are dedicated in detail to specific closure operations and their “natural” Grothendieck topologies and can be read independently from each other.

Part 3 deals with the radical and the surjective topology. In (3.1) we recall the radical of an ideal and introduce the *radical of a submodule*. The surjective topology is introduced in (3.2) and it is shown that it induces the radical (Proposition 3.2.3). In (3.3) we discuss the *Jacobson radical* and the *Jacobson topology*, where a cover need only be surjective on the closed points. It follows from a version of the Nakayama Lemma that for a finitely generated module the Jacobson radical and the radical is not the whole module (Corollary 3.3.8). In (3.4) we deal with the surjective topology over a field and show that the ring of global sections is the perfect closure of the field. Note that though over a field a ring is faithfully flat if and only if it is spec-surjective, the surjective topology over a field is not the same as the faithfully flat topology, since also the covers of $A \otimes_K A$ are crucial for the topology. In (3.5) we show that the direct product over the algebraic closures of all residue class fields of $\text{Spec } R$ provides an absolute stalk (Proposition 3.5.1).

In (3.6) we discuss *constructible partitions* of a scheme, which are the easiest non-trivial surjective mappings, and which are rich enough to induce also the radical. Constructible partitions are easier to handle than arbitrary surjections, yet they contain often the full information of the surjective topology and many questions can be reduced to the *constructible topology*. Section (3.7) provides some lemmata for this. In (3.8) we show that the ring of global sections in the surjective topology for a ring of finite (Krull) dimension over a field of characteristic 0 is the same as in the constructible topology (Theorem 3.8.2).

So a global function is given by a collection of algebraic functions on a constructible partition of the scheme. The exactness in the surjective topology is characterized in (3.9) and is equivalent to the exactness after tensoration with an arbitrary field. It is also equivalent to global exactness, which is rather untypical for a Grothendieck topology. Finally, in (3.10) we prove that the first cohomology of coherent sheaves vanishes in the surjective topology (Theorem 3.10.2 and Corollary 3.10.3).

Part 4 deals with the *Frobenius topology* and the *Frobenius closure* in positive characteristic. In (4.1) we recall the definition of Frobenius closure of modules and in (4.2) we introduce the Frobenius topology which induces the Frobenius closure (Proposition 4.2.3) and we characterize the purity of the Frobenius topology (Proposition 4.2.4). In (4.3) we compute the ring of global sections in the Frobenius topology to be the perfect closure (Theorem 4.3.3). In (4.4) we characterize exactness in the Frobenius topology and in (4.5) we show that the first cohomology of coherent modules vanishes (Proposition 4.5.1 and Corollary 4.5.2).

Part 5 deals with the *plus closure* and the *finite topology*. In (5.1) we recall the plus closure and introduce the finite topology which induces the plus closure. In (5.2) we show that the ring of global sections for a domain of characteristic zero in the finite topology is the seminormalization, i.e. the maximal subring of the normalization which does not separate points (Proposition 5.2.3). The absolute stalks in the finite topology, which are just products of the absolute integral closures, are described in (5.3). Our results on the computation of cohomology in the finite topology are rather modest (5.4).

Part 6 is concerned with *torsion theories* and topologies given by *Zariski filters*. These topologies are flat but not surjective (and not affine). A topological filter in the Zariski topology defines a Grothendieck topology by declaring the open sets inside the filter to be covers (6.1). For a filter given by just one open set the closure of 0 is nothing but the module of sections with support in the closed complement, as used in the definition of local cohomology (6.2). Section (6.3) provides the link between Grothendieck topologies given by a Zariski filter and hereditary torsion theories (Theorem 6.3.1). In (6.4) we look at the *divisorial topology*, where the Zariski filter consists of all open subsets which contain all points of codimension one. The module of global sections is under certain condition the *reflexive hull* of the module, i.e. its bidual (Lemma 6.4.2), and the closure of a submodule inside a reflexive module over a normal ring is also the reflexive hull (Proposition 6.4.3). We also describe the relation to *symbolic powers* and we show that the divisor class group is the Picard group in the divisorial topology (Proposition 6.4.4).

In the final part 7 we treat some further Grothendieck topologies. In (7.1) we show how to define, starting from an ideal $\mathfrak{a}$ in a ring R , a Grothendieck topology such that the ring of global sections is the $\mathfrak{a}$ -adic completion of R . Here a covering needs infinitely many objects. In (7.2) we treat the *proper topology* (first introduced by S.-I. Kimura in [42]), where a cover is given by a proper morphism. We show in Proposition 7.2.5 that the induced closure is the *integral closure*. We will come back to this in the study of submersive topologies [4]. Section (7.3) finally studies the so-called $\triangle$ -closure of Ratliff, which is given by a multiplicatively closed set of ideals. This closure is induced by taking the blow-ups of these ideals as covers (Proposition 7.3.1).

We do not touch in this paper the following topics: the topos of modules defined by a Grothendieck topology, e.g. structure theory for the $\mathcal{O}_{\text{top}}$ -modules, the question when scheme morphisms induce isomorphisms of sites, free resolutions in the topology,

minimal number of generators, reductions, core, projective dimension, Picard group in the Grothendieck topology (the multiplicative theory), other non-quasicoherent sheaves like constant sheaves, the algebro-topological or homotopical content of the Grothendieck topologies, extension problems for covers.

I thank R. Buchweitz, A. Conca, G. Dietz, D. Eisenbud, N. Epstein, T. Gaffney, S. Goto, R. Hartshorne, M. Hashimoto, R. Heitmann, A. Kaid, M. Katzman, E. Kunz, M. Morales, P. Roberts, E. Sbarra, R. Sharp, V. Srinivas, M. Vitulli, K.I. Watanabe for their interest and useful remarks. Most of all I thank Manuel Blickle for critical remarks and many helpful discussions during the whole project.

1. GROTHENDIECK TOPOLOGIES

1.1. Presheaves, Grothendieck topologies and sheaves.

An (abelian) *presheaf* on a category $\mathcal{T}$ is a contravariant functor $\mathcal{F} : \mathcal{T} \rightarrow \mathcal{A}$, where $\mathcal{A}$ is the category of abelian groups (or another abelian category). For a morphism $j : U \rightarrow V$ we will often denote $\mathcal{F}(j) : \mathcal{F}(V) \rightarrow \mathcal{F}(U)$ by $\rho_U^V(j)$ or simply by ρ_U^V and call this the *restriction* from V to U . For a topological space, a sheaf is a presheaf which satisfies certain properties with respect to coverings. To talk about sheaves in a given category one has to define a notion of coverings with certain structural properties (see [1], [24], [49], [66]). This is the basic idea of a Grothendieck topology.

Recall that a *product* in a category for $U \rightarrow W, V \rightarrow W$ is an object $U \times_W V$ with projections to U and V compatible over W such that for every $T \rightarrow U, T \rightarrow V$ compatible over W there exists a unique morphism $T \rightarrow U \times_W V$.

Definition 1.1.1. Let $\mathcal{T}$ be a category with products. A *Grothendieck topology* (sometimes called a pretopology, see [24, Définition II.1.1] or [66, Definition 2.23 and Remark 2.24]) on $\mathcal{T}$ consists of a collection of coverings, that is, families $\varphi_i : U_i \rightarrow U, i \in I$, where in each family the target $U \in \mathcal{T}$ is fixed. These coverings should satisfy.

- (i) Isomorphisms are coverings.
- (ii) If $\varphi_i : U_i \rightarrow U, i \in I$, and $\varphi_{ij} : U_{ij} \rightarrow U_i, j \in J_i$, are coverings for every i , then so is the composition $\varphi_i \circ \varphi_{ij} : U_{ij} \rightarrow U, (i, j) \in \bigsqcup_{i \in I} J_i$.
- (iii) If $U_i \rightarrow U$ is a covering and $V \rightarrow U$ is any morphism in $\mathcal{T}$, then $V \times_U U_i \rightarrow V$ is a covering.

Each covering $U_i \rightarrow U$ in a Grothendieck topology induces the diagram (which is the first part of the Čech diagram)

$$\bigsqcup_{i,j} U_i \times_U U_j \rightrightarrows \bigsqcup_i U_i \rightarrow U$$

(the disjoint unions do not have to exist in $\mathcal{T}$), where the arrows $\bigsqcup_{i,j} U_i \times_U U_j \rightarrow \bigsqcup_i U_i$ are on each open component either the first or the second projection. For a presheaf $\mathcal{F}$ we get the Čech diagram of the covering with values in $\mathcal{F}, \mathcal{F}(U) \rightarrow \prod_i \mathcal{F}(U_i) \rightrightarrows \prod_{i,j} \mathcal{F}(U_i \times_U U_j)$.

Definition 1.1.2. Let $\mathcal{T}$ be a Grothendieck topology. Then a presheaf $\mathcal{F}$ is called a *sheaf*, if for all coverings $U_i \rightarrow U$ in the topology the sequence

$$\mathcal{F}(U) \rightarrow \prod_{i \in I} \mathcal{F}(U_i) \rightrightarrows \prod_{(i,j) \in I \times I} \mathcal{F}(U_i \times_U U_j)$$

is exact (which also means that the first map is injective).

Remark 1.1.3. Explicitly, a sheaf must satisfy the following two properties.

- (i) If $s \in \mathcal{F}(U)$ is such that $\rho_{U_i}^U(s) = 0$ for all $i \in I$, then $s = 0$.
- (ii) If $s_i \in \mathcal{F}(U_i)$ is such that $(\rho_{U_i \times_U U_j}^{U_i})(s_i) = (\rho_{U_i \times_U U_j}^{U_j})(s_j)$ for all i, j , then there exists an $s \in \mathcal{F}(U)$ such that $\rho_{U_i}^U(s) = s_i$.

A presheaf which satisfies only the first condition is called a *separated presheaf*. Note that the sheaf condition is in this general setting, contrary to the case of a topological space, also non-trivial for a single cover $V \rightarrow U$ (we will usually prefer the term cover for a covering consisting of only one object), because the two projections $p_1, p_2 : V \times_U V \rightarrow V$ are different.

1.2. Sheafification.

One needs to construct a *sheafification* of a presheaf $\mathcal{F}$, that is, a sheaf $\mathcal{F}^\#$ together with a presheaf homomorphism $\theta : \mathcal{F} \rightarrow \mathcal{F}^\#$ which is universal for homomorphisms from $\mathcal{F}$ to sheaves. That is, whenever the solid arrow to the sheaf $\mathcal{G}$ is given,

$$\begin{array}{ccc} & & \mathcal{G} \\ & \nearrow & \uparrow \text{dotted} \\ \mathcal{F} & \xrightarrow{\theta} & \mathcal{F}^\# \end{array}$$

the dotted arrow exists uniquely. As we will use sheafification explicitly we recall its construction. The sheafification is a two step process, see [49, Theorem II 2.11], [24, Exposé II.3] or [66, Section 2.3.7]. In the first step one defines a subpresheaf $\mathcal{F}_0 \subseteq \mathcal{F}$ by

$$\mathcal{F}_0(U) := \{s \in \mathcal{F}(U) : \exists \text{ a covering } U_i \rightarrow U, i \in I, \text{ such that } \rho_{U_i}^U(s) = 0 \ \forall i\}.$$

Then one sets $\mathcal{F}_1(U) := \mathcal{F}(U)/\mathcal{F}_0(U)$. This presheaf $\mathcal{F}_1$ is separated.

To describe the second step we pause to recall some categorical notions, in particular colimits over a (quasi)filtered system. Beside the sheafification procedure we need this to define the inverse presheaf under a site morphism (Section 1.3) and to talk about (absolute) filters and stalks in a Grothendieck topology (Section 1.7). We are explicit about this point also since not everything written down in the literature is correct (see Example 1.7.4).

Definition 1.2.1. Let Λ be a category. We call Λ *quasifiltered*, if the following property holds.

- (i) For every pair $\lambda \rightarrow \mu$ and $\lambda' \rightarrow \mu$ in Λ there exists $\kappa \in \Lambda$ and morphisms $\kappa \rightarrow \lambda$ and $\kappa \rightarrow \lambda'$ such that the compositions are the same; and for every $\lambda, \lambda' \in \Lambda$ there exists κ with arrows $\kappa \rightarrow \lambda, \kappa \rightarrow \lambda'$ (this follows from the first condition if Λ has a final object).

We call it *cofiltered* if it has also the property that

- (ii) For every pair $\psi_1, \psi_2 : \lambda \rightrightarrows \mu$ in Λ there exists $\varphi : \kappa \rightarrow \lambda$ such that $\psi_1 \circ \varphi = \psi_2 \circ \varphi$ (so φ ‘unifies’ the two morphisms).

Remark 1.2.2. If the arrows are reversed the corresponding notions are called *quasifiltered* and *filtered*. A Λ -*diagram* is a functor defined on a category Λ . We will deal with covariant topological diagrams which lead by evaluating a (pre)sheaf to contravariant diagrams in an abelian category. The *colimit* of such a contravariant diagram $\Lambda \rightarrow \mathcal{A}$ is the object A in $\mathcal{A}$ characterized by the property that there exist homomorphisms $\rho_\lambda : A_\lambda \rightarrow A$, which are compatible with respect to the homomorphisms $A_\mu \rightarrow A_\lambda$ indexed by $\lambda \rightarrow \mu$,

and that for every object $\tilde{A}$ in $\mathcal{A}$ with such compatible homomorphisms there exists a unique factorization $A \rightarrow \tilde{A}$.

The existence of colimits and their description depend on properties of the indexing category and of the category $\mathcal{A}$. Colimits exist for R -modules over a small category [12, Proposition A 6.2]. Over a small quasifiltered category the colimit of sets is the disjoint union of all sets A_λ modulo the equivalence relation which identifies two elements $a \in A_\lambda$ and $a' \in A_{\lambda'}$ if they map to the same element in some A_κ under $\psi, \psi' : A_\lambda, A_{\lambda'} \rightarrow A_\kappa$. For a quasifiltered diagram of abelian groups (or R -modules) the colimit in the category of groups is different from the colimit in the category of sets ([12, Proposition A 6.3] is wrong, see Example 1.7.4). For a filtered diagram however the two notions coincide, and the colimit in the category of groups has the easy set theoretical description.

We return now to the construction of the sheafification. In the second step one sets $\mathcal{P}^+(U) = \varinjlim \check{H}(\mathcal{U}, \mathcal{P})$ (applied then to the separated presheaf $\mathcal{P} = \mathcal{F}_1$), where $\check{H}(\mathcal{U}, \mathcal{P})$ consists of the elements $s_i \in \Gamma(U_i, \mathcal{P})$ which are compatible in the sense that $\rho_{U_i \times_U U_j}^{U_i}(s_i) = \rho_{U_i \times_U U_j}^{U_j}(s_j)$ and where the colimit is taken over all coverings of U ordered by refinements. So the objects in the indexing category are coverings $\mathcal{U} = (U_i \rightarrow U, i \in I)$, and the morphisms are refinements of coverings. A covering $V_j \rightarrow U, j \in J$, is a *refinement* of $U_i \rightarrow U, i \in I$, if there exists a mapping $\psi : J \rightarrow I$ and morphisms (in $\mathcal{T}_X$) $\theta_j : V_j \rightarrow U_{\psi(j)}$ ([49, III §2] or [66, Definition 2.43]).

A refinement $\mathcal{V} \rightarrow \mathcal{U}$ induces a homomorphism $\check{H}(\mathcal{U}, \mathcal{P}) \rightarrow \check{H}(\mathcal{V}, \mathcal{P})$ by sending $(s_i), i \in I$, to $(\theta_j(s_{\psi(j)})), j \in J$, $\theta_j = \rho(\theta_j) : \Gamma(U_{\psi(j)}, \mathcal{P}) \rightarrow \Gamma(V_j, \mathcal{P})$. Because of the compatibility property this homomorphism is independent of ψ and of θ_j [24, Corollaire V.2.3.5]. Therefore this is a filtered diagram in the category of abelian groups (though on the topological level it is not even quasicofiltered) and its colimit equals the colimit in the category of sets. In particular, the homomorphisms $\mathcal{P}(U) \rightarrow \mathcal{P}^+(U)$ are injective for a separated presheaf.

Combining the two steps gives us the sheafification $\mathcal{F}^\# = (\mathcal{F}_1)^+$ of a given presheaf $\mathcal{F}$. It is also true that $\mathcal{F}^\# = (\mathcal{F}^+)^+$, but we will use the first description.

The kernel presheaf of a sheaf morphism $\mathcal{G} \rightarrow \mathcal{F}$ is a sheaf, but the presheaf image is not. The *image sheaf* of a sheaf morphism is by definition the sheafification of the presheaf image. A complex of sheaves $\mathcal{G} \xrightarrow{\alpha} \mathcal{F} \xrightarrow{\beta} \mathcal{E}$ is called *exact* if the identity of sheaves $\text{kern}(\beta) = \text{im}(\alpha)$ holds; this means that for every $U \in \mathcal{T}_X$ and every element $s \in \Gamma(U, \mathcal{F})$ mapping to 0 in $\Gamma(U, \mathcal{E})$ there exists a covering $U_i \rightarrow U, i \in I$, and sections $t_i \in \Gamma(U_i, \mathcal{G})$, $i \in I$, (without compatibility condition) mapping to the restrictions $\rho_{U_i}^U(s) \in \Gamma(U_i, \mathcal{F})$. With these definitions the category of sheaves on $\mathcal{T}$ is an abelian category.

A complex of presheaves is called *exact* if for every U in $\mathcal{T}$ the evaluated complex of abelian groups is exact. This notion is often too strong, and it is independent of the Grothendieck topology. We will use also the following definition.

Definition 1.2.3. Let $\mathcal{T}$ be a Grothendieck topology and let $U \in \mathcal{T}$ be an object. Let $\mathcal{G} \xrightarrow{\alpha} \mathcal{F} \xrightarrow{\beta} \mathcal{E}$ be a complex of presheaves of abelian groups. Then we say that the complex is $\mathcal{T}$ -*exact* on U or U -*exact* if for every $s \in \Gamma(U, \mathcal{F})$ mapping to 0 in $\Gamma(U, \mathcal{E})$ there exists a covering $U_i \rightarrow U, i \in I$, and sections $t_i \in \Gamma(U_i, \mathcal{G})$ mapping to the restrictions $\rho_{U_i}^U(s)$ in $\Gamma(U_i, \mathcal{F})$. We say that the complex is $\mathcal{T}$ -*exact* if it is U -exact for every $U \in \mathcal{T}$.

Proposition 1.2.4. *Let $\mathcal{G} \xrightarrow{\alpha} \mathcal{F} \xrightarrow{\beta} \mathcal{E}$ be a complex of presheaves on a Grothendieck topology $\mathcal{T}$. Then the following holds.*

- (i) *If the complex is exact as a complex of presheaves, then it is $\mathcal{T}$ -exact.*
- (ii) *If the sheafified complex is U -exact, then the complex of presheaves is U -exact.*
- (iii) *The complex is $\mathcal{T}$ -exact if and only if the associated complex of sheaves is $\mathcal{T}$ -exact.*
- (iv) *A complex of sheaves is exact if and only if it is $\mathcal{T}$ -exact.*

Proof. (i) is clear. (ii). Suppose that the sheafified complex is U -exact, and let $s \in \Gamma(U, \mathcal{F})$ mapping to 0. Then there exists a covering $U_i \rightarrow U$, $i \in I$, and elements $t_i \in \Gamma(U_i, \mathcal{G}^\#)$ mapping to $s_i = \rho_{U_i}^U(\theta(s)) \in \Gamma(U_i, \mathcal{F}^\#)$. The elements t_i are represented by sections of $\mathcal{G}$ in a finer covering and also the identities hold already in $\mathcal{F}$ in a finer covering.

(iii). One direction follows from (ii). For the other direction let $s \in \Gamma(U, \mathcal{F}^\#)$ mapping to 0 in $\Gamma(U, \mathcal{E}^\#)$. Then there exists a covering $U_i \rightarrow U$, $i \in I$, and $s_i \in \Gamma(U_i, \mathcal{F})$ representing s and coverings $V_j \rightarrow U_i$, $j \in J_i$, such that the restrictions of $\beta(s_i)$ in $\Gamma(V_j, \mathcal{E})$ are 0. Since the statement is local we may assume that $U_i = V_j = U$. But then again the assumptions on the presheaves yield immediately a covering $U_i \rightarrow U$ and sections $t_i \in \Gamma(U_i, \mathcal{G}) \rightarrow \Gamma(U_i, \mathcal{G}^\#)$ which map to the restrictions of s . (iv) follows from the definitions. $\square$

1.3. Site morphisms.

We will call a category $\mathcal{T}$ with a final object X , with arbitrary products and endowed with a Grothendieck topology a *site*, and we will denote it by $(X, \mathcal{T}_X)$ or sometimes just by X . Later on X will be a scheme, the products will be products of schemes and $\mathcal{T}_X$ will be a subcategory of schemes over X .

If $\mathcal{T}_X$ and $\mathcal{T}_Y$ are two sites with final objects X and Y , then a *site morphism* $\varphi^* : \mathcal{T}_X \rightarrow \mathcal{T}_Y$, for which we will briefly write $\varphi : Y \rightarrow X$, is a functor $\mathcal{T}_X \rightarrow \mathcal{T}_Y$ with $\varphi^*(X) = Y$ and which respects products and coverings, so that if $U_i \rightarrow U$, $i \in I$, is a covering in $\mathcal{T}_X$, then $\varphi^*(U_i) \rightarrow \varphi^*(U)$, $i \in I$, is a covering in $\mathcal{T}_Y$.

Construction 1.3.1. A site morphism $\varphi : Y \rightarrow X$ associates to a presheaf $\mathcal{P}$ of abelian groups on X a presheaf $\varphi^{-1}(\mathcal{P})$ on Y (called its *inverse image*) in the following way (cf. the construction of u_i in [24, Proposition I.5.1] in connection with [24, Proposition III.1.6]). For $V \in \mathcal{T}_Y$, let I_V^φ be the category of all pairs (U, ρ) with $U \in \mathcal{T}_X$ and $\rho \in \text{Mor}_{\mathcal{T}_Y}(V, \varphi^*(U))$. The morphisms $\text{Mor}_{I_V^\varphi}((U_1, \rho_1), (U_2, \rho_2))$ in this category are the maps $\sigma \in \text{Mor}_{\mathcal{T}_X}(U_1, U_2)$ such that $\varphi^*(\sigma) \circ \rho_1 = \rho_2$. This category is quasicofiltered, since for $\rho_1 : V \rightarrow \varphi^*(U_1)$ and $\rho_2 : V \rightarrow \varphi^*(U_2)$ mapping to $\rho_0 : V \rightarrow \varphi^*(U_0)$ via $\sigma_1 : U_1 \rightarrow U_0$ and $\sigma_2 : U_2 \rightarrow U_0$ there exists $\rho : V \rightarrow \varphi^*(U_1) \times_{\varphi^*(U_0)} \varphi^*(U_2) = \varphi^*(U_1 \times_{U_0} U_2)$ over ρ_1 and ρ_2 . We take the colimit

$$\varphi^{-1}(\mathcal{P})(V) := \varinjlim_{(U, \rho) \in I_V^\varphi} \Gamma(U, \mathcal{P}).$$

in the category of abelian groups. This indeed defines a presheaf on $\mathcal{T}_Y$, since a map $\tau \in \text{Mor}_{\mathcal{T}_Y}(V, W)$ induces a functor $I_W^\varphi \rightarrow I_V^\varphi$ by sending (U, ρ) to $(U, \rho \circ \tau)$ and thus induces a map of the colimits above.

The categories I_V^φ are in general not cofiltered, but they are under the following *cofiltered condition*. For every pair $\psi_1, \psi_2 : U \rightrightarrows U'$ in $\mathcal{T}_X$ and $\rho : V \rightarrow \varphi^*(U)$ in $\mathcal{T}_Y$ there exists $\gamma : \tilde{U} \rightarrow U$ which unifies ψ_1 and ψ_2 and $\tilde{\rho} : V \rightarrow \varphi^*(\tilde{U})$ such that $\varphi^*(\gamma) \circ \tilde{\rho} = \rho$. Under this condition $\varphi^{-1}(\mathcal{P})$ has the easy set-theoretical description.

The presheaf $\varphi^{-1}(\mathcal{P})$ constructed in 1.3.1 can be sheafified, but for us it will be more important to sheafify with respect to a structure sheaf of rings on the sites (we need the ‘image réciproque au sens modules’ instead of the ‘image réciproque ensembliste’, cf. [24, Exposés IV.11-13]).

Definition 1.3.2. A site $\mathcal{T}$ is called a *(pre-)ringed site* if there exists a fixed (pre)sheaf $U \mapsto \Gamma(U, \mathcal{O})$, $U \in \mathcal{T}$, of commutative rings, called its *structure (pre)sheaf*.

A presheaf $\mathcal{P}$ of abelian groups on such a (pre-)ringed site is called an $\mathcal{O}$ -premodule, if it has additionally an $\mathcal{O}$ -multiplication such that $\Gamma(U, \mathcal{P})$ is an $\Gamma(U, \mathcal{O})$ -module for every $U \in \mathcal{T}$. If moreover $\mathcal{P}$ is a sheaf, then we call it an $\mathcal{O}$ -module.

The sheafification of the structure presheaf $\mathcal{O}$ on a pre-ring site makes it into a ringed site. The sheafification of an $\mathcal{O}$ -premodule $\mathcal{F}$ is an $\mathcal{O}$ -module and an $\mathcal{O}^\#$ -module, since the multiplication $\mathcal{O} \times \mathcal{F} \rightarrow \mathcal{F}$ sheafifies to $\mathcal{O}^\# \times \mathcal{F}^\# \rightarrow \mathcal{F}^\#$.

Definition 1.3.3. If $(\mathcal{T}_X, \mathcal{O}_X)$ and $(\mathcal{T}_Y, \mathcal{O}_Y)$ are two (pre-)ringed sites, then a site morphism $\varphi^* : \mathcal{T}_X \rightarrow \mathcal{T}_Y$, for which we will write again $\varphi : Y \rightarrow X$, is called a *morphism of (pre)ringed sites*, if there is given a family of ring homomorphisms $\Gamma(U, \mathcal{O}_X) \rightarrow \Gamma(\varphi^*(U), \mathcal{O}_Y)$ compatible with the restrictions in the topologies.

Remark 1.3.4. A morphism of (pre-)ringed sites $\varphi : Y \rightarrow X$ induces on $\mathcal{T}_Y$ a morphism of presheaves of rings $\varphi^{-1}(\mathcal{O}_X) \rightarrow \mathcal{O}_Y$ by

$$\Gamma(V, \varphi^{-1}(\mathcal{O}_X)) = \varinjlim_{(U, \rho) \in I_V^\varphi} \Gamma(U, \mathcal{O}_X) \longrightarrow \varinjlim_{(U, \rho) \in I_V^\varphi} \Gamma(\varphi^*(U), \mathcal{O}_Y) \longrightarrow \Gamma(V, \mathcal{O}_Y).$$

(Note that $\varphi^{-1}(\mathcal{O}_X)$ is here the inverse sheaf in the category of rings, so the colimit we have to take in its construction is the colimit inside the category of rings.) On the other hand, if such a presheaf morphism is given, then the composition $\Gamma(U, \mathcal{O}_X) \rightarrow \Gamma(\varphi^*(U), \varphi^{-1}(\mathcal{O}_X)) \rightarrow \Gamma(\varphi^*(U), \mathcal{O}_Y)$ yields the ring homomorphisms between the sites.

In the case of a morphism between (pre-)ringed sites $\varphi : Y \rightarrow X$ the inverse image $\varphi^{-1}(\mathcal{F})$ of an $\mathcal{O}_X$ -(pre)module $\mathcal{F}$ is in general not an $\mathcal{O}_Y$ -(pre)module.

Definition 1.3.5. Let $\varphi : Y \rightarrow X$ be a morphism between (pre-)ringed sites. We define $\varphi^{\text{pre}}(\mathcal{F})$ to be the $\mathcal{O}_Y$ -premodule given by

$$V \longmapsto \Gamma(V, \varphi^{-1}(\mathcal{F})) \otimes_{\Gamma(V, \varphi^{-1}(\mathcal{O}_X))} \Gamma(V, \mathcal{O}_Y).$$

We denote its sheafification by $\varphi^*(\mathcal{F})$ (cf. [24, IV.13.2.2]).

$\varphi^{\text{pre}}(\mathcal{F})$ is an $\mathcal{O}_Y$ -premodule and $\varphi^*(\mathcal{F})$ is an $\mathcal{O}_Y$ -module.

Lemma 1.3.6. Let $\varphi : Y \rightarrow X$ be a morphism between ringed sites. Then φ^* is a right exact functor for $\mathcal{O}_X$ -modules.

Proof. See [24, Corollaire IV.13.6]. □

1.4. Extension and contraction under a site morphism.

A ring homomorphism $\varphi : R \rightarrow S$ defines for an ideal $I \subseteq R$ the *extended ideal* IS in S and then the *contraction* $\varphi^{-1}(IS)$, which contains I . Basically the same construction applies to an arbitrary site morphism. Later on we will specialize to the situation where the site morphism $(\text{Spec } R)_{\text{top}} \rightarrow \text{Spec } R$ is given by a Grothendieck topology on an affine scheme $\text{Spec } R$. This gives then a closure operation for ideals in R .

Definition 1.4.1. Let X and Y stand for two ringed sites $(\mathcal{T}_X, \mathcal{O}_X)$ and $(\mathcal{T}_Y, \mathcal{O}_Y)$ and let $\varphi : Y \rightarrow X$ denote a morphism of ringed sites. Let $\mathcal{G} \subseteq \mathcal{F}$ denote $\mathcal{O}_X$ -modules. This induces an $\mathcal{O}_Y$ -module homomorphisms $\varphi^*(\mathcal{G}) \rightarrow \varphi^*(\mathcal{F})$. The image sheaf of $\varphi^*(\mathcal{G})$ inside $\varphi^*(\mathcal{F})$ is called the *extended submodule* of $\mathcal{G}$ and is denoted by $\mathcal{G}^{\text{ext}}$. In particular, for an ideal sheaf $\mathcal{I} \subseteq \mathcal{O}_X$ we call the image sheaf of $\varphi^*(\mathcal{I})$ inside $\mathcal{O}_Y = \varphi^*(\mathcal{O}_X)$ the *extended ideal sheaf*, denoted by $\mathcal{I}^{\text{ext}}$.

Recall that a site morphism $\varphi : Y \rightarrow X$ defines a functor $\mathcal{H} \mapsto \varphi_*(\mathcal{H})$ for presheaves by setting $\Gamma(U, \varphi_*(\mathcal{H})) := \Gamma(\varphi^*(U), \mathcal{H})$ for $U \in \mathcal{T}_X$. This *push forward* functor sends sheaves to sheaves and is left exact. It also sends $\mathcal{O}_Y$ -modules to $\mathcal{O}_X$ -modules. In particular, for an $\mathcal{O}_Y$ -submodule $\mathcal{H} \subseteq \mathcal{H}'$ also $\varphi_*(\mathcal{H}) \subseteq \varphi_*(\mathcal{H}')$ is an $\mathcal{O}_X$ -submodule. For an $\mathcal{O}_X$ -module $\mathcal{F}$ we have a natural sheaf homomorphism $\theta : \mathcal{F} \rightarrow \varphi_*(\varphi^*(\mathcal{F}))$ given by

$$\theta : \Gamma(U, \mathcal{F}) \rightarrow \Gamma(U, \varphi_*(\varphi^*(\mathcal{F}))) = \Gamma(\varphi^*(U), \varphi^*(\mathcal{F})).$$

Definition 1.4.2. Let $\varphi : Y \rightarrow X$ be a site morphism between two ringed sites, let $\mathcal{F}$ be an $\mathcal{O}_X$ -module. An $\mathcal{O}_Y$ -submodule $\mathcal{H} \subseteq \varphi^*(\mathcal{F})$ yields an $\mathcal{O}_X$ -submodule $\varphi_*(\mathcal{H}) \subseteq \varphi_*(\varphi^*(\mathcal{F}))$. We call the $\mathcal{O}_X$ -submodule of $\mathcal{F}$ given as the preimage sheaf of $\varphi_*(\mathcal{H})$ under the natural sheaf homomorphism $\theta : \mathcal{F} \rightarrow \varphi_*(\varphi^*(\mathcal{F}))$ the *contracted module*, denoted by $\mathcal{H}^{\text{cont}}$. Explicitly, we have $\Gamma(U, \mathcal{H}^{\text{cont}}) = \{s \in \Gamma(U, \mathcal{F}) : \theta(s) \in \Gamma(U, \varphi_*(\mathcal{H})) = \Gamma(\varphi^*(U), \mathcal{H})\}$.

Definition 1.4.3. Let $\varphi : Y \rightarrow X$ be a site morphism between two ringed sites, let $\mathcal{G} \subseteq \mathcal{F}$ be $\mathcal{O}_X$ -modules. We call the contraction of the extension the *closure submodule* of $\mathcal{G}$ inside $\mathcal{F}$ via φ , and denote it by $\mathcal{G}^{\varphi-c} = \mathcal{G}^c = (\mathcal{G}^{\text{ext}})^{\text{cont}}$.

Proposition 1.4.4. Let $\varphi : Y \rightarrow X$ be a site morphism between two ringed sites, let $\mathcal{G} \subseteq \mathcal{F}$ be $\mathcal{O}_X$ -modules. Then the following holds.

- (i) The closure submodule $\mathcal{G}^c$ is an $\mathcal{O}_X$ -module and in particular a sheaf.
- (ii) We have $\mathcal{G} \subseteq \mathcal{G}^c = (\mathcal{G}^c)^c$.
- (iii) Let $\mathcal{G}' \subseteq \mathcal{F}'$ be another pair of $\mathcal{O}_X$ -modules and let $\mu : \mathcal{F} \rightarrow \mathcal{F}'$ be an $\mathcal{O}_X$ -module homomorphism with $\mu(\mathcal{G}) \subseteq \mathcal{G}'$. Then $\mu(\mathcal{G}^c) \subseteq (\mathcal{G}')^c$.
- (iv) If $\mathcal{G} \subseteq \mathcal{H} \subseteq \mathcal{F}$ is another $\mathcal{O}_X$ -submodule, then $\mathcal{G}^c \subseteq \mathcal{H}^c$.
- (v) Consider the short exact sequence $0 \rightarrow \mathcal{G} \xrightarrow{\alpha} \mathcal{F} \xrightarrow{\beta} \mathcal{E} \rightarrow 0$ of $\mathcal{O}_X$ -modules, where $\mathcal{E} = \mathcal{F}/\mathcal{G}$. Then $\mathcal{G}^c = \beta^{-1}(0^c)$.

Proof. (i) is clear from the corresponding properties of extension and contraction.

(ii). The inclusion $\mathcal{G} \subseteq \mathcal{G}^c$ and so also $\mathcal{G}^c \subseteq (\mathcal{G}^c)^c$ is clear. For the other direction note that in general $(\mathcal{H}^{\text{cont}})^{\text{ext}} \subseteq \mathcal{H}$ for a submodule $\mathcal{H} \subseteq \varphi^*(\mathcal{F})$. This is because $\varphi^*(\mathcal{H}^{\text{cont}}) \rightarrow \varphi^*(\mathcal{F})$ factors through $\mathcal{H}$, since we have $\mathcal{H}^{\text{cont}} \rightarrow \varphi_*(\mathcal{H})$ by definition and so $\varphi^*(\mathcal{H}^{\text{cont}}) \rightarrow \varphi^*\varphi_*(\mathcal{H})$, and by adjunction we get $\varphi^*\varphi_*(\mathcal{H}) \rightarrow \mathcal{H} \rightarrow \varphi^*(\mathcal{F})$.

(iii) follows from applying φ^* to the commutative diagram

$$\begin{array}{ccc} \mathcal{G} & \longrightarrow & \mathcal{F} \\ \downarrow & & \downarrow \mu \\ \mathcal{G}' & \longrightarrow & \mathcal{F}' \end{array}$$

(iv) is a special case of (iii).

(v). The inclusion $\subseteq$ follows from (iii). For the other inclusion note that we have by Lemma 1.3.6 an exact sequence of sheaves $\varphi^*(\mathcal{G}) \xrightarrow{\alpha} \varphi^*(\mathcal{F}) \xrightarrow{\beta} \varphi^*(\mathcal{E}) \rightarrow 0$ in $\mathcal{T}_Y$, where $\mathcal{G}^{\text{ext}} = \text{im}(\alpha) = \text{kern}(\beta)$ holds. If $s \in \Gamma(U, \mathcal{F})$ is such that $\beta(s) \in 0^c$ (inside $\mathcal{E} = \mathcal{F}/\mathcal{G}$), then $\theta(\beta(s)) = 0$ in $\Gamma(\varphi^*(U), \varphi^*(\mathcal{E}))$, since $0^{\text{ext}} = 0$. But then $\theta(s) \in \Gamma(\varphi^*(U), \varphi^*(\mathcal{F}))$ maps to 0 in $\Gamma(\varphi^*(U), \varphi^*(\mathcal{E}))$ and belongs to $\Gamma(\varphi^*(U), \mathcal{G}^{\text{ext}})$. $\square$

We describe the containment in the closure submodule more explicitly.

Lemma 1.4.5. *Let $\varphi : Y \rightarrow X$ be a morphism between two (pre-)ringed sites and suppose that it fulfills the cofiltered condition of 1.3.1. Let $\mathcal{G} \subseteq \mathcal{F}$ be $\mathcal{O}_X$ -(pre-) modules. Let $s \in \Gamma(U, \mathcal{F})$, $U \in \mathcal{T}_X$. Set $V = \varphi^*(U)$. Then $s \in \Gamma(U, \mathcal{G}^c)$ if and only if there exists a covering $V_i \xrightarrow{\tau_i} V$, $i \in I$, and morphisms $\rho_i : V_i \rightarrow \varphi^*(U_i)$ in $\mathcal{T}_Y$ and $\sigma_i : U_i \rightarrow U$ in $\mathcal{T}_X$ such that $\tau_i = \varphi^*(\sigma_i) \circ \rho_i$ and sections $t_i \in \Gamma(U_i, \mathcal{G}) \otimes_{\Gamma(U_i, \mathcal{O}_X)} \Gamma(V_i, \mathcal{O}_Y)$ such that $\rho_{U_i}^U(s) \otimes 1 = \alpha(t_i)$ in $\Gamma(U_i, \mathcal{F}) \otimes_{\Gamma(U_i, \mathcal{O}_X)} \Gamma(V_i, \mathcal{O}_Y)$, where α denotes the inclusion $\alpha : \mathcal{G} \rightarrow \mathcal{F}$.*

Proof. The element $s \in \Gamma(U, \mathcal{F})$ defines the element $\theta(s) \in \Gamma(V, \varphi^*(\mathcal{F}))$ as the image of $s \otimes 1$ under

$$\Gamma(U, \mathcal{F}) \otimes_{\Gamma(U, \mathcal{O}_X)} \Gamma(V, \mathcal{O}_Y) \rightarrow \Gamma(V, \varphi^{-1}(\mathcal{F})) \otimes_{\Gamma(V, \varphi^{-1}(\mathcal{O}_X))} \Gamma(V, \mathcal{O}_Y) \rightarrow \Gamma(V, \varphi^*(\mathcal{F})).$$

This element is in the image sheaf of $\varphi^*(\mathcal{G}) \rightarrow \varphi^*(\mathcal{F})$ if and only if there exists a covering $V_i \rightarrow V$, $i \in I$, and sections $r_i \in \Gamma(V_i, \varphi^*(\mathcal{G}))$ mapping to $\rho_{V_i}^V(\theta(s))$. Each r_i is represented by a covering $W_j \rightarrow V_i$, $j \in J_i$, and elements $\tilde{r}_j \in \Gamma(W_j, \varphi^{\text{pre}}(\mathcal{G})) = \Gamma(W_j, \varphi^{-1}(\mathcal{G})) \otimes_{\Gamma(W_j, \varphi^{-1}(\mathcal{O}_X))} \Gamma(W_j, \mathcal{O}_Y)$ (fulfilling some compatible condition, which we do not need). We denote the composed covering of V again by V_i . By the definition of inverse presheaf, the sections $r_i \in \Gamma(V_i, \varphi^{-1}(\mathcal{G})) \otimes_{\Gamma(V_i, \varphi^{-1}(\mathcal{O}_X))} \Gamma(V_i, \mathcal{O}_Y)$ are represented by $t_i \in \Gamma(U_i, \mathcal{G}) \otimes_{\Gamma(U_i, \mathcal{O}_X)} \Gamma(V_i, \mathcal{O}_Y)$, where $\rho_i : V_i \rightarrow \varphi^*(U_i)$ are morphisms in $\mathcal{T}_Y$ and $U_i \in \mathcal{T}_X$. These elements define sections $\alpha(t_i) \in \Gamma(U_i, \mathcal{F}) \otimes_{\Gamma(U_i, \mathcal{O}_X)} \Gamma(V_i, \mathcal{O}_Y)$, which map by assumption to $\rho_{V_i}^V(\theta(s)) \in \Gamma(V_i, \varphi^*(\mathcal{F}))$.

We may replace U_i by $U_i \times_X U$ (denoted again by U_i) so that we have the projections to U , the morphisms commute in the described way and $s_i = \rho_{U_i}^U(s) \in \Gamma(U_i, \mathcal{F})$ makes sense. So we have elements $\alpha(t_i)$ and $s_i \otimes 1$ in $\Gamma(U_i, \mathcal{F}) \otimes_{\Gamma(U_i, \mathcal{O}_X)} \Gamma(V_i, \mathcal{O}_Y)$ and we know that they define the same element in the sheafification $\Gamma(V_i, \varphi^*(\mathcal{F}))$. This means that there exist coverings $W_j \rightarrow V_i$, $j \in J_i$, such that their difference is 0 in $\Gamma(W_j, \varphi^{\text{pre}}(\mathcal{F}))$. This means by the cofiltered condition that there exist morphisms $W_j \rightarrow \varphi^*(X_j)$, $X_j \in \mathcal{T}_X$, such that the restriction of $s_i \otimes 1$ in $\Gamma(U_i \times_X X_j, \mathcal{F}) \otimes_{\Gamma(U_i \times_X X_j, \mathcal{O}_X)} \Gamma(W_j, \mathcal{O}_Y)$ equals the restriction of $\alpha(t_i)$. So the covering $W_j \rightarrow V$, $j \in \bigcup_i J_i$, has all the desired properties. $\square$

Example 1.4.6. Let X denote an algebraic variety over the complex numbers $\mathbb{C}$ and let $X(\mathbb{C})$ denote the corresponding complex space with the sheaf of holomorphic functions. Then we have a morphism of ringed spaces $X(\mathbb{C}) \rightarrow X$, which is in particular a site morphism. Extension and contraction yields the same ideal back, since the algebraic stalks and the analytic stalks at a closed point have the same completion, which is faithfully flat. So if $f \in I^{\text{an}}$, then this holds in every analytic stalk, in their completions and therefore already in the algebraic setting.

If we consider however $X(\mathbb{C})$ with the sheaf of continuous functions, denoted by X_{cont} , then extension and contraction may give back a larger ideal I^{cont} , which we call the *continuous closure*. For example, in $\mathbb{C}[z, w]$ we have $z^2 w^2 \notin (z^3, w^3)$, but $z^2 w^2 \in (z^3, w^3)^{\text{cont}}$.

We deal with this continuous closure in [7], where we give a precise description of it for monomial ideals.

1.5. Grothendieck topologies on a scheme.

Let X denote a scheme, and let Sch_X denote the category of schemes over X , where the morphisms are supposed to be compatible with the structure morphisms. A Grothendieck topology on a subcategory $\mathcal{T}_X \subseteq Sch_X$ containing X and closed under products is called a Grothendieck topology on X . We will denote the scheme X endowed with such a topology by X_{top} , where top is a name for the corresponding topology. Note that the products appearing in Definition 1.1.1 are now products of schemes. The site $(X_{\text{top}}, \mathcal{T}_X)$ is immediately a pretopos by setting $\Gamma(U, \mathcal{O}^{\text{pre}}) := \Gamma(U, \mathcal{O}_U)$ for $U \in \mathcal{T}_X$, and sheafification in the topology gives a ringed site $(X_{\text{top}}, \mathcal{T}_X, \mathcal{O}_{\text{top}})$.

A site morphism between schemes Y_{top} and X_{top} is a scheme morphism $\varphi : Y \rightarrow X$ such that $U \in \mathcal{T}_X$ implies that $\varphi^*(U) = Y \times_X U \in \mathcal{T}_Y$ and such that a covering $U_i \rightarrow U$, $i \in I$, yields a covering $Y \times_X U_i \rightarrow Y \times_X U$, $i \in I$. It is then also a morphism of (pre)ringed sites by taking $\Gamma(U, \mathcal{O}^{\text{pre}}) = \Gamma(U, \mathcal{O}_U) \rightarrow \Gamma(\varphi^*(U), \mathcal{O}_{\varphi^*(U)}) = \Gamma(\varphi^*(U), \mathcal{O}^{\text{pre}})$ and its sheafification. A scheme morphism may or may not induce a site morphism. For our treatment we will also adopt the following convention.

Convention 1.5.1. In saying that a scheme X has a Grothendieck topology $\mathcal{T}_X$ we mean that there exists a basic (or standard) topology $\mathcal{B}_X$ on X which is always inside the Zariski topology (it will be usually either the trivial topology (Example 1.5.2) or the Zariski topology) and which we will not mention in general, of which the given Grothendieck topology is a refinement. In particular we have a site morphism $\varphi : X_{\text{top}} \rightarrow X$. We assume also that if $j : V \rightarrow X$ belongs to $\mathcal{T}_X$ and factors through an open subscheme $V \rightarrow U \subseteq X$, then also $V \rightarrow U \in \mathcal{T}_X$. We denote the $\mathcal{O}_{\text{top}}$ -sheafification of an $\mathcal{O}_X$ -module M by $M_{\text{top}} = \varphi^*(M)$.

Example 1.5.2. Let R be a commutative ring and consider the category consisting of one object X ('Spec R ') and one identical morphism. We define a sheaf of rings by $\Gamma(X, \mathcal{O}_X) = R$ and call this the *trivial site* for R . An R -module M yields immediately an $\mathcal{O}_X$ -module, which we also denote by M .

Let $\mathcal{T}_R$ be a subcategory of R -algebras (or rather its opposite category, so we will write $\text{Spec } S$ instead of S) containing R , closed under tensor products and endowed with a certain Grothendieck topology. Then $\mathcal{T}_R$ is immediately a pretopos by setting $\Gamma(\text{Spec } S, \mathcal{O}^{\text{pre}}) = S$, and we get a morphism of pretopos by the inclusion $\varphi^* : \{X\} \rightarrow \mathcal{T}_R$ (with $\varphi^*(X) = X$). The ring homomorphism is given by the identity $\Gamma(X, \mathcal{O}_X) = R \rightarrow \Gamma(\varphi^*(X), \mathcal{O}^{\text{pre}}) = R$. The inverse image presheaf of an $\mathcal{O}_X$ -module M is given by the constant presheaf since

$$\Gamma(\text{Spec } S, \varphi^{-1}(M)) = \varinjlim_{(U, \rho) \in I_S^{\varphi}} \Gamma(U, M) = \varinjlim_{\rho : R \rightarrow S} \Gamma(X, M) = M.$$

Here the (only) indexing morphism $R \rightarrow S$ is the structure homomorphism of the R -algebra S . The $\mathcal{O}_X$ -module M yields the corresponding $\mathcal{O}^{\text{pre}}$ -premodule $\varphi^{\text{pre}}(M)$ by sending $\text{Spec } S \mapsto M \otimes_R S$. The sheafification M_{top} (and also $\mathcal{O}_{\text{top}}$) might be difficult to compute, depending on the topology.

Example 1.5.3. Let $X = \text{Spec } R$ be an affine scheme with the trivial topology and let X_{Zar} be the affine scheme with the Zariski topology and with the structure sheaf $\mathcal{O}_{\text{Zar}}$,

so that $\varphi : X_{\text{Zar}} \rightarrow X_{\text{triv}}$ is a morphism of ringed sites. If M is an R -module, then the inverse image presheaf $\varphi^{-1}(M)$ is the constant presheaf $V \mapsto M$. The associated $\mathcal{O}_{\text{Zar}}$ -premodule is given by $V \mapsto M \otimes_R \Gamma(V, \mathcal{O}_{\text{Zar}})$. The pull-back of M is the sheaf associated to this presheaf and gives the quasicoherent module $\tilde{M}$ [27, Proposition II.5.1].

Definition 1.5.4. Let $\mathcal{T}_X$ denote a Grothendieck topology on a scheme X .

We call $\mathcal{T}_X$ or X_{top} *affine*, if all structural morphisms $U \rightarrow X$ in $\mathcal{T}_X$ are affine (then all morphisms are affine).

We call it *of finite type* if all morphisms $U \rightarrow V$ in $\mathcal{T}_X$ are of finite type.

We call it *quasicompact* if for every covering $U_i \rightarrow U$, $i \in I$, there exists a finite subset $J \subseteq I$ such that $U_i \rightarrow U$, $i \in J$, is also a covering.

We call it *single-handed* if it is of finite type and if every covering is given by a single morphism $U \rightarrow V$.

We call it *covering* if every morphism in $\mathcal{T}_X$ is a cover.

Proposition 1.5.5. *Let X be a scheme and let $\mathcal{T}_X$ be a Grothendieck topology on X (recall Convention 1.5.1). Let $\varphi : X_{\text{top}} \rightarrow X$ be the corresponding site morphism. Let $j : V \rightarrow X$ be in $\mathcal{T}_X$, considered as a Zariski morphism, and let M be an $\mathcal{O}_X$ -module. Then the following hold.*

- (i) *We have $\Gamma(V, \varphi^{-1}(M)) = \Gamma(V, j^{-1}(M))$.*
- (ii) *We have $\Gamma(V, \varphi^{\text{pre}}(M)) = \Gamma(V, j^{\text{pre}}(M))$.*
- (iii) *The sheafification $\varphi^*(M) = M_{\text{top}}$ equals the $\mathcal{T}_X$ -sheafification of the $\mathcal{O}^{\text{pre}}$ -premodule $V \mapsto \Gamma(V, j^{\text{pre}}(M))$.*
- (iv) *If $\mathcal{T}_X$ is a refinement of the Zariski topology or an affine topology on an affine scheme, then M_{top} equals also the sheafification of $V \mapsto \Gamma(V, j^*(M))$, where j^* is the pull-back of modules in the Zariski topology.*

Proof. (i). We have

$$\Gamma(V, \varphi^{-1}(M)) = \varinjlim_{(U, \rho) \in I_V^\varphi} \Gamma(U, M) = \varinjlim_{j(V) \subseteq U} \Gamma(U, M) = \Gamma(V, j^{-1}(M)),$$

because there exists by our convention a morphism $V \rightarrow U = \varphi^*(U)$ in $\mathcal{T}_X$ if and only if $j(V) \subseteq U$.

(ii). We have by Definition 1.3.5 and part (i)

$$\begin{aligned} \Gamma(V, \varphi^{\text{pre}}(M)) &= \Gamma(V, \varphi^{-1}(M)) \otimes_{\Gamma(V, \varphi^{-1}(\mathcal{O}_{\text{Zar}}))} \Gamma(V, \mathcal{O}^{\text{pre}}) \\ &= \Gamma(V, j^{-1}(M)) \otimes_{\Gamma(V, j^{-1}(\mathcal{O}_{\text{Zar}}))} \Gamma(V, \mathcal{O}_V) = \Gamma(V, j^{\text{pre}}(M)). \end{aligned}$$

(iii) follows from (ii). (iv) follows also from (ii), since in the first case $V \mapsto \Gamma(V, j^*(M))$ is the Zariski sheafification of the presheaf $V \mapsto \Gamma(V, \varphi^{-1}(M)) = \Gamma(V, j^{-1}(M))$, and the topology is a refinement of the Zariski topology. In the second case $\Gamma(V, j^*(M)) = \Gamma(V, j^{\text{pre}}(M))$ anyway. $\square$

Corollary 1.5.6. *Let X denote a scheme endowed with a Grothendieck topology $\mathcal{T}_X$, and suppose that it is either a refinement of the Zariski topology or affine over an affine scheme. Let $j : Y \rightarrow X$ be in $\mathcal{T}_X$ and let M be an $\mathcal{O}_X$ -module on X . Let M_{top} be its sheafification in X_{top} and let $j^*(M)$ be the Zariski pull-back to Y . Let $\mathcal{T}_Y$ be the category consisting of all morphisms $Z \rightarrow Y$ in $\mathcal{T}_X$ as objects and the morphisms in $\mathcal{T}_X$ over Y as morphisms (and the induced coverings). Then the restriction sheaf $(M_{\text{top}})|_{\mathcal{T}_Y}$ given by $(Z \rightarrow Y) \mapsto \Gamma(Z, M_{\text{top}})$ equals the sheafification $(j^*(M))_{\text{top}}$ in $\mathcal{T}_Y$.*

Proof. We have to show for $\rho : Z \rightarrow Y$ (in $\mathcal{T}_Y$) that $\Gamma(Z, (j^*(M))_{\text{top}}) = \Gamma(Z, M_{\text{top}})$. By Proposition 1.5.5(iv) both sheaves come from the presheaf $(\rho : Z \rightarrow Y) \mapsto \rho^*(j^*(M))$, one time sheafified in $\mathcal{T}_X$ and one time sheafified in $\mathcal{T}_Y$. The objects $Z \rightarrow Y$ have the same coverings in $\mathcal{T}_X$ and in $\mathcal{T}_Y$, because a covering $Z_i \rightarrow Z$ over X can also be considered as a covering with basis Y via $Z \rightarrow Y$. Hence the associated separated presheaves are the same and then also their sheafifications. $\square$

Example 1.5.7. Proposition 1.5.5 and Corollary 1.5.6 are not true without Convention 1.5.1 on Zariski factorizations. To give an example, let R be a normal domain of dimension ≥ 2 . Let $X = \text{Spec } R$ have the trivial topology and let $\mathcal{T}_X$ be given by the Zariski open subsets which contain all points of codimension one (and Zariski coverings). Then $\Gamma(V, \mathcal{O}_X) = R$ for all $V \in \mathcal{T}_X$, and X_{top} is a ringed site with constant structure sheaf. For an R -module M the inverse image presheaf under $\varphi : X_{\text{top}} \rightarrow X$ is constant, and so also $\varphi^{\text{pre}}(M)$ is constant $= M$. Let now $Y \in \mathcal{T}_X$ and M be an R -module such that $\Gamma(Y, \tilde{M}) \neq M$, e.g. take $j : Y = X - \{P\} \rightarrow X$ and $M = \kappa(P)$, where $P \in X$ is a closed point. Then $j^*(M) = 0$ and so $(j^*(M))_{\text{top}} = 0$, but $\Gamma(Y, M_{\text{top}}) = M$.

To a large extent we will restrict to affine, single-handed Grothendieck topologies on an affine scheme $X = \text{Spec } R$, so that $\mathcal{T}_R = \mathcal{T}_X$ is given by a category of R -algebras and the coverings are given by a certain class of R -algebra homomorphisms.

Remark 1.5.8. We will often consider the Zariski topology on a scheme X as an (affine) single-handed Grothendieck topology, called the *(affine) single-handed Zariski topology*, where $\mathcal{T}_X$ consists of all finite disjoint unions of (affine) open immersions $U \rightarrow X$ and where the only coverings are given by $\bigsqcup U_i \rightarrow U'$, where the U_i do cover U' as a set. If $Y = \bigsqcup_{i \in I} U_i \rightarrow X$ is a Zariski covering and $\mathcal{F}$ is a quasicohherent sheaf on X , then

$$\begin{aligned} \Gamma(Y \times_X Y, \mathcal{F}) &= \Gamma\left(\left(\bigsqcup_{i \in I} U_i\right) \times_X \left(\bigsqcup_{i \in I} U_i\right), \mathcal{F}\right) \\ &= \Gamma\left(\bigsqcup_{(i,j) \in I \times I} U_i \times_X U_j, \mathcal{F}\right) \\ &= \bigoplus_{(i,j) \in I \times I} \Gamma(U_i \times_X U_j, \mathcal{F}) = \bigoplus_{(i,j) \in I \times I} \Gamma(U_i \cap U_j, \mathcal{F}). \end{aligned}$$

This holds more generally for every sheaf which respects disjoint Zariski unions (as in Lemma 1.5.10) and it holds also for products with more than two factors. From this it follows that the Čech cohomology for such sheaves is the same in the Zariski site and in the single-handed Zariski site.

Although we will mainly deal with single-handed Grothendieck topologies it is good to know when they respect Zariski covers.

Definition 1.5.9. Let X be a scheme with a Grothendieck topology $\mathcal{T}_X$. We say that the Grothendieck topology *respects (disjoint) Zariski coverings* if a family of morphisms $U_i \rightarrow U$, $i \in I$, (inside $\mathcal{T}_X$) is a covering if and only if for an arbitrary (disjoint) Zariski cover $U = \bigsqcup_{j \in J} V_j$ ($U = \bigsqcup_{j \in J} V_j$) the family $U_i \times_V V_j \rightarrow V_j$, $i \in I$, is a covering of V_j for every j .

Lemma 1.5.10. Let $X = \bigsqcup_{i \in I} U_i$ be a finite disjoint union of open subschemes of a scheme X . Let an affine single-handed Grothendieck topology X_{top} be given which is a refinement of the single-handed Zariski topology and which respects disjoint Zariski coverings. Let M

be an $\mathcal{O}_X$ -module on X . Then $\Gamma(X_{\text{top}}, M_{\text{top}}) = \bigoplus_{i \in I} \Gamma(U_i, M_{\text{top}}) = \bigoplus_{i \in I} \Gamma(U_i, (\varphi_i^* M)_{\text{top}})$ ($\varphi_i : U_i \rightarrow X$).

Proof. The second equation follows from Corollary 1.5.6. The morphisms $\varphi_i : U_i \rightarrow X$ induce the restrictions $\Gamma(X_{\text{top}}, M_{\text{top}}) \rightarrow \Gamma(U_i, M_{\text{top}})$. An element $s \in \Gamma(X_{\text{top}}, M_{\text{top}})$ is represented by $t \in \Gamma(Y, \psi^*(M))$ (Proposition 1.5.5(iv)), where $\psi : Y \rightarrow X$ is a cover in the topology. Suppose that s restricts to 0 in $\Gamma(U_i, M_{\text{top}})$ for all open subsets U_i . This means that $\theta(t) \in \Gamma(Y, M_{\text{top}})$ restricts to 0 in $\Gamma(U_i \times_X Y, M_{\text{top}})$ and so there exist covers $Z_i \rightarrow U_i \times_X Y$ such that the pull-backs of t are 0. These covers glue together to the cover $\biguplus_{i \in I} Z_i \rightarrow Y$ (since being a cover can be tested on the disjoint subsets of the union), hence also $\theta(t) = 0$ in $\Gamma(Y, M_{\text{top}})$ and so $s = 0$ in $\Gamma(X, M_{\text{top}})$. On the other hand, let $s_i \in \Gamma(U_i, M_{\text{top}})$ be given. Say s_i is represented by compatible elements $t_i \in \Gamma(Z_i, \psi_i^*(M))$, where $\psi_i : Z_i \rightarrow U_i$ is a cover for each $i \in I$. Then again these covers glue together to a cover of X , and the tuple (t_i) is also compatible over X , so it defines a global section. $\square$

1.6. Rings and modules of global sections.

In the flat (or the étale) Grothendieck topology on an affine scheme $\text{Spec } R$ we have $\Gamma(\text{Spec } R, \mathcal{O}_{\text{top}}) = R$. This does not hold for non-pure topologies in general. We have to compute in a given Grothendieck topology $\mathcal{T}_X$ the sheafification $U \mapsto \Gamma(U, \mathcal{O}_{\text{top}})$, $U \in \mathcal{T}_X$, associated to the presheaf $U \mapsto \Gamma(U, \mathcal{O}_U)$, and this computation is a non-trivial matter.

Proposition 1.6.1. *Suppose that we have a Grothendieck topology on a certain subcategory G of schemes, meaning that for every $X \in G$ we have a Grothendieck topology $\mathcal{T}_X$ and every scheme morphism $Y \rightarrow X$ in G induces a site morphism $Y_{\text{top}} \rightarrow X_{\text{top}}$. Then we get a functor $X \mapsto \Gamma(X, \mathcal{O}_{\text{top}})$, $X \in G$, together with a natural transformation $\theta : \Gamma(X, \mathcal{O}_X) \rightarrow \Gamma(X, \mathcal{O}_{\text{top}})$.*

Proof. This is clear. $\square$

Remark 1.6.2. In this way we will encounter the perfect closure (Theorem 4.3.3), the semi-normalization (Proposition 5.2.3), the ring of constructible sections (Theorem 3.8.2), the completion of a local ring (Proposition 7.1.1) and more. In particular, we have $\Gamma((\text{Spec } R)_{\text{top}}, \mathcal{O}_{\text{top}}) \neq R$. From this it follows that the additive group scheme $\mathbb{A}^1$ does not define a sheaf in the topology, because $\text{Mor}(X, \mathbb{A}^1) \cong \Gamma(X, \mathcal{O}_X) \neq \Gamma(X_{\text{top}}, \mathcal{O}_{\text{top}})$ [27, Exercise 2.4]. Hence such topologies are not *subcanonical*, i.e. not every representable functor is a sheaf (see [66, Definition 2.54] or [24, Définition II.2.5]).

A global section $f \in \Gamma(X, \mathcal{O}_{\text{top}})$ is represented by a compatible family $f_i \in \Gamma(U_i, \mathcal{O}_{U_i})$, $i \in I$, where $U_i \rightarrow X$, $i \in I$, is a covering in the Grothendieck topology, and where the compatibility condition means that $p_1^*(f_i) - p_2^*(f_j) \in \Gamma(U_i \times_X U_j)$ has the property that it vanishes on a covering of $U_i \times_X U_j$. A similar characterization holds for an $\mathcal{O}_X$ -module M . Quite often we will deal with a single cover $Y \rightarrow X$, and a global section is then given by $f \in \Gamma(Y, \mathcal{O}_Y)$ such that $p_1^*(f) - p_2^*(f) \in \Gamma(Y \times_X Y, \mathcal{O})$ is zero in a covering of $Y \times_X Y$.

Lemma 1.6.3. *Suppose that a Grothendieck topology on a scheme X has the property that every covering $\varphi_i : U_i \rightarrow X$, $i \in I$, has a section $\psi_i : X \rightarrow U_i$ for some i (M. Walker calls such X acyclic; cf. [69, Lemma 3.7]). Then for an $\mathcal{O}_X$ -module M the natural homomorphism $\theta : \Gamma(X, M) \rightarrow \Gamma(X, M_{\text{top}})$ is an isomorphism.*

Proof. Let $s \in \Gamma(X, M)$. If $s = 0$ in $\Gamma(X, M_{\text{top}})$, then there exists a covering $\varphi_i : U_i \rightarrow X$, $i \in I$, such that $\varphi_i^*(s) = 0$ in $\Gamma(U_i, \varphi_i^*(M))$ for all $i \in I$. The factorization $X \rightarrow U_i \rightarrow X$ shows that $s = 0$.

On the other hand, let $s \in \Gamma(X, M)$ be represented by $s_i \in \Gamma(U_i, \varphi_i^*(M))$ for a covering $U_i \rightarrow X$, $i \in I$. Then via $X \rightarrow U_i \rightarrow X$ the element s_i gives an element $\psi_i^*(s_i) \in \Gamma(X, M)$, and since $\text{id} : X \rightarrow X$ is a covering, we may represent s also by $\psi_i^*(s_i)$. $\square$

Lemma 1.6.4. *Let $Y \rightarrow X$ be a cover in a Grothendieck topology on X , and let $f \in \Gamma(Y, \mathcal{O}_Y)$ be a compatible element, defining a global element in $\Gamma(X, \mathcal{O}_{\text{top}})$. If $\alpha : Y \rightarrow Y$ is an X -automorphism, then $\alpha^*(f)$ is also compatible and it defines the same global element.*

Proof. It is clear that $\alpha^*(f)$ is also compatible, since an automorphism preserves the compatibility conditions. Suppose that the pull-backs of $p_1^*(f)$ and $p_2^*(f)$ are the same under the covering $Z_i \rightarrow Y \times_X Y$, $i \in I$. The automorphism $\text{id} \times \alpha : Y \times_X Y \rightarrow Y \times_X Y$ shows that also $(\text{id} \times \alpha)^*(p_1^*(f)) = p_1^*(f)$ and $(\text{id} \times \alpha)^*(p_2^*(f)) = p_2^*(\alpha^*(f))$ are the same in some covering. Therefore also $p_2^*(f)$ and $p_2^*(\alpha^*(f))$ are the same in some covering. Hence f and $\alpha^*(f)$ are the same in some covering and so they define the same element in $\Gamma(X, \mathcal{O}_{\text{top}})$. $\square$

If a Grothendieck topology $\mathcal{T}_X$ on $X = \text{Spec } R$ has the property that the reduction defines a cover, then it is important to know for an element $f \in S$, $R \rightarrow S$ a cover, whether $f \otimes 1 - 1 \otimes f \in S \otimes_R S$ is nilpotent. In this case f is compatible and defines a global element in $\Gamma(R_{\text{top}}, \mathcal{O}_{\text{top}})$.

Lemma 1.6.5. *Let S denote an R -algebra, $f \in S$. Then $f \otimes 1 - 1 \otimes f \in S \otimes_R S$ is nilpotent if and only if the following hold: For all homomorphisms $\varphi : R \rightarrow K$, K a field, and any two extensions $\psi_1, \psi_2 : S \rightarrow K$ of φ , we have $\psi_1(f) = \psi_2(f)$.*

Proof. The element $\delta(f) = f \otimes 1 - 1 \otimes f \in S \otimes_R S$ is nilpotent if and only if for all ring homomorphisms $S \otimes_R S \rightarrow K$, K a field, the image of $\delta(f)$ is 0. These homomorphisms are in one to one correspondence with the data described in the lemma, so the result follows. $\square$

Lemma 1.6.6. *Let a Grothendieck topology be given on a subcategory G of schemes such that every morphism in G induces a site morphism. Suppose that for $X \in G$ all $U \rightarrow X$ (in $\mathcal{T}_X$) and all morphisms $\text{Spec } \kappa(x) = x \rightarrow X$, $x \in X$ a point, belong to G . Suppose also that $\Gamma(x, \mathcal{O}_{\text{top}}) = \kappa(x)$ (or $\kappa(x)^\infty$) for every point x in G . Then a compatible element $f \in \Gamma(Y, \mathcal{O}_Y)$, $Y \rightarrow X$ a cover, has the property that for two points $y_1, y_2 \in Y$ over $x \in X$ the values of f in $\kappa(y_1), \kappa(y_2) \supseteq \kappa(x)$ lie in $\kappa(x)$ (or in their perfect closures) and are identical.*

Proof. We have the commutative diagram

$$\begin{array}{ccc} \Gamma(X, \mathcal{O}_{\text{top}}) & \xrightarrow{\quad} & \Gamma(Y, \mathcal{O}_{\text{top}}) \\ \downarrow & & \downarrow \\ \Gamma(x, \mathcal{O}_{\text{top}}) = \kappa(x) & \longrightarrow & \Gamma(y_1, \mathcal{O}_{\text{top}}) = \kappa(y_1), \Gamma(y_2, \mathcal{O}_{\text{top}}) = \kappa(y_2). \end{array}$$

In this diagram the element $f \in \Gamma(Y, \mathcal{O}_Y) \rightarrow \Gamma(Y, \mathcal{O}_{\text{top}})$ comes from an element $\tilde{f} \in \Gamma(X, \mathcal{O}_{\text{top}})$, since $f \in \Gamma(Y, \mathcal{O}_Y)$ is compatible over X (both rings of global sections are colimits of compatible elements over coverings). Hence the image of f in $\kappa(y_1)$ and $\kappa(y_2)$

comes from $\tilde{f}$ via $\kappa(x)$. The same argument works if the fields are replaced by their perfect closures. $\square$

The next lemma describes a situation where a lot of “new” global elements arise in the Grothendieck topology. We will use this argument in Corollary 3.6.6, Theorem 4.3.3, Proposition 6.1.6 and Proposition 7.3.8.

Lemma 1.6.7. *Suppose that $Y \rightarrow X$ is a cover in a Grothendieck topology on X such that the diagonal $\Delta : Y \rightarrow Y \times_X Y$ is a cover. Let $\mathcal{F}$ denote a sheaf of abelian groups on X_{top} . Then the restriction homomorphism $\Gamma(X, \mathcal{F}) \rightarrow \Gamma(Y, \mathcal{F})$ is an isomorphism. This holds in particular if one (then both) projection $Y \times_X Y \rightarrow Y$ is an isomorphism.*

Proof. We look at the exact complex

$$\Gamma(X, \mathcal{F}) \longrightarrow \Gamma(Y, \mathcal{F}) \xrightarrow{\rho_1 - \rho_2} \Gamma(Y \times_X Y, \mathcal{F})$$

and have to show that every element $s \in \Gamma(Y, \mathcal{F})$ maps to 0 (the first mapping is injective anyway). The restriction homomorphism $\rho(\Delta) : \Gamma(Y \times_X Y, \mathcal{F}) \rightarrow \Gamma(Y, \mathcal{F})$ comes from a cover and $\rho(\Delta)(\rho_1(s) - \rho_2(s)) = 0$, so $\rho_1(s) - \rho_2(s) = 0$ as well.

If one projection is an isomorphism, then also the diagonal $\Delta : Y \rightarrow Y \times_X Y$ is an isomorphism, hence a cover. Then both projections equal Δ^{-1} . $\square$

1.7. Filters and stalks.

In the étale topology, the *Henselization* of the local ring $\mathcal{O}_x$ of a point $x \in X$ with separably closed residue class field can be recovered by taken a suitable colimit over all *étale neighborhoods* of that point ([24, Définition VIII.4.3], [20, Définition 18.6.5], [49, Remark I.4.11]). We will describe here how to get suitable colimits over reasonably filtered systems in $\mathcal{T}_X$ and in which sense it is possible to bring all coverings together into a single absolute cover. Motivating examples for this are that all finite extensions of a domain are inside its absolute integral closure [36], or all Frobenius powers are inside the perfect closure.

Definition 1.7.1. Let Λ denote a small quasicolimated category (Definition 1.2.1) with a final object and let $\mathcal{T}_X$ be a Grothendieck topology on a scheme X . Then we call a covariant functor $F : \Lambda \rightarrow \mathcal{C}$ which respects the final object a *quasifilter* in $\mathcal{T}_X$. If moreover Λ is cofiltered, then we call F a *filter*. We write often U_λ or $U_\lambda \rightarrow X$ instead of $F(\lambda)$.

Suppose that $\mathcal{P}$ is a presheaf on a Grothendieck topology $\mathcal{T}_X$ with values in a category $\mathcal{A}$ and that $F : \Lambda \rightarrow \mathcal{T}_X$ is a quasifilter. Then we get also a contravariant (quasifiltered) diagram in $\mathcal{A}$ by looking at

$$\mathcal{P} \circ F : \Lambda \longrightarrow \mathcal{A}, \lambda \longmapsto \Gamma(U_\lambda, \mathcal{P}).$$

If $\rho : \lambda \rightarrow \mu$ belongs to Λ , then we will denote the induced restriction morphism $\Gamma(U_\mu, \mathcal{P}) \rightarrow \Gamma(U_\lambda, \mathcal{P})$ also by ρ or ρ_λ^μ . If $\mathcal{A}$ is such that colimits over (certain) diagrams exist, then we denote the colimit by

$$\mathcal{P}_F := \varinjlim_{\lambda \in \Lambda} \Gamma(U_\lambda, \mathcal{P})$$

and call it the *stalk* of $\mathcal{P}$ in F .

Construction 1.7.2. Let $\mathcal{T}_X$ be a Grothendieck topology on X and fix a scheme morphism $Z \rightarrow X$ (we call this a *fixation*). This defines a quasifilter in the following way: Let the indexing category $\Lambda = \Lambda_Z = \Lambda_{Z \rightarrow X}$ be given by (isomorphism types of) all commutative diagrams

$$\begin{array}{ccc} & U & \\ \nearrow & \downarrow & \\ Z & \longrightarrow & X \end{array}$$

as objects, where $U \rightarrow X$ is in $\mathcal{T}_X$ (so the objects are the Z -pointed open sets in $\mathcal{T}_X$), and by all the morphisms $U \rightarrow U'$ in $\mathcal{T}_X$ which commute with the fixations $Z \rightarrow U$ and $Z \rightarrow U'$. The final object is given by $Z \rightarrow X \xrightarrow{\text{id}} X$. If two diagrams $Z \rightarrow U \rightarrow X$ and $Z \rightarrow U' \rightarrow X$ are given over $Z \rightarrow U_0 \rightarrow X$ (that is, $U, U' \rightarrow U_0$ fixing Z), then they come together in $Z \rightarrow U \times_{U_0} U' \rightarrow X$ (arbitrary products exist in $\mathcal{T}_X$). Hence Λ is a quasicofiltered category. The corresponding quasifilter is given by sending

$$\Lambda \longrightarrow \mathcal{T}_X, (Z \rightarrow U \rightarrow X) \longmapsto (U \rightarrow X),$$

i.e. by forgetting $Z \rightarrow U$.

If $\mathcal{T}_X$ is the full category of finite type over X (a big site) and the topology is affine, then an affine fixation $Z = \text{Spec } H$ yields even a filter. If $\psi_1, \psi_2 : U \rightrightarrows U'$ are given corresponding to $\psi_1, \psi_2 : B' \rightrightarrows B$, then $B \otimes_{B'} B / (\psi_1(c) \otimes 1 - 1 \otimes \psi_2(c), c \in B')$ gives the equalizer which belongs then to the category.

Example 1.7.3. An important case is when $Z = \text{Spec } K$ is the affine spectrum of an algebraically closed field K , in which case $Z \rightarrow X$ is called a *geometric point*. Goodwillie and Lichtenbaum call in [17, Section 2] a scheme P a *point* in a Grothendieck topology if for every covering $U_i \rightarrow U$, $i \in I$, and every morphism $P \rightarrow U$ there exists a lifting $P \rightarrow U_i$ for some i . If the topology is such that all coverings are surjective and of finite type, then every geometric point is also a point in this sense.

Example 1.7.4. In a Grothendieck topology the quasifilter given by a fixation is in general not a filter. Consider the category of finite non-empty sets where only surjective morphisms are allowed and are covers. This is the same as the affine single-handed covering Zariski topology on $X = \text{Spec } K$, K a field. The identity $X \rightarrow X$ gives a fixation and an (absolute) quasifilter, the objects in this quasifilter are the pointed sets. Let W consist of the two points P_0 (fixed) and P_1 and let V consist of three points Q_0 (fixed), Q_1 and Q_2 . Let ψ_1 and ψ_2 be given by $\psi_1(Q_0) = \psi_2(Q_0) = P_0$, $\psi_1(Q_1) = \psi_2(Q_1) = P_1$, and $\psi_1(Q_2) = P_0$, $\psi_2(Q_2) = P_1$. Then there does not exist a surjective map $U \rightarrow V$ which unifies the two morphisms.

Consider the sheaf $V \rightarrow K^V = \Gamma(V, \mathcal{O}_{\text{top}})$ and the diagram induced by the given quasifilter. The colimit of this system as a diagram of sets is $(\bigoplus_V K^V) / \sim$, where $\sim$ is the equivalence relation in which $f \in K^V$ and $g \in K^W$ are equivalent if and only if there exists a set U and surjective mappings $\varphi : U \rightarrow V$, $\psi : U \rightarrow W$ with $\varphi^*(f) = \psi^*(g)$. This is true if and only if f and g have the same values. Therefore the equivalence classes are given by finite unordered tuples of different elements in K .

This is however not a group, since it is not possible to define a group operation. Two elements $f \in K^V$ and $g \in K^W$ come together in some K^U , but there are several choices, so their sum is not well defined. For each summand it is possible to bring the different

choices together, but not for both summands simultaneously. This example shows that Proposition [12, Proposition A6.3] is wrong. The colimit in the category of groups is just K given by the evaluation at the fixed point.

1.8. Irreducible filter.

Definition 1.8.1. We call a (quasi)filter $F : \Lambda \rightarrow \mathcal{T}_X$ in a Grothendieck topology $\mathcal{T}_X$ on a scheme X *irreducible* if for every $\lambda \in \Lambda$ and every covering $U_i \rightarrow U_\lambda$, $i \in I$, there exists $i \in I$ and $(\kappa \rightarrow \lambda) \in \Lambda$ such that $F(\kappa \rightarrow \lambda) = U_\kappa \rightarrow U_\lambda$ factors through $U_i \rightarrow U_\lambda$ via some $(U_\kappa \rightarrow U_i) \in \mathcal{T}_X$.

Example 1.8.2. In an usual topological space, the filter of open neighborhoods of a point is irreducible, because if $\bigcup_{i \in I} U_i$ is a neighborhood of a point x , then U_i must be a neighborhood of x for at least one i .

Example 1.8.3. If a Grothendieck topology over a scheme X has the property that every covering is also set-theoretically a covering, then the quasifilter corresponding to a geometric point $Z = \text{Spec } K \rightarrow X$ is irreducible. For if $\text{Spec } K \rightarrow U \rightarrow X$ is a given neighborhood of x and $U_i \rightarrow U$, $i \in I$, is a covering, then there exists i and $\text{Spec } K \rightarrow U_i$ compatible with the given neighborhood.

The following lemma is a variant of the fact that for a point in a topological space the stalk of a presheaf and of its sheafification are the same.

Lemma 1.8.4. Let $\mathcal{T}_X$ denote a Grothendieck topology on X , let $F : \Lambda \rightarrow \mathcal{T}_X$ denote an irreducible filter. Let $\mathcal{P}$ denote a presheaf of abelian groups with sheafification $\mathcal{P}^\#$. Then $\mathcal{P}_F = (\mathcal{P}^\#)_F$.

Proof. We have a natural homomorphism

$$\mathcal{P}_F = \varinjlim_{\lambda \in \Lambda} \Gamma(U_\lambda, \mathcal{P}) \longrightarrow \varinjlim_{\lambda \in \Lambda} \Gamma(U_\lambda, \mathcal{P}^\#) = (\mathcal{P}^\#)_F.$$

Suppose that $s \in \mathcal{P}_F$, represented by $s \in \Gamma(U_\lambda, \mathcal{P})$, maps to 0 on the right. We may then assume that $s = 0$ in $\Gamma(U_\lambda, \mathcal{P}^\#)$ (because the colimit over a filter is the set theoretical colimit). This means that there exists a covering $U_i \rightarrow U_\lambda$, $i \in I$, such that the restrictions $\rho_{U_i}^{U_\lambda}(s) \in \Gamma(U_i, \mathcal{P})$ are zero. Since $(U_\kappa \rightarrow U_i \rightarrow U_\lambda) \in F$ for some $i \in I$, it follows that $s = 0$ in $\Gamma(U_\kappa, \mathcal{P})$ and hence in $\mathcal{P}_F$.

Now suppose that $s \in (\mathcal{P}^\#)_F$ is represented by $s \in \Gamma(U_\lambda, \mathcal{P}^\#)$, $\lambda \in \Lambda$. This means that there exists a covering $U_i \rightarrow U_\lambda$, $i \in I$, and elements $s_i \in \Gamma(U_i, \mathcal{P})$ which are compatible in the sense that $s_i = s_j$ in $\Gamma(U_i \times_{U_\lambda} U_j, \mathcal{P}_1)$. Let $(U_\kappa \rightarrow U_i \rightarrow U_\lambda) \in F$. Then $s = s_i$ on U_i and also on U_κ , hence s comes from the left. $\square$

Corollary 1.8.5. Let $X = \text{Spec } R$ be endowed with an affine Grothendieck topology, let $F : \Lambda \rightarrow \mathcal{T}_X$ denote an irreducible filter with structure stalk $\mathcal{O}_F$. Then for an R -module M we have $(M_{\text{top}})_F = M \otimes_R \mathcal{O}_F$.

Proof. We have by Lemma 1.8.4 and since tensor products commute with colimits the identities (setting $U_\lambda = \text{Spec } R_\lambda$)

$$\begin{aligned} (M_{\text{top}})_F &= \varinjlim_{\lambda \in \Lambda} \Gamma(U_\lambda, M_{\text{top}}) = \varinjlim_{\lambda \in \Lambda} M \otimes_R R_\lambda = M \otimes_R (\varinjlim_{\lambda \in \Lambda} R_\lambda) \\ &= M \otimes_R (\varinjlim_{\lambda \in \Lambda} \Gamma(U_\lambda, \mathcal{O}_{\text{top}})) = M \otimes_R \mathcal{O}_F. \end{aligned}$$

$\square$

Proposition 1.8.6. *Let $\mathcal{T}_R$ denote a Grothendieck topology on $\text{Spec } R$, let $R \rightarrow H$ be a ring homomorphism defining a quasifilter F by the small indexing category*

$$\Lambda_{R \rightarrow H} = \{[R \rightarrow B \rightarrow H], (R \rightarrow B) \in \mathcal{T}_R\}$$

(here $[\]$ denotes the isomorphism class). Suppose that F is irreducible and let $\mathcal{O}_F$ denote the stalk of the structure sheaf in F . Suppose in the first two statements that F is a filter. Then the following holds.

- (i) *There exists a natural ring homomorphism $\mathcal{O}_F \rightarrow H$.*
- (ii) *Suppose that $R \rightarrow R[T]$ is a cover in the topology. Then $\mathcal{O}_F \rightarrow H$ is a surjection.*
- (iii) *Suppose now that for each $R \rightarrow B \rightarrow H$ in $\Lambda_{R \rightarrow H}$ the image $\overline{B}$ of B in H has the property that the induced homomorphism $B \rightarrow \overline{B}$ belongs also to $\mathcal{T}_R$. Then F is a filter and the stalk $\mathcal{O}_F$ is a subring of H .*

Proof. (i). We always have a ring homomorphism $\varinjlim_{\lambda \in \Lambda} B_\lambda \rightarrow H$, and $\mathcal{O}_F = \varinjlim_{\lambda \in \Lambda} B_\lambda$ by Lemma 1.8.4 if F is an irreducible filter. (ii). For $a \in H$ consider the index λ given by

$$R \rightarrow R[T] \rightarrow H, T \mapsto a.$$

Then the image of $T \in B_\lambda$ in the colimit $\varinjlim_{\lambda \in \Lambda} B_\lambda$ (which is $\mathcal{O}_F$) maps to a .

We have to verify property (ii) in Definition 1.2.1 for the indexing category Λ_H . So assume that the first two rows of the following diagram commute.

$$\begin{array}{ccccc} R & \longrightarrow & B' & \longrightarrow & H \\ \downarrow = & & \downarrow \psi_1 & \downarrow \psi_2 & \downarrow = \\ R & \longrightarrow & B & \longrightarrow & H \\ \downarrow = & & \downarrow \varphi & & \downarrow = \\ R & \longrightarrow & \overline{B} & \longrightarrow & H \end{array} \quad \begin{array}{c} \mu \\ \uparrow \uparrow \\ \lambda \\ \uparrow \\ \kappa \end{array}$$

This can be commutatively extended to the third row, and the homomorphisms $\varphi \circ \psi_1, \varphi \circ \psi_2 : B' \rightarrow \overline{B}$ are identical, since they are identical as homomorphisms to H .

Suppose that $f \in \mathcal{O}_F$ maps to 0 in H , represented by $f \in B_\lambda$. Then the image of f in $B_\lambda \rightarrow \overline{B}_\lambda \rightarrow H$ is 0 and so $f = 0$ in $\mathcal{O}_F$. $\square$

1.9. Absolute filters and absolute stalks.

With absolute filters and absolute stalks we want to introduce a concept which is supposed to control all covers in a single-handed Grothendieck topology and contains in some sense all the information of the topology. The absolute stalk contains in particular the information about the induced closure operation (Corollary 2.4.10). A leading and motivating example is the role of the absolute integral closure R^+ of a domain which controls the plus closure, see Part 5.

Definition 1.9.1. We call a (quasi)filter $F : \Lambda \rightarrow \mathcal{T}_X$ in a Grothendieck topology $\mathcal{T}_X$ over X *absolute* if $U_\lambda \rightarrow X$ is a cover for every $\lambda \in \Lambda$ and if for every cover $Y \rightarrow U_\lambda$ in $\mathcal{T}_X$ ($\lambda \in \Lambda$) there exists $\kappa \in \Lambda$ and a factorization $U_\kappa \rightarrow Y \rightarrow U_\lambda$ (so that the composition comes from $\kappa \rightarrow \lambda$). We call the stalk of a presheaf $\mathcal{S}$ in an absolute filter an *absolute stalk*.

Remark 1.9.2. An absolute quasifilter has in particular the property that for every cover $Y \rightarrow X$ there exists $U_\lambda \rightarrow Y \rightarrow X$, as $X = U_{\lambda_0}$ for the final element $\lambda_0 \in \Lambda$. We will also encounter the condition that every morphism $U_\kappa \rightarrow U_\lambda$ indicated by F is a cover, and the stronger property that every morphism $Y \rightarrow U_\lambda$ in $\mathcal{T}_X$ is a cover for every $\lambda \in \Lambda$. See Proposition 1.10.4 and Examples 5.3.2 and 5.3.3 in the finite topology, where these conditions occur naturally. If F is given in an affine single-handed Grothendieck topology by a fixation $R \rightarrow H$, then this quasifilter is absolute if $R \rightarrow S \rightarrow H$ implies that $R \rightarrow S$ is a cover and that for given $S \rightarrow H$ and a cover $S \rightarrow T$ there exists an S -algebra homomorphism $T \rightarrow H$.

Definition 1.9.3. A scheme morphism $Z \rightarrow X$ (not necessarily in $\mathcal{T}_X$) is called an *absolute cover* if a morphism $U \rightarrow X$ in $\mathcal{T}_X$ is a cover if and only if there exists a factorization $Z \rightarrow U \rightarrow X$.

Lemma 1.9.4. *Let $\mathcal{T}_X$ be a single-handed Grothendieck topology on a scheme X . Then the following hold.*

- (i) *An absolute quasifilter $F : \Lambda \rightarrow \mathcal{T}_X$ is irreducible.*
- (ii) *An irreducible quasifilter $F : \Lambda \rightarrow \mathcal{T}_X$ with the property that every U_λ , $\lambda \in \Lambda$, is a cover of X , is absolute.*

Proof. (i). If $U \rightarrow U_\lambda$ is a cover (there are only single covers), $\lambda \in \Lambda$, then there exists $\kappa \in \Lambda$ and morphisms $U_\kappa \rightarrow U \rightarrow U_\lambda$, where $(U_\kappa \rightarrow U_\lambda) = F(\kappa \rightarrow \lambda)$.

(ii). We have to check the second condition of an absolute filter, so let $Y \rightarrow U_\lambda$ be a cover. Then by irreducibility there exists a $\kappa \in \Lambda$ and a factorization $U_\kappa \rightarrow Y \rightarrow U_\lambda$. $\square$

Lemma 1.9.4(i) is not true for non single-handed Grothendieck topologies, see Proposition 7.1.1(iv) for an example.

Lemma 1.9.5. *Let X denote a scheme with a Grothendieck topology $\mathcal{T}_X$ and an absolute filter $F : \Lambda \rightarrow \mathcal{T}_X$. Let $\mathcal{S}$ denote a sheaf of abelian groups on X_{top} . Then the natural homomorphism $\Gamma(X_{\text{top}}, \mathcal{S}) \rightarrow \mathcal{S}_F$ is injective.*

Proof. If $s \in \Gamma(X_{\text{top}}, \mathcal{S})$ becomes 0 in the absolute stalk, then it must be 0 in $\Gamma(U_\lambda, \mathcal{S})$ for some $U_\lambda \rightarrow X$, $\lambda \in \Lambda$. Because $U_\lambda \rightarrow X$ is a cover, s must itself be 0 by the local nature of a sheaf. $\square$

Corollary 1.9.6. *Let $\mathcal{T}_X$ denote a single-handed Grothendieck topology on a scheme X . Let M denote an $\mathcal{O}_X$ -module and let M_{top} denote the corresponding sheaf (of abelian groups) in the topology. Let $\Lambda \rightarrow \mathcal{T}_X$ denote an absolute filter. Then the absolute stalk is $\varinjlim_{\lambda \in \Lambda} \Gamma(U_\lambda, j_\lambda^*(M)) = \varinjlim_{\lambda \in \Lambda} \Gamma(U_\lambda, M_{\text{top}})$. In the affine case this means that $\varinjlim_{\lambda \in \Lambda} M \otimes_R R_\lambda = \varinjlim_{\lambda \in \Lambda} \Gamma(R_\lambda, M_{\text{top}})$.*

Proof. This follows from Lemma 1.8.4, Lemma 1.9.4(i) and Proposition 1.5.5(iv). $\square$

Example 1.9.7. Let $X = \text{Spec } R$ be an affine scheme with the single-handed Zariski topology (Example 1.5.8). Then every global function $g = (g_i)$, $g_i \in \Gamma(U_i, \mathcal{O})$, over a cover $\bigsqcup_{i \in I} U_i \rightarrow X$ will yield an element in an absolute stalk, so this cannot be an easy object. We look at the fixation given by $Z = \text{Spec}(\prod_{\mathfrak{p} \in X} R_{\mathfrak{p}})$. An object in the corresponding quasifilter is given (after refinement) by a covering $\bigsqcup_i U_i \rightarrow X$, $U_i = \text{Spec } R_{f_i}$, together with a morphism from Z to the covering. Such a morphism is given by a family of ring homomorphisms $\prod_{i \in I} R_{f_i} \rightarrow R_{\mathfrak{p}}$ for every $\mathfrak{p}$, and they are given by an i th projection

(where $i = i(\mathfrak{p})$ is such that $\mathfrak{p} \in D(f_i)$) and followed by $R_{f_i} \rightarrow R_{\mathfrak{p}}$. This quasifilter is absolute, since for a homomorphism $\prod_{i \in I} R_{f_i} \rightarrow R_{\mathfrak{p}}$ and a cover $\prod_{i \in I} R_{f_i} \rightarrow \prod_{j \in J} R_{g_j}$ there exists $\prod_{j \in J} R_{g_j} \rightarrow R_{\mathfrak{p}}$.

Assume now that R is a domain. The image of $\prod_{i \in I} R_{f_i}$ inside $\prod R_{\mathfrak{p}}$ is $\prod_{i \in J} R_{f_i}$ for a subset $J \subseteq I$ so that $\biguplus_{i \in J} U_i \rightarrow X$ is also cover. So by Proposition 1.8.6(iii) this quasifilter is a filter. Hence the absolute stalk is given by

$$\{(h_{\mathfrak{p}}), h_{\mathfrak{p}} \in R_{\mathfrak{p}}, \exists(f_1, \dots, f_n) = (1) \text{ and } a_i, k_i \text{ such that } h_{\mathfrak{p}} = a_i / f_i^{k_i} \text{ for some } i\}.$$

In this construction one can also restrict to the product over all local rings at maximal ideals, but the resulting absolute stalk is different. For a local ring R the fixation by R is an absolute filter with absolute stalk R . If we fix by $R \rightarrow R/\mathfrak{m}_R$ then this defines the same filter with again R as stalk (but the condition in Proposition 1.8.6(iii) is not fulfilled). If say R is a discrete valuation domain and we fix by $R \times Q(R)$ (as described in the first paragraph), then the resulting absolute stalk is $R \times Q(R)$.

1.10. Exactness.

An exact sequence of $\mathcal{O}_X$ -modules yields an exact sequence of sheaves in the flat (or the étale) topology (in the small site), but not in an arbitrary Grothendieck topology. The exactness of a complex of sheaves in a non-flat topology is a phenomenon in its own right, which we study here and in later sections for several topologies.

The exactness in the topology is in general quite a strong condition, in particular if $\mathcal{T}_X$ is big (this is already true for the big Zariski site). For example, if $f \in R$ is a non-zero divisor, then $0 \rightarrow R \xrightarrow{f} R$ is exact, but as soon as there is one object $S \in \mathcal{T}_R$ where f is a zero divisor, the corresponding complex of sheaves is not exact anymore. Therefore we will also deal with weaker exactness properties.

Corollary 1.10.1. *Let $X = \text{Spec } R$ denote an affine scheme endowed with an affine single-handed Grothendieck topology $\mathcal{T}_X$. Let $N \xrightarrow{\alpha} M \xrightarrow{\beta} L$ denote a complex of R -modules, and let $N_{\text{top}} \xrightarrow{\alpha} M_{\text{top}} \xrightarrow{\beta} L_{\text{top}}$ denote the corresponding complex of sheaves in the topology. Then the following are equivalent.*

- (i) *The sheaf complex $N_{\text{top}} \xrightarrow{\alpha} M_{\text{top}} \xrightarrow{\beta} L_{\text{top}}$ is exact.*
- (ii) *For every ring homomorphism $R \rightarrow S$ in $\mathcal{T}_R$ and every $s \in \Gamma(S, M_{\text{top}})$ such that $\beta(s) = 0$ in $\Gamma(S, L_{\text{top}})$ there exists a cover $S \rightarrow S'$ and $t \in \Gamma(S', N_{\text{top}})$ such that $\alpha(t)$ equals the restriction of s to $\Gamma(S', M_{\text{top}})$.*
- (iii) *For every ring homomorphism $R \rightarrow S$ in $\mathcal{T}_R$ and every $s \in M \otimes_R S$ such that $\beta(s) = 0$ in $L \otimes_R S$ there exists a cover $S \rightarrow S'$ and $s \in N \otimes_R S'$ such that $s = \alpha(t)$ in $M \otimes_R S'$.*

Proof. (ii) is just the formulation of the exactness of a complex of sheaves (Section 1.2) in the given affine single-handed Grothendieck topology. (iii) is the formulation for $\mathcal{T}_R$ -exactness (Definition 1.2.3) for the complex of presheaves given by tensoration with $R \rightarrow S$. Hence the equivalence (ii) $\Leftrightarrow$ (iii) follows from Proposition 1.2.4(iii) and Proposition 1.5.5(iii). $\square$

A complex $\mathcal{G} \rightarrow \mathcal{F} \rightarrow \mathcal{E}$ of sheaves on a Grothendieck topology $\mathcal{T}_X$ on X yields also a complex in a (quasi-)filter $F : \Lambda \rightarrow \mathcal{T}_X$. Its exactness can be characterized as follows.

Lemma 1.10.2. *Let X denote a scheme together with a Grothendieck topology $\mathcal{T}_X$, let $F : \Lambda \rightarrow \mathcal{T}_X$ denote a filter and let $\mathcal{G} \xrightarrow{\alpha} \mathcal{F} \xrightarrow{\beta} \mathcal{E}$ denote a complex of abelian sheaves. Then the following are equivalent.*

- (i) *The complex is exact in F , that is, the stalk complex $\mathcal{G}_F \xrightarrow{\alpha} \mathcal{F}_F \xrightarrow{\beta} \mathcal{E}_F$ is exact.*
- (ii) *For all $s \in \Gamma(U_\lambda, \mathcal{F})$ such that $\beta(s) = 0$ in $\Gamma(U_\lambda, \mathcal{E})$ ($\lambda \in \Lambda$) there exists $U_\kappa \rightarrow U_\lambda$ (indexed by $\kappa \rightarrow \lambda$ in Λ) and $t \in \Gamma(U_\kappa, \mathcal{G})$ such that $\alpha(t) = \rho_\kappa^\lambda(s)$ in $\Gamma(U_\kappa, \mathcal{F})$.*

Proof. (i) $\Rightarrow$ (ii). Since $\beta(s) = 0$ in $\Gamma(U_\lambda, \mathcal{E})$, this holds also in $\mathcal{E}_F$. Hence there exists $t \in \mathcal{G}_F$ such that $\alpha(t) = s$ in $\mathcal{F}_F$. So there exists $U_\kappa, \kappa \in \Lambda$, such that t is represented by $t \in \Gamma(U_\kappa, \mathcal{G})$ and such that $\alpha(t) = \rho_\kappa^\lambda(s)$ holds in $\Gamma(U_\kappa, \mathcal{F})$.

(ii) $\Rightarrow$ (i). Let $s \in \mathcal{F}_F$ map to 0 in $\mathcal{E}_F$. We may assume that $s \in \Gamma(U_\lambda, \mathcal{F})$ and that $\beta(s) = 0$ in $\Gamma(U_\lambda, \mathcal{E})$. So by (ii) there exists $U_\kappa \rightarrow U_\lambda$ and $t \in \Gamma(U_\kappa, \mathcal{G})$ mapping to $\rho_\kappa^\lambda(s)$, and this is then also true in the stalk. $\square$

Lemma 1.10.3. *Let $\mathcal{T}_X$ denote a Grothendieck topology on X and let $\mathcal{G} \xrightarrow{\alpha} \mathcal{F} \xrightarrow{\beta} \mathcal{E}$ denote an exact complex of sheaves of abelian groups on X_{top} . Then the complex is exact in every irreducible filter. In particular it is exact in an absolute filter, if $\mathcal{T}_X$ is single-handed.*

Proof. Let $F : \Lambda \rightarrow \mathcal{T}_X$ denote an irreducible filter. Let $s \in \mathcal{F}_F$ be represented by $s \in \Gamma(U_\lambda, \mathcal{F})$, $\lambda \in \Lambda$, and suppose that $\beta(s) = 0$ in $\mathcal{E}_F$. We may assume that $\beta(s) = 0$ in $\Gamma(U_\lambda, \mathcal{E})$. Due to exactness there exists a covering $U_i \rightarrow U_\lambda$, $i \in I$, and sections $t_i \in \Gamma(U_i, \mathcal{G})$, $i \in I$, such that $\alpha(t_i) = \rho_{U_i}^{U_\lambda}(s)$ in $\Gamma(U_i, \mathcal{F})$. Since F is irreducible there exists $i \in I$ and $\kappa \in \Lambda$ such that $(U_\kappa \rightarrow U_i \rightarrow U_\lambda) \in F$. Then $\rho_{U_\kappa}^{U_i}(t_i)$ maps to s in $\mathcal{F}_F$. $\square$

We characterize the exactness in an absolute filter. This is in general a weaker property than the exactness in the topology, but under certain conditions they are equivalent.

Proposition 1.10.4. *Let $X = \text{Spec } R$ denote an affine scheme endowed with an affine single-handed Grothendieck topology $\mathcal{T}_X$. Let $N \xrightarrow{\alpha} M \xrightarrow{\beta} L$ denote a complex of R -modules and let $F : \Lambda \rightarrow \mathcal{T}_X$ be an absolute filter with absolute stalk H . Then the following are equivalent.*

- (i) *The sheaf complex $N_{\text{top}} \xrightarrow{\alpha} M_{\text{top}} \xrightarrow{\beta} L_{\text{top}}$ is exact in the absolute filter.*
- (ii) *The complex $N \otimes_R H \xrightarrow{\alpha} M \otimes_R H \xrightarrow{\beta} L \otimes_R H$ is exact.*
- (iii) *For every $\lambda \in \Lambda$ and every $s \in \Gamma(U_\lambda, M_{\text{top}})$ mapping to 0 in $\Gamma(U_\lambda, L_{\text{top}})$ there exists $(\kappa \rightarrow \lambda) \in \Lambda$ and $t \in \Gamma(U_\kappa, N_{\text{top}})$ mapping to $\rho_\kappa^\lambda(s) \in \Gamma(U_\kappa, M_{\text{top}})$.*
- (iv) *For every $\lambda \in \Lambda$ and every $s \in M \otimes_R R_\lambda$ mapping to 0 in $L \otimes_R R_\lambda$ there exists $(\kappa \rightarrow \lambda) \in \Lambda$ and $t \in N \otimes_R R_\kappa$ mapping to $s \otimes 1 \in M \otimes_R R_\kappa$.*

If the filter is covering (Remark 1.9.2) in the sense that for every $\kappa \rightarrow \lambda$ in Λ the morphism $U_\kappa \rightarrow U_\lambda$ is a cover, then this is also equivalent to the following.

- (v) *The sheaf complex $N_{\text{top}} \xrightarrow{\alpha} M_{\text{top}} \xrightarrow{\beta} L_{\text{top}}$ is $\mathcal{T}$ -exact for every $U_\lambda, \lambda \in \Lambda$.*

If the filter is covering in the sense that for every $\lambda \in \Lambda$ every morphism $Y \rightarrow U_\lambda$ in $\mathcal{T}_X$ is a cover, then this is also equivalent to the following.

- (vi) For every cover $Y \rightarrow X$ and every $s \in \Gamma(Y, M_{\text{top}})$ mapping to 0 in $\Gamma(Y, L_{\text{top}})$ there exists $\kappa \in \Lambda$, a morphism $U_\kappa \rightarrow Y$ in $\mathcal{T}_X$ and $t \in \Gamma(U_\kappa, N_{\text{top}})$ mapping to $\rho_{U_\kappa}^Y(s) \in \Gamma(U_\kappa, M_{\text{top}})$.
- (vii) For every cover $Y \rightarrow X$ and every $s \in \Gamma(Y, M_{\text{top}})$ mapping to 0 in $\Gamma(Y, L_{\text{top}})$ there exists a cover $Y' \rightarrow X$ and a morphism $Y' \rightarrow Y$ and $t \in \Gamma(Y', N_{\text{top}})$ mapping to $\rho_{Y'}^Y(s) \in \Gamma(Y', M_{\text{top}})$.

If the topology is covering (Definition 1.5.4), then this is also equivalent to

- (viii) The sheaf complex $N_{\text{top}} \xrightarrow{\alpha} M_{\text{top}} \xrightarrow{\beta} L_{\text{top}}$ is exact.

Proof. (i) and (ii) are equivalent by Corollary 1.8.5. (iii) is the explicit version of (i) (Lemma 1.10.2) and (iv) is the explicit version of (ii). The implications (viii) $\Rightarrow$ (v), (v) $\Rightarrow$ (i)-(iv), (i)-(iv) $\Rightarrow$ (vi) and (vi) $\Leftrightarrow$ (vii) hold without further conditions, so we prove them first. (viii) $\Rightarrow$ (v) is a restriction (Proposition 1.2.4).

(v) $\Rightarrow$ (iii). In the given situation of (iii) there exists by (v) a cover $Y \rightarrow U_\lambda$ such that the restriction of s comes from the left. By definition of an absolute filter there exists a morphism $U_\kappa \rightarrow Y$, and so the restriction of s to $\Gamma(U_\kappa, M_{\text{top}})$ comes from an element in $\Gamma(U_\kappa, N_{\text{top}})$.

(iii) $\Rightarrow$ (vi). In the given situation there exists $\lambda \in \Lambda$ and a morphism $U_\lambda \rightarrow Y$, since F is an absolute filter. The condition holds also for $\rho_{U_\lambda}^Y(s)$ and by (iii) there exists $U_\kappa \rightarrow U_\lambda$ such that s comes from an element in $\Gamma(U_\kappa, N_{\text{top}})$. (vi) $\Rightarrow$ (vii) is trivial ($Y' = U_\kappa$). For (vii) $\Rightarrow$ (vi) just take $U_\kappa \rightarrow Y' \rightarrow Y$.

Now assume that the indicated morphisms in the filter are covers. Then (iii) $\Rightarrow$ (v) is trivial.

(vii) $\Rightarrow$ (iii) under the given assumption. Suppose the situation of (iii). There exists by (vii) a morphism $Y' \rightarrow U_\lambda$, $Y' \rightarrow X$ a cover, such that the restriction of s to Y' comes from the left. By assumption on the filter, $Y' \rightarrow U_\lambda$ is a cover. Hence by the definition of an absolute filter there exists $\kappa \rightarrow \lambda$ and a factorization $U_\kappa \rightarrow Y' \rightarrow U_\lambda$. If the topology is covering, then (vii) $\Rightarrow$ (viii) is trivial. $\square$

For an example where Proposition 1.10.4(vi) and (vii) hold, but not (i)-(v), see Example 5.3.3.

1.11. Cohomology.

The category of sheaves of abelian groups in a Grothendieck topology is an abelian category [49, Theorem II.2.15] and every abelian sheaf can be embedded into an injective sheaf [49, Proposition III.1.1]. In particular, a sheaf has an injective resolution and so it is possible to define *cohomology* of a sheaf as the derived functor for the functor of global sections $\mathcal{F} \mapsto \Gamma(X, \mathcal{F})$ [27, Section III.1].

We will be interested in cohomology applied to a short exact sequence $0 \rightarrow \text{Syz} \rightarrow \mathcal{O}_{\text{top}}^n \xrightarrow{f_1, \dots, f_n} I^{\text{top}} \rightarrow 0$ given by ideal generators $I = (f_1, \dots, f_n) \subseteq R$, as this gives under certain conditions a bijection $I^{\text{top-cl}}/I = H^1((\text{Spec } R)_{\text{top}}, \text{Syz})$ (Proposition 2.8.1). So the first cohomology of the syzygy sheaf measures the difference between an ideal and its closure induced by the Grothendieck topology. We will also compute that in the surjective topology (in characteristic zero) and in the Frobenius topology the first cohomology of a quasicoherent module is zero (Theorem 3.10.2, Proposition 4.5.1) using Čech cohomology.

To review *Čech cohomology* let X be a site and let $\mathcal{U} = (U_i \rightarrow U, i \in I)$ be a covering of $U \in \mathcal{T}_X$. We write $U_{i_0 \dots i_k} = U_{i_0} \times_U \dots \times_U U_{i_k}$. The *Čech diagram* (of this covering) is

$$\dots \bigoplus_{i_0, i_1, i_2, i_3} U_{i_0 i_1 i_2 i_3} \rightrightarrows \bigoplus_{i_0, i_1, i_2} U_{i_0 i_1 i_2} \rightrightarrows \bigoplus_{i_0, i_1} U_{i_0 i_1} \rightrightarrows \bigoplus_{i_0} U_{i_0} \rightarrow U,$$

where the t th arrow $\bigoplus_{i_0, \dots, i_{k+1}} U_{i_0 \dots i_{k+1}} \rightarrow \bigoplus_{i_0, \dots, i_k} U_{i_0 \dots i_k}$ ($(i_0, \dots, i_{k+1}) \in I^{k+1}$) is on every open component just the projection $q_t = p_{i_0 \dots \check{i}_t \dots i_{k+1}} : U_{i_0 \dots i_{k+1}} \rightarrow U_{i_0 \dots \check{i}_t \dots i_{k+1}}$ (see [24, V.1.10] or [49, III §2]).

Applying a presheaf $\mathcal{P}$ of abelian groups to this diagram we get the Čech diagram of the covering with values in $\mathcal{P}$ [24, V.2.3.3], and taking the alternating sum over the arrows of this diagram we get the *Čech complex* of $\mathcal{P}$ with respect to this covering. Explicitly, $C^k(\mathcal{U}, \mathcal{P}) = \prod_{i_0, \dots, i_k} \Gamma(U_{i_0 \dots i_k}, \mathcal{P})$ and the derivation is $d^k : C^k(\mathcal{U}, \mathcal{P}) \rightarrow C^{k+1}(\mathcal{U}, \mathcal{P})$ by sending $(d^k s)_{i_0, \dots, i_{k+1}} = \sum_{t=0}^{k+1} (-1)^t \rho_t(s_{i_0 \dots \check{i}_t \dots i_{k+1}})$, where $\rho_t = \rho_{i_0, \dots, \check{i}_t \dots i_{k+1}}^{i_0 \dots \check{i}_t \dots i_{k+1}}$ stands for the restriction to the projection q_t . The homology groups of this complex are called the *Čech cohomology groups* and are denoted by $\check{H}^k(U_i \rightarrow U, \mathcal{P})$.

If $V_j \rightarrow U$, $j \in J$, is a second covering which refines $\mathcal{U}$, then there exists a refinement homomorphism $\check{H}^k(\mathcal{U}, \mathcal{P}) \rightarrow \check{H}^k(V_j \rightarrow U, \mathcal{P})$. The Čech cohomology groups are defined as the colimit $\check{H}^k(U, \mathcal{P}) = \varinjlim_{\mathcal{U}} \check{H}^k(\mathcal{U}, \mathcal{P})$ over all coverings $\mathcal{U}$ of U [24, V.2.4.5]. For a sheaf $\mathcal{P}$, $\check{H}^1(U, \mathcal{P})$ equals the derived functor cohomology $H^1(U, \mathcal{P})$ [49, Corollary III.2.10].

The assumption in the next lemma holds in the constructible topology (Lemma 3.6.5, Proposition 3.10.1) and in the Frobenius topology (Proposition 4.5.1) and generalizes Lemma 1.6.7.

Lemma 1.11.1. *Suppose that a cover $Y \rightarrow X$ in a Grothendieck topology on a scheme X has the property that also the diagonal $\Delta : Y \rightarrow Y \times_X Y$ is a cover (this is in particular the case if a projection $Y \times_X Y \rightarrow Y$ is an isomorphism). Then for a sheaf $\mathcal{F}$ of abelian groups on X_{top} we get in the Čech complex either the 0-map or the identity,*

$$\Gamma(X, \mathcal{F}) \rightarrow \Gamma(Y, \mathcal{F}) \xrightarrow{0} \Gamma(Y \times_X Y, \mathcal{F}) \xrightarrow{\text{id}} \Gamma(Y \times_X Y \times_X Y, \mathcal{F}) \xrightarrow{0} \dots$$

It follows that the Čech cohomology is trivial, i.e. $\check{H}^0(Y \rightarrow X, \mathcal{F}) = \Gamma(Y, \mathcal{F})$ and $\check{H}^k(Y \rightarrow X, \mathcal{F}) = 0$ for $k \geq 1$.

Proof. We show that $\rho_t = \rho(q_t)$ is the same mapping for every t . Since $q_t \circ \Delta_{t, t+1} = q_{t+1} \circ \Delta_{t, t+1}$ we get $\rho(\Delta_{t, t+1}) \circ \rho_t = \rho(\Delta_{t, t+1}) \circ \rho_{t+1}$. As Δ is a cover, also $\Delta_{t, t+1}$ is a cover, hence it follows that $\rho_t = \rho_{t+1}$. All these mappings are identities, since for a cover $U \rightarrow V$ with a covering section $V \rightarrow U$ the induced maps on a sheaf are always isomorphisms. In the Čech complex we have either an even or an odd alternating sum of identities, so the sum is either 0 or the identity. The statements about the cohomology follow. $\square$

It is also possible to define *local cohomology* for an open subset $U \rightarrow X$ in a Grothendieck topology on X (see [24, V.6.3]). For an abelian sheaf $\mathcal{F}$ define

$$\Gamma_{X-U}(\mathcal{F}) = \{s \in \Gamma(X, \mathcal{F}) : s|_U = 0\} = \ker(\rho_U^X : \Gamma(X, \mathcal{F}) \rightarrow \Gamma(U, \mathcal{F}))$$

($X - U$ is just a symbol for us). This is again a left exact functor and gives rise to a derived functor, denoted by $H_{X-U}^k(-)$. There exists a long exact sequence (as in [22, Corollaire

I.2.9]) $0 \rightarrow \Gamma_{X-U}(\mathcal{F}) \rightarrow \Gamma(X, \mathcal{F}) \rightarrow \Gamma(U, \mathcal{F}) \rightarrow H_{X-U}^1(\mathcal{F}) \rightarrow H^1(X, \mathcal{F}) \rightarrow H^1(U, \mathcal{F})$ [24, Proposition V.6.5]. We will use this in Section 2.8.

2. CLOSURE OPERATIONS AND GROTHENDIECK TOPOLOGIES

We describe now how a Grothendieck topology on an affine scheme $X = \operatorname{Spec} R$ defines a closure operation for R -submodules $N \subseteq M$ and we characterize which closure operations can be obtained in this way. In general there will be several more or less natural Grothendieck topologies which realize the same closure operation. These Grothendieck topologies come along with a variety of gadgets such as cohomology, ring and modules of global sections, notions of stalks and exactness, which we try to understand in the context of the closure operation.

2.1. Forcing algebras.

We recall the definition of (universal) forcing algebras following Hochster [33]. Let R be a commutative ring and let $N \subseteq M$ denote R -modules such that M/N is finitely presented, $s \in M$. This means that we have a resolution

$$R^\nu \xrightarrow{D} R^\mu \rightarrow M/N \rightarrow 0,$$

where $D = (r_{ij})_{ij}$ is a $\mu \times \nu$ -matrix with entries in R . Let $\tilde{s} = (s_1, \dots, s_\mu) \in R^\mu$ denote a lifting of s . In this situation we make the following definition.

Definition 2.1.1. The finitely generated R -algebra

$$B = R[T_1, \dots, T_\nu] / (DT - \tilde{s}) = R[T_1, \dots, T_\nu] / (\sum r_{1j}T_j - s_1, \dots, \sum r_{\mu j}T_j - s_\mu)$$

is called a *forcing algebra* for (M, N, s) . Accordingly, $\operatorname{Spec} B$ is called the *forcing scheme*.

Remark 2.1.2. Note that the three sets of data (M, N, s) , $(R^\mu, \tilde{N}, \tilde{s})$, where $\tilde{N}$ is the kernel of $R^\mu \rightarrow M/N$, and $(\overline{M}, 0, \overline{s})$, where $\overline{M} = M/N$ and $\overline{s}$ denotes the image of s , have the same forcing algebras. This means that we may simplify the situation by either passing to a torsion-free submodule inside a free module or to the 0-submodule inside a module. Or the other way round: forcing algebras can only be helpful to understand such closure operations which are independent of the presentation, see Proposition 2.2.6(iii).

Remark 2.1.3. The construction of the forcing algebra is functorial with respect to ring homomorphisms $R \rightarrow R'$. By right exactness we get immediately the resolution

$$R'^\nu \xrightarrow{D'} R'^\mu \rightarrow (M/N) \otimes_R R' = M'/N' \rightarrow 0,$$

where N' is the image of $N \otimes_R R'$ inside $M' = M \otimes_R R'$, and the forcing algebra over R' built with D' is just $B \otimes_R R'$.

The forcing algebra has the following universal property.

Proposition 2.1.4. *Let R denote a commutative ring and let B denote a forcing algebra for R -modules $N \subseteq M$ such that M/N is finitely presented, $s \in M$. Then the following holds.*

- (i) *The element $s \otimes 1$ lies in the image of $N \otimes_R B \rightarrow M \otimes_R B$.*
- (ii) *If A is an R -algebra such that $s \otimes 1 \in \operatorname{im}(N \otimes_R A \rightarrow M \otimes_R A)$, then there exists an R -algebra homomorphism $B \rightarrow A$.*

Proof. We use the notation of Remark 2.1.2. The condition that $s \otimes 1 \in \text{im}(N \otimes_R A \rightarrow M \otimes_R A)$ is equivalent to the property that $\bar{s} \otimes 1$ becomes 0 in $(M/N) \otimes_R A$, and this is equivalent to the property that $\tilde{s} \otimes 1$ in A^μ is in the image of $A^\nu \rightarrow A^\mu$. This means that the inhomogeneous linear system of equations $\sum r_{1j}T_j - s_1, \dots, \sum r_{\mu j}T_j - s_\mu$ has a solution in A . For the forcing algebra B this is true by the very definition, which gives (i). (ii). If a solution over A exists, then we just have to send the variables T_j to the corresponding coefficients in A , so there exists an R -algebra homomorphism $B \rightarrow A$. $\square$

Recall that a ring homomorphism $R \rightarrow A$ is called *pure* if for every R -module M the induced map $M \rightarrow M \otimes_R A$ is injective. For a pure homomorphism the contraction of an extended ideal is the ideal itself. Pure ring homomorphisms induce a spec-surjective morphism. We note the following basic observation.

Lemma 2.1.5. *Let R be a commutative ring. Let $N \subseteq M$ be R -modules such that M/N is finitely presented, let $s \in M$ be an element. Let*

$$B = R[T_1, \dots, T_\nu]/(DT - \tilde{s})$$

be the forcing algebra for (M, N, s) . Then the following are equivalent.

- (i) $s \in N$.
- (ii) *There is a ring section $B \rightarrow R$ for $R \rightarrow B$.*
- (iii) *$R \rightarrow B$ splits as a homomorphism of R -modules (that is, R is a direct summand in B).*
- (iv) *$R \rightarrow B$ is pure.*

Proof. (i) $\Rightarrow$ (ii) follows from the universal property of a forcing algebra (Proposition 2.1.4(ii)). The implications (ii) $\Rightarrow$ (iii) and (iii) $\Rightarrow$ (iv) hold for arbitrary ring homomorphisms. So suppose that (iv) holds. Apply the purity property to the R -module M/N . Let $\bar{s}$ denote the image of s in M/N . Then by Proposition 2.1.4(i) $\bar{s}$ becomes 0 in $(M/N) \otimes_R B$, so it must be 0 by purity. $\square$

Remark 2.1.6. Let R denote a commutative ring, and let $I = (f_1, \dots, f_n)$ denote a finitely generated ideal, let $f \in R$. In this case the resolution is just $R^n \xrightarrow{f_i} R \rightarrow R/I \rightarrow 0$ and the forcing algebra

$$B = R[T_1, \dots, T_n]/(f_1T_1 + \dots + f_nT_n + f)$$

is given by one affine-linear equation. In this case we can add to the equivalent conditions in Lemma 2.1.5 the property that $R \rightarrow B$ is cyclic pure, which means by definition that $IB \cap R = I$ for every ideal $I \subseteq R$.

2.2. Admissible closure operations.

The closure operations we will consider will be defined for submodules in a certain class of modules and will be persistent with respect to certain ring homomorphisms. We fix the following situation.

Situation 2.2.1. Let G denote a subcategory of commutative rings and suppose that a subcategory $\mathcal{M} = \mathcal{M}_R$ of R -modules is given for each $R \in G$, such that for every ring homomorphism $\varphi : R \rightarrow S$ in G the module $M \otimes_R S$ belongs to $\mathcal{M}_S$. A *closure operation* for $\mathcal{M}$ is an assignment $N \mapsto N^c$ from $\mathcal{M}_R$ to $\mathcal{M}_R$ with $N \subseteq N^c = (N^c)^c$. The closure operation is *G-persistent* if for $\varphi : R \rightarrow S$ in G one has $\text{im}(N^c) \subseteq (N')^c$ in $M \otimes_R S$, where $N' = \text{im}(N \otimes_R S)$.

Remark 2.2.2. The category G will be the full category of all rings, or all noetherian rings, all domains, all rings of characteristic p etc. We may also restrict the morphisms to homomorphisms of finite type, inclusion of domains, etc.

The category of modules $\mathcal{M}$ will be either the category of all ideals (then we do not insist that $I \otimes_R S$ belongs to $\mathcal{M}_S$), or the category of all finitely generated R -module, or all torsion-free modules, or the category of all modules.

Definition 2.2.3. Let an assignment c on $\mathcal{M}$ as described in 2.2.1 be given. We call a forcing algebra B for (M, N, s) for R -modules $N \subseteq M$ such that M/N is finitely presented *admissible* (with respect to the closure operation) or *c-admissible* if $s \in N^c$.

We call a ring homomorphism $\varphi : R \rightarrow S$ *c-pure* if for every $U \subseteq V$ in $\mathcal{M}_R$ we have that $z \otimes 1 \in \text{im}(U \otimes_R S \rightarrow V \otimes_R S)$ implies that $z \in U^c$.

Proposition 2.2.4. Suppose the situation described in 2.2.1. Then the following holds.

- (i) $R \rightarrow S$ is *c-pure* if and only if for every forcing algebra $R \rightarrow B$ such that $S \rightarrow B \otimes_R S = B_S$ is *c-admissible*, then also $R \rightarrow B$ is *c-admissible*.
- (ii) The composition of two *c-pure* homomorphisms $R \xrightarrow{\varphi} S \xrightarrow{\psi} T$ is also *c-pure*.

Proof. (i). Let B be the forcing algebra for (M, N, s) . Then $s \otimes 1$ is in the image of $N \otimes_R S \rightarrow M \otimes_R S$ if and only if $s \otimes 1$ is in the closure of the extended module N' . This is by definition equivalent to $S \rightarrow B_S$ being *c-admissible*. That $R \rightarrow B$ is *c-admissible* is equivalent to $s \in N^c$.

(ii) follows direct or from (i), since if $R \rightarrow B$ is a forcing algebra such that $T \rightarrow B_T$ is admissible, then first $S \rightarrow B_S$ is admissible and then $R \rightarrow B$ is admissible. $\square$

Definition 2.2.5. Suppose the situation described in 2.2.1. We call c *G-admissible*, if the following conditions hold.

- (i) c is a closure operation, that is, $N \subseteq N^c = (N^c)^c$.
- (ii) The admissible forcing algebras are *c-pure*.
- (iii) The closure operation is *G-persistent*.

Proposition 2.2.6. Suppose that we have an admissible closure operation on a subcategory of noetherian rings and for finitely generated modules. Then the following hold.

- (i) (*Module-persistence*) Let $N \subseteq M$, $s \in M$ and let $\varphi : M \rightarrow M'$ be an R -module homomorphism. Set $N' := \varphi(N)$ and suppose that $s \in N^c$. Then also $\varphi(s) \in (N')^c$.
- (ii) (*Independence of presentation*) Let $N \subseteq M$, $s \in M$. Let $p : \tilde{M} \rightarrow M \rightarrow 0$ and $\tilde{N} := p^{-1}(N)$ and let $\tilde{s} \in \tilde{M}$ be such that $p(\tilde{s}) = s$. Then $s \in N^c$ if and only if $\tilde{s} \in \tilde{N}^c$. In particular, $s \in N^c$ if and only if $\bar{s} \in 0^c$ in M/N .
- (iii) The closure operation is order preserving, that is, for submodules $N \subseteq L \subseteq M$ of a finitely generated R -module M we have $N^c \subseteq L^c$.
- (iv) Suppose also that the closure is persistent with respect to homomorphisms of finite type. Let $N_i \subseteq M_i, i = 1, \dots, k$, be submodules and set $N = N_1 \times \dots \times N_k \subseteq M_1 \times \dots \times M_k = M$. Then $N^c = N_1^c \times \dots \times N_k^c$, where N_i^c is the closure inside M_i .

Proof. (i). Let $R \rightarrow B$ denote the (admissible) forcing algebra for (M, N, s) . Then $s \otimes 1$ is in the image of $N \otimes_R B \rightarrow M \otimes_R B$. But then $\varphi(s) \otimes 1$ is in the image of $N' \otimes_R B \rightarrow M' \otimes_R B$.

(ii). One direction follows from (i). For the other direction suppose that $s \in N^c$, let B denote a forcing algebra for these data. Then B is also the forcing algebra for $(\tilde{M}, \tilde{N}, \tilde{s})$.

(iii) follows from (i) and (ii).

(iv). By (ii) we may assume that $N = 0$. Let $s = (s_1, \dots, s_k)$. If $s_i \in 0^c$ in M_i for every i , then the corresponding forcing algebras B_i are c -admissible, hence c -pure. Because of the persistence property this holds in the universal sense. Therefore also their tensor product B (which is the forcing algebra of s) is c -pure and since $s = 0$ in $M \otimes_R B$ we get $s \in 0^c$. On the other hand, if $s \in 0^c$, then also $s_i \in 0^c$ by applying (i) to the projections $M \rightarrow M_i$. $\square$

Remark 2.2.7. The module persistence of an admissible closure operation implies that the assignment $M \mapsto 0^c$ is a functor from $R\text{-Mod}$ to itself (which is not a projection). An inclusion $N \subseteq M$ yields to an inclusion $0_N^c \subseteq 0_M^c$, but this functor is neither exact in the middle nor does it respect surjections. The left exactness is related to hereditary torsion theories; see Lemma 2.4.9 and Section 6.3.

Lemma 2.2.8. *An order preserving closure operation for ideals in a local noetherian ring $(R, \mathfrak{m})$ is trivial if it is trivial for all $\mathfrak{m}$ -primary ideals.*

Proof. Let $I \subseteq R$ and suppose that $f \in I^c$, $f \notin I$. If f is a unit, then also $f \in \mathfrak{m}^c$. So we may assume that f is not a unit. Then there exists $k \in \mathbb{N}$ such that $f \notin I + \mathfrak{m}^k$ (by Krull's intersection theorem [12, Corollary 5.4] applied to R/I). Then $f \in I^c \subseteq (I + \mathfrak{m}^k)^c$, and the $\mathfrak{m}$ -primary ideal $I + \mathfrak{m}^k$ is not c -closed. $\square$

Remark 2.2.9. If an order preserving closure operation is given on an R -module M , then an arbitrary intersection of a family of closed submodules $N_j = N_j^c$, $j \in J$, is again closed. Therefore one can define a topology on M (in the usual sense) by declaring finite unions of closed submodules to be the closed subsets. As the union of two submodules is a submodule only if one of them is contained in the other, the closed submodules in this topology are exactly the closed submodules for the operation.

Example 2.2.10. There are several examples of closure operations for ideals which are trivial for $\mathfrak{m}$ -primary ideals, but not in general, so these are not order preserving and hence not admissible (and not induced by a Grothendieck topology). E.g. the so-called *top-dimensional closure* (or *equidimensional hull* [39, Definition 5.7]) of an ideal $I \subseteq R = K[X_1, \dots, X_m]$ (or more general) is given in the following way: let $I = \mathfrak{q}_1 \cap \dots \cap \mathfrak{q}_r$ be a primary decomposition such that $\mathfrak{q}_i$ is $\mathfrak{p}_i$ -primary and such that the $\mathfrak{p}_1, \dots, \mathfrak{p}_s$ ($s \leq r$) have minimal height (or $R/\mathfrak{p}_i$ has maximal dimension) among the $\mathfrak{p}_i$. Then $I^{\text{top}} = \mathfrak{q}_1 \cap \dots \cap \mathfrak{q}_r$. If I is $(X_1, \dots, X_m)$ -primary, then of course $I^{\text{top}} = I$.

Example 2.2.11. The “*symbolic closure operation*” $I \mapsto \bigcap_{I \subseteq \mathfrak{p} \text{ minimal prime}} R \cap IR_{\mathfrak{p}}$ is not admissible, since it is trivial on the primary ideals, but not in general. For a prime ideal $\mathfrak{p}$ this assignment maps the ordinary power $\mathfrak{p}^n$ to the symbolic power $\mathfrak{p}^{(n)}$, [12, Section 3.9]. See also Section 6.4, where the divisorial topology induces the symbolic powers for prime ideals of height one.

Example 2.2.12. The Ratliff-Rush *closure* of an ideal $I \subseteq R$ is defined as $\tilde{I} = \bigcup_{n \geq 1} (I^{n+1} : I^n)$, see [56]. This closure is not an admissible closure operation, since it is not order preserving

Example 2.2.13. The socle $\text{soc}(M) = \{s \in M : sm = 0\}$ for an R -module over a local ring $(R, \mathfrak{m})$ is not a closure operation. E.g. for $R = K[x]/(x^3)$ we get $\text{soc}(R) = (x^2)$, but its “socle-closure” would be (x) .

Definition 2.2.14. Let c denote a closure operation on $\mathcal{M}$. Consider for a pair $N \subseteq M$, $s \in M$, $s \in N^c$ the properties

- (i) There exists an R -submodule $N \subseteq M' \subseteq M$ such that $s \in M'$, $s \in N^c$ inside M' and M'/N is finitely presented.
- (ii) There exists a finitely generated R -submodule $N' \subseteq N$ such that $s \in (N')^c$.

We say that the closure operation has the first finiteness property, if (i) holds (for every $N \subseteq M$) and the second finiteness property, if (ii) holds.

2.3. Examples of admissible closure operations.

We give some examples of admissible closure operations.

Example 2.3.1. We consider the identical closure operation, where of course the conditions (i) and (iii) (for the category of all rings) in Definition 2.2.5 are fulfilled. The ring homomorphisms $R \rightarrow A$ which are pure with respect to the identical closure operation are exactly the pure ring homomorphisms. An admissible forcing algebra for the trivial closure operation is pure by Lemma 2.1.5, hence the identical closure is admissible.

Definition 2.3.2. Let $\mathcal{D}$ be a subcategory of rings. Such a category defines a closure operation for R -submodules $N \subseteq M$ by declaring for $s \in M$ that $s \in N^c$ if and only if

$$s \otimes 1 \in \text{im}(N \otimes_R T \rightarrow M \otimes_R T) \text{ for all ring homomorphisms } R \rightarrow T, T \in \mathcal{D}.$$

We call this the *closure operation associated to $\mathcal{D}$* (the rings in $\mathcal{D}$ are the testrings).

Lemma 2.3.3. Let R be a commutative ring, let $\mathcal{D}$ be a category of rings with the induced closure operation c . Let $N \subseteq M$ be R -modules such that M/N is finitely presented, $s \in M$, let B be the forcing algebra. Then the following are equivalent.

- (i) $s \in N^c$.
- (ii) For every $R \rightarrow T$, $T \in \mathcal{D}$, the induced forcing algebra $T \rightarrow B_T = B \otimes_R T$ has a ring section.
- (iii) For every $R \rightarrow T$, $T \in \mathcal{D}$, the induced forcing algebra $T \rightarrow B_T$ has an T -module section.
- (iv) For every $R \rightarrow T$, $T \in \mathcal{D}$, the induced forcing algebra $T \rightarrow B_T$ is pure.

Proof. This follows from the functoriality of forcing algebras (Remark 2.1.3) and Lemma 2.1.5. $\square$

Proposition 2.3.4. Let $N \mapsto N^c$ denote a closure operation given by a subcategory $\mathcal{D}$ of rings. Then this closure operation is admissible with respect to all ring homomorphisms.

Proof. It is clear that $N \subseteq N^c = (N^c)^c$ and that the closure is persistent for all ring homomorphisms $R \rightarrow R'$. By Lemma 2.3.3 a forcing algebra $R \rightarrow B$ is c -admissible if and only if for every $R \rightarrow T$, $T \in \mathcal{D}$, the induced mapping $T \rightarrow B_T = B \otimes_R T$ has a ring-section. But an R -algebra B with this last property is in general c -pure. For if $s \otimes 1 \in \text{im}(U \otimes_R B \rightarrow V \otimes_R B)$ and if $R \rightarrow T$, $T \in \mathcal{D}$ is given, then we have a factorization $R \rightarrow B \rightarrow T$ showing that $s \otimes 1 \in \text{im}(U \otimes_R T \rightarrow V \otimes_R T)$. $\square$

We describe some closure operations coming from such a subcategory of rings.

Example 2.3.5. Take $\mathcal{D}$ to be the category of fields. The corresponding ideal closure operation is the radical $\text{rad}(I)$ of the ideal I , see Section 3.1. This closure operation is related to the surjective topology, where the coverings are given by spec-surjective morphisms.

Example 2.3.6. Suppose that R is a (noetherian) ring and let $\mathcal{D}$ be the category of (discrete) valuation domains. Then the induced closure operation is the integral closure, which we will study in its relation to the submersive topology in [4].

Example 2.3.7. The weak subintegral closure was introduced by Leahy and Vitulli ([67], [59]). In [8] we show that *crosses*, i.e. one dimensional schemes consisting of two normal components meeting in one point transversally give a category of test rings for this closure operation. The *scheme of axes*, i.e. one dimensional schemes consisting of normal components meeting transversally in one point and having maximal embedding dimension, yield also an interesting category of test rings, which induce the so-called *axes closure*, which we study in its relation to the continuous closure in [7].

Example 2.3.8. Let $\mathcal{D}$ be the category of regular rings. Then the induced closure operation is the so-called *regular closure* [35]. The regular closure is clearly contained in the integral closure and in the radical. It does contain the tight closure of an ideal, since tight closure is persistent and every ideal in a regular ring is tightly closed [38, Theorem 2.3 and Theorem 1.3(e)]. Therefore $I^* \subseteq I^{\text{reg}} \subseteq \bar{I} \subseteq \text{rad}(I)$.

2.4. The closure operation induced by a Grothendieck topology.

Let X denote a scheme endowed with a Grothendieck topology which is a refinement of the trivial topology (Example 1.5.2) (or of the single-handed Zariski topology (Remark 1.5.8); which one should be clear from the context). This defines a site morphism $\varphi : X_{\text{top}} \rightarrow X$, and we will specialize now the construction of extension and contraction described in Section 1.4 to this situation. We could essentially develop all our topologies as refinements of the Zariski topology, however, this is sometimes unnecessary ballast, in particular in the affine case.

Notation 2.4.1. Let X be a scheme endowed with a Grothendieck topology such that $\varphi : X_{\text{top}} \rightarrow X$ is a site morphism. For an $\mathcal{O}_X$ -module M we denote the module pull-back (the sheafification) by $M_{\text{top}} = \varphi^*M$. Let $N \subseteq M$ denote $\mathcal{O}_X$ -modules. Then the image sheaf of the induced $\mathcal{O}_{X_{\text{top}}}$ -module morphisms $\varphi^*(N) = N_{\text{top}} \rightarrow \varphi^*(M) = M_{\text{top}}$ on X_{top} is called the *extended submodule* of $N \subseteq M$ in M_{top} , denoted by $N^{\text{top}} = N^{\text{ext}}$. In particular, for an ideal sheaf $\mathcal{I} \subseteq \mathcal{O}_X$ we call the image sheaf of $\mathcal{I}_{\text{top}}$ inside $\mathcal{O}_{\text{top}}$ the *extended ideal sheaf*, denoted by $\mathcal{I}^{\text{top}}$.

For an $\mathcal{O}_{\text{top}}$ -submodule $\mathcal{H} \subseteq M_{\text{top}}$ we call the push forward sheaf on X given by $U \mapsto \theta^{-1}(\Gamma(\varphi^*(U), \mathcal{H}))$, where $\theta : \Gamma(U, M) \rightarrow \Gamma(\varphi^*(U), M_{\text{top}})$ is the natural mapping, the *contracted sheaf*, denoted by $\mathcal{H}^{\text{cont}}$.

For $N \subseteq M$ we call the contraction of the extension the *closure sheaf* or *closure submodule* (induced by the topology) of N inside M and denote it by $N^{\text{top-cl}} = (N^{\text{top}})^{\text{cont}}$.

Proposition 2.4.2. *Let X be a scheme endowed with a Grothendieck topology $X_{\text{top}} \rightarrow X$, let $N \subseteq M$ be $\mathcal{O}_X$ -modules. Then the following holds.*

- (i) *The closure submodule $N^{\text{top-cl}}$ is an $\mathcal{O}_X$ -module (but in general not quasicohent).*
- (ii) *We have $N \subseteq N^{\text{top-cl}} = (N^{\text{top-cl}})^{\text{top-cl}}$.*

- (iii) If $\varphi : M \rightarrow M'$ is an $\mathcal{O}_X$ -module homomorphism to another $\mathcal{O}_X$ -module M' and if $N' \subseteq M'$ is a submodule, then $\varphi(N^{\text{top-cl}}) \subseteq (\varphi(N'))^{\text{top-cl}}$.
- (iv) Let $N \subseteq N' \subseteq M$ be another submodule. Then $N^{\text{top-cl}} \subseteq (N')^{\text{top-cl}}$.
- (v) Let $\pi : M \rightarrow M/N$. Then $N^{\text{top-cl}} = \pi^{-1}(0^{\text{top-cl}})$.

Proof. This is a special case of Proposition 1.4.4. $\square$

Remark 2.4.3. We will apply Proposition 2.4.2 mainly in the case of a commutative ring (an affine scheme $X = \text{Spec } R$ endowed with the trivial topology) and an affine Grothendieck topology for R given by a certain class of R -algebras. For R -submodules $N \subseteq M$ we get

$$N \mapsto N^{\text{pre}} \mapsto N_{\text{top}} \mapsto N^{\text{top}} \mapsto \Gamma(X_{\text{top}}, N^{\text{top}}) \mapsto \theta^{-1}(\Gamma(X_{\text{top}}, N^{\text{top}})).$$

We denote this R -submodule of M by $N^{\text{top-cl}}$. For an ideal $I \subseteq R$ we get the assignment

$$I \mapsto I^{\text{pre}} \mapsto I_{\text{top}} \mapsto I^{\text{top}} \mapsto \Gamma(X_{\text{top}}, I^{\text{top}}) \mapsto \theta^{-1}(\Gamma(X_{\text{top}}, I^{\text{top}}))$$

which is an ideal $I^{\text{top-cl}} \subseteq R$. In the case that $N = 0 \subseteq M$, we have $N^{\text{pre}} = N_{\text{top}} = N^{\text{top}} = 0$ and the procedure simplifies to $0^{\text{top-cl}} = \text{kern}(\theta)$, i.e. an element $s \in M$ is in $0^{\text{top-cl}}$ if and only if it is annihilated by the global sheafification homomorphism $\theta : M \rightarrow \Gamma(X_{\text{top}}, M_{\text{top}})$.

The following Lemma describes explicitly the closure operation given by an affine Grothendieck topology.

Lemma 2.4.4. *Let $\mathcal{T}_X$ denote an affine Grothendieck topology on an affine scheme $X = \text{Spec } R$. Let $N \subseteq M$ denote R -modules. Let $s \in M$. Then $s \in N^{\text{top-cl}}$ if and only if there exists a covering $R \rightarrow R_i$, $i \in I$, and elements $t_i \in N \otimes_R R_i$ which map to $s \otimes 1$ under the natural homomorphisms $N \otimes_R R_i \rightarrow M \otimes_R R_i$ for all $i \in I$. If $\mathcal{T}_X$ is quasicompact, then we may restrict to finite index sets.*

Proof. This follows from Lemma 1.4.5 applied to $(X_{\text{top}}, \mathcal{T}_X, \mathcal{O}^{\text{pre}}) \rightarrow X$. Note that in the notation of that Lemma we have $U = X = V = U_i$ for all $i \in I$, $V_i = \text{Spec } R_i$ and $\Gamma(V_i, \mathcal{O}^{\text{pre}}) = R_i$. $\square$

Remark 2.4.5. The condition that $s \in \Gamma(X_{\text{top}}, N^{\text{top}})$ does only mean that s is locally in the image of $N_{\text{top}} \rightarrow M_{\text{top}}$. The property that there exists a (global) element $t \in \Gamma(X_{\text{top}}, N_{\text{top}})$ mapping to s is a stronger condition, because such an element t is represented by compatible elements $t_i \in N \otimes_R R_i$ for some covering $R \rightarrow R_i$, $i \in I$. Also this deserves interest, and there are topologies like the Frobenius topology and the surjective topology where both properties coincide (see Example 3.6.9, Theorem 3.9.1 and Proposition 4.4.1).

We are going to establish finiteness properties of the closure operation induced by a Grothendieck topology.

Lemma 2.4.6. *Let R denote a commutative ring, M an R -module, A an R -algebra. Let $s \in M$ and suppose that $s \otimes 1 = 0$ in $M \otimes_R A$. Then there exists a finitely generated R -submodule $M' \subseteq M$, $s \in M'$, and an R -subalgebra $A' \subseteq A$ of finite type such that $s \otimes 1 = 0$ in $M' \otimes_R A'$.*

Proof. For this we have to go back to the definition of the tensor product, which is constructed by finite sums of formal tensor expressions $v \otimes a$ modulo certain relations. That an element in the tensor product is zero means that it is in the submodule generated by the relations, and for that only finitely many data are needed. $\square$

Corollary 2.4.7. *Suppose that $\mathcal{T}_X$ is a quasicompact Grothendieck topology on a scheme X . Then the closure operation induced by the topology fulfills both finiteness properties from Definition 2.2.14.*

Proof. Suppose that $s \in N^{\text{top-cl}}$, where $N \subseteq M$ are R -modules. By Lemma 2.4.4 there exists a covering $R \rightarrow R_i$, $i \in I$ finite, such that $s \otimes 1 \in M \otimes_R R_i$ is the image of $t_i \in N \otimes_R R_i$.

For the first finiteness property we may assume that $N = 0$. Since $s \in 0^{\text{top-cl}}$ we have that $s \otimes 1 = 0$ in $M \otimes_R R_i$ for every $i \in I$. By Lemma 2.4.6 there exist finitely generated R -submodules $M_i \subseteq M$ such that $s \in M_i$ and such that $s_i = 0$ in $M_i \otimes_R R_i$. Then we can take the finitely generated R -submodule $R\langle M_i, i \in I \rangle \subseteq M$.

For the second finiteness property there exist finitely many elements $y_j \in N$ and $r_j \in R_i$, $j \in J_i$, such that $t_i = \sum_{j \in J_i} y_j \otimes r_j$. This means then that s is already in the closure of the finitely generated submodule $L = R\langle y_j, j \in J_i, i \in I \rangle \subseteq N$. $\square$

Theorem 2.4.8. *Let G denote a subcategory of rings, let $\mathcal{T}_R$ denote an affine Grothendieck topology of finite type on $\text{Spec } R$, $R \in G$, and suppose that every ring homomorphism $R \rightarrow S$ in G induces a site morphism $(\text{Spec } S)_{\text{top}} \rightarrow (\text{Spec } R)_{\text{top}}$. Then the closure operation top-cl induced by the Grothendieck topology is a G -admissible closure operation for all submodules.*

Proof. That top-cl is a closure operation was mentioned in Proposition 2.4.2(ii).

Admissibility. Let $s \in N^{\text{top-cl}}$ and let B denote a forcing algebra (of finite type) for (M, N, s) , where $N \subseteq M$ is a finitely presented R -module and where $s \in N^{\text{top-cl}}$. By Lemma 2.4.4 there exists a covering $R \rightarrow R_i$, $i \in I$, such that $s \otimes 1 \in M \otimes_R R_i$ lies in the image of $N \otimes_R R_i$. Then by the universal property (Proposition 2.1.4) of the forcing algebra we have factorizations $R \rightarrow B \rightarrow R_i$ for all $i \in I$. We have to show that B is top-cl -pure. So let $U \subseteq V$ and $z \in V$ be given and suppose that $z \otimes 1$ is in the image of $U \otimes_R B \rightarrow V \otimes_R B$. Then it is also in the image of $(U \otimes_R B) \otimes_B R_i \cong U \otimes_R R_i$ in $V \otimes_R R_i$ for every $i \in I$, and so $z \in U^{\text{top-cl}}$ by Lemma 2.4.4.

G-persistence. Again, let $N \subseteq M$ and $s \in M$. Let $R \rightarrow S$ denote a ring homomorphism in G with corresponding site morphism $(\text{Spec } S)_{\text{top}} \rightarrow (\text{Spec } R)_{\text{top}}$. The condition $s \in N^{\text{top-cl}}$ means by Lemma 2.4.4 that there exists a covering $R \rightarrow R_i$, $i \in I$, such that there exist $t_i \in N \otimes_R R_i$ mapping to $s \otimes 1$ in $M \otimes_R R_i$. The pull-back $S \rightarrow S_i := R_i \otimes_R S$, $i \in I$, defines a covering of $(\text{Spec } S)_{\text{top}}$. For every $i \in I$ we have the commutative diagram

$$\begin{array}{ccc} N \otimes_R R_i & \longrightarrow & M \otimes_R R_i \\ \otimes_R S \downarrow & & \downarrow \otimes_R S \\ N \otimes_R R_i \otimes_R S & \longrightarrow & M \otimes_R R_i \otimes_R S. \end{array}$$

Note that $N \otimes_R R_i \otimes_R S \cong N \otimes_R S_i \cong (N \otimes_R S) \otimes_S S_i$ (and also for M), so the diagram shows that also $s \otimes 1 \in M \otimes_R S$ has the property that $(s \otimes 1) \otimes 1 \in (M \otimes_R S) \otimes_S S_i$ is in the image of $(N \otimes_R S) \otimes_S S_i$. This means by Lemma 2.4.4 that $s \otimes 1 \in (N_S)^{\text{top-cl}}$, where N_S denotes the image module under $N \otimes_R S \rightarrow M \otimes_R S$. $\square$

As mentioned in Lemma 1.3.6 the sheafification with respect to a Grothendieck topology is right exact. In general it is not left exact (Example 6.2.4). It is however exact for flat topologies and for hereditary torsion theories, see Section 6.3.

Lemma 2.4.9. *If the sheafification in a Grothendieck topology X_{top} is left-exact, then also the closure operation for the 0-submodule is left exact.*

Proof. Let $0 \rightarrow N \rightarrow M \rightarrow L \rightarrow 0$ be an exact sequence of $\mathcal{O}_X$ -modules. Suppose that $s \in 0^{\text{top-cl}} \subseteq \Gamma(U, M)$ maps to 0 in $\Gamma(U, L)$. Then s is the image of an element $t \in \Gamma(U, N)$. Since by assumption $\Gamma(U, N_{\text{top}}) \rightarrow \Gamma(U, M_{\text{top}})$ is injective and $\theta(s) = 0$, it follows that $\theta(t) = 0$ in $\Gamma(U, M_{\text{top}})$, hence $t \in 0^{\text{top-cl}}$ in N . $\square$

An absolute stalk in the affine single-handed case contains the full information about the closure operation. It is however in general not so easy to find an absolute stalk explicitly.

Corollary 2.4.10. *Let $X = \text{Spec } R$ and let an affine single-handed Grothendieck topology $\mathcal{T}_X$ be given on X with induced closure operation top-cl on submodules. Suppose that there exists an absolute filter $F : \Lambda \rightarrow \mathcal{T}_X$ with absolute stalk $R \rightarrow H = \varinjlim_{\lambda \in \Lambda} \Gamma(R_\lambda, \mathcal{O}_{\text{top}})$. Let $N \subseteq M$ denote R -modules, $s \in M$. Then $s \in N^{\text{top-cl}}$ if and only if $s \otimes 1 \in \text{im}(N \otimes_R H \rightarrow M \otimes_R H)$.*

Proof. Suppose that $s \in N^{\text{top-cl}}$. This means by Lemma 2.4.4 that there exists a cover $R \rightarrow R'$ such that $s \otimes 1 \in \text{im}(N \otimes_R R' \rightarrow M \otimes_R R')$. By the definition of an absolute filter, there exists $\lambda \in \Lambda$ and $R' \rightarrow R_\lambda \rightarrow \varinjlim_{\lambda \in \Lambda} R_\lambda = \varinjlim_{\lambda \in \Lambda} \Gamma(R_\lambda, \mathcal{O}_{\text{top}}) = H$ (Corollary 1.9.6). Hence $s \otimes 1 \in \text{im}(N \otimes_R H \rightarrow M \otimes_R H)$. Suppose now on the other hand that this holds. This means that there exist elements $t_j \in N$ and $h_j \in H$, $j \in J$ finite, such that $s \otimes 1 = \sum_{j \in J} h_j t_j$ in $M \otimes_R H$. The h_j are represented in some R_{λ_j} and hence also in one R_λ , $\lambda \in \Lambda$. Since we have a filter, the equality $s \otimes 1 = \sum_{j \in J} h_j t_j$ in the colimit holds already in some covering $R_\lambda \rightarrow R_\kappa$, hence $s \in N^{\text{top-cl}}$ by Lemma 2.4.4. $\square$

2.5. Pure Grothendieck topologies.

If a Grothendieck topology is given on a subcategory G of rings, then there will be certain rings $R \in G$ such that the induced closure operation is trivial, like F -pure rings in the context of Frobenius closure/Frobenius topology or F -regular rings in the context of tight closure. Here we characterize the Grothendieck topologies where the induced closure operation is trivial.

Definition 2.5.1. We call a family of scheme-morphisms $\varphi_i : U_i \rightarrow X$, $i \in I$, *pure* if for every quasicohherent $\mathcal{O}_X$ -module M the natural module homomorphism $\Gamma(X, M) \rightarrow \bigoplus_{i \in I} \Gamma(U_i, \varphi_i^*(M))$ is injective.

We call a Grothendieck topology on a scheme X *pure* if every covering of X in the topology is pure.

Remark 2.5.2. An affine single-handed Grothendieck topology is pure if and only if every cover $R \rightarrow R'$ is a pure homomorphism.

Proposition 2.5.3. *Let X_{top} be an affine scheme endowed with a Grothendieck topology. Then the following are equivalent.*

- (i) *The induced closure operation on submodules is trivial.*
- (ii) *The sheafification homomorphism $\theta : M \rightarrow \Gamma(X_{\text{top}}, M_{\text{top}})$ is injective for every R -module M .*
- (iii) *The Grothendieck topology is pure.*

If the Grothendieck topology is quasicompact, then this is also equivalent to

(iv) *The induced closure operation on submodules of finitely generated modules is trivial.*

Moreover, if X is noetherian, then this is also equivalent to

(v) *The induced closure operation on submodules in artinian modules is trivial.*

Proof. (i) $\Leftrightarrow$ (ii) is clear from Proposition 2.4.2(v). (ii) $\Rightarrow$ (iii). Suppose that $s \in M$ becomes 0 in every $\Gamma(U_i, \varphi_i^* M)$, where $\varphi_i : U_i \rightarrow X$, $i \in I$, is a covering. Then this is also true for the element s considered in $\Gamma(X_{\text{top}}, M_{\text{top}})$. Hence $s = 0$ by (ii).

(iii) $\Rightarrow$ (ii). If $s = 0$ in $\Gamma(X_{\text{top}}, M_{\text{top}})$, then it must be 0 on the presheaf level on some covering (by the definition of sheafification).

The equivalence (i) $\Leftrightarrow$ (iv) follows from Corollary 2.4.7. Suppose now that X is also noetherian, and that (v) holds. Let $s \in M$, a finitely generated module, $s \neq 0$. Then $s \notin \mathfrak{m}^r M$ for some power of some maximal ideal $\mathfrak{m}$. Therefore $\bar{s} \neq 0$ in the artinian module $M/\mathfrak{m}^r M$ and therefore $\bar{s} \notin 0^{\text{top-cl}}$ by (v). Hence $s \notin (\mathfrak{m}^r M)^{\text{top-cl}}$ in M by Proposition 2.2.6(iii) and in particular $s \notin 0^{\text{top-cl}}$ in M by Proposition 2.2.6(i) (or Proposition 2.4.2). $\square$

Remark 2.5.4. In a pure topology the identity $\Gamma(X_{\text{top}}, M_{\text{top}}) = M$ does not hold in general. This is only true if for every object in $\mathcal{T}_X$ the induced topology is pure. For example, for the F -pure rings, i.e. the rings of positive characteristic which are pure in the Frobenius topology (see Proposition 4.2.4 and Theorem 4.3.3) we have $\Gamma(R_{\text{frob}}, \mathcal{O}_{\text{frob}}) = R^\infty$.

Example 2.5.5. In the pure affine single-handed Grothendieck topology only the pure ring homomorphisms $R \rightarrow S$ are covers. Then $M \rightarrow M \otimes_R S$ is by definition injective for all R -modules M . Hence $(M^{\text{pre}})_0 = 0$ and the associated presheaf $M^{\text{pre}} = (M^{\text{pre}})_1$ is already separated. Therefore the closure operation induced by the pure topology is trivial. As soon as an affine single-handed Grothendieck topology encompasses one non-pure covering $R \rightarrow S$ the closure operation is not trivial anymore.

2.6. Grothendieck topologies which induce given closure operations.

In this section we show that every admissible closure operation for ideals (or submodules) comes from the closure operation induced by a suitable (not uniquely determined) affine single-handed Grothendieck topology.

Theorem 2.6.1. *Let G denote a subcategory of commutative rings which is closed under the formation of forcing algebras. Suppose that we have an admissible G -persistent closure operation $N \mapsto N^c$ defined for submodules in a certain category $\mathcal{M}$ of finitely presented modules. Then there exists an affine single-handed Grothendieck topology on every $X = \text{Spec } R$, $R \in G$, such that the induced closure operation is $N^{\text{top-cl}} = N^c$ and such that the homomorphisms $R \rightarrow S$ in G yield site morphisms $(\text{Spec } S)_{\text{top}} \rightarrow (\text{Spec } R)_{\text{top}}$.*

Proof. We look at the following topology on $X = \text{Spec } R$, $R \in G$. Let the objects in $\mathcal{T}_X$ be given by (morphisms which are isomorphic to) compositions

$$X_k \longrightarrow \dots \longrightarrow X_1 \longrightarrow X,$$

where each $X_{i+1} \rightarrow X_i$ is an admissible forcing algebra for the closure operation. We declare such compositions to be the covers in $\mathcal{T}_X$ and the only morphisms.

We have to check that this gives indeed a Grothendieck topology. The identity $X \rightarrow X$ is the forcing algebra of the zero-data. The coverings are of course closed under composition. If $Y \rightarrow X$ is a G -morphism (in particular if $Y \in \mathcal{T}_X$), then we set $Y_i = Y \times_X X_i$ and because of $Y_{i-1} \times_{X_{i-1}} X_i = (Y \times_X X_{i-1}) \times_{X_{i-1}} X_i = Y \times_X X_i = Y_i$ the chain

$$Y_k \rightarrow \dots \rightarrow Y_1 \rightarrow Y$$

is again a composition of forcing algebras over Y . Note that these forcing algebras are again c -admissible due to the G -persistence of the closure operation. Therefore $\mathcal{T}_X$ is closed under products, the covers respect base changes and every G -morphism yields a site morphism.

We have to show that the closure operation top-cl given by this topology is the same as the given closure operation c . Let $N \subseteq M$, $s \in M$, M/N finitely presented. Suppose first that $s \in N^c$. Then the forcing algebra $R \rightarrow B$ for these data is admissible and yields cover (by definition of the topology) such that $s \otimes 1$ is in the image of $N \otimes_R B$. Therefore $s \in N^{\text{top-cl}}$ by Lemma 2.4.4.

On the other hand, if $s \in N^{\text{top-cl}}$, then by Lemma 2.4.4 there exists a covering, that is, a composition

$$R \rightarrow B_1 \rightarrow \dots \rightarrow B_k$$

of admissible forcing algebras, such that $s \otimes 1$ is in the image of $N \otimes_R B_k \rightarrow M \otimes_R B_k$. Since by assumption the admissible forcing algebras are c -pure and since this is true for a composition (Proposition 2.2.4(ii)), it follows that $s \in N^c$. $\square$

Remark 2.6.2. Different topologies may yield the same closure operation. The (somewhat artificial) topology constructed in the proof of Theorem 2.6.1 is the coarsest topology such that the admissible forcing algebras are covers. In general we will deal with other more natural topologies which yield the same closure operation.

Proposition 2.6.3. *Let $\mathcal{D}$ be a subcategory of test rings in the sense of Definition 2.3.2 and let R be a commutative ring. For $X = \text{Spec } R$ let $\mathcal{T}_X$ be the full subcategory of finitely generated R -algebras. Let $(X, \mathcal{T}_X)$ be endowed with an affine single-handed Grothendieck topology, where the covers $\varphi : S \rightarrow S'$ are given by one of the following conditions.*

- (i) *For every ring homomorphism $\psi : S \rightarrow T$, $T \in \mathcal{D}$, the induced ring homomorphism $\varphi_T : T \rightarrow S' \otimes_S T$ has a ring section.*
- (ii) *For every ring homomorphism $\psi : S \rightarrow T$, $T \in \mathcal{D}$, the induced ring homomorphism $\varphi_T : T \rightarrow S' \otimes_S T$ is an direct summand.*
- (iii) *For every ring homomorphism $\psi : S \rightarrow T$, $T \in \mathcal{D}$, the induced ring homomorphism $\varphi_T : T \rightarrow S' \otimes_S T$ is pure (so φ is a $\mathcal{D}$ -pure homomorphisms).*

Then for finitely presented R -modules $N \subseteq M$ we have $N^{\text{top-cl}} = N^{\mathcal{D}\text{-cl}}$. If the closure induced by $\mathcal{D}$ fulfills the first finiteness condition then this holds for arbitrary R -modules.

Proof. Note that the allowed covers are different, but the allowed forcing algebras in the topology are the same under (i), (ii) and (iii) because of Lemma 2.1.5. Hence the three Grothendieck topologies induce the same closure operation, and we restrict to the pure case (iii).

Suppose that $s \in N^{\text{top-cl}}$. This means by Lemma 2.4.4 that there is a cover $R \rightarrow A$ such that $s \otimes 1 \in \text{im}(N \otimes_R A \rightarrow M \otimes_R A)$. This means for $\psi : R \rightarrow T$, $T \in \mathcal{D}$, that $\psi(s) \in \text{im}(N \otimes_R A \otimes_R T \rightarrow M \otimes_R A \otimes_R T)$. Since $T \rightarrow A \otimes_R T$ is pure it follows that already $\psi(s) \in \text{im}(N \otimes_R T \rightarrow M \otimes_R T)$. Hence $s \in N^{\mathcal{D}\text{-cl}}$.

If, on the other hand, $s \in N^{\mathcal{D}-\text{cl}}$, then we consider the (finitely generated) forcing algebra B . It follows that $R \rightarrow B$ is an allowed covering, since after base change to $\psi : R \rightarrow T$ it has even a ring section. Therefore $s \in N^{\text{top-cl}}$ again by Lemma 2.4.4.

The last statement is clear since the topological closure fulfills the finiteness conditions by Corollary 2.4.7 and so the identity reduces to the finite case. $\square$

2.7. Exactness and closure operations.

We relate now for a complex $N \xrightarrow{\alpha} M \xrightarrow{\beta} L$ of R -modules several notions of exactness in a given Grothendieck topology with the property that the kernel is contained in the closure of the image.

Proposition 2.7.1. *Let $\mathcal{T}_X$ denote an affine single-handed Grothendieck topology on $X = \text{Spec } R$. Let $N \xrightarrow{\alpha} M \xrightarrow{\beta} L$ denote a complex of R -modules. Consider the following properties.*

- (i) *The complex of sheaves $N_{\text{top}} \xrightarrow{\alpha} M_{\text{top}} \xrightarrow{\beta} L_{\text{top}}$ on X_{top} is exact.*
- (ii) *The complex of sheaves $N_{\text{top}} \xrightarrow{\alpha} M_{\text{top}} \xrightarrow{\beta} L_{\text{top}}$ on X_{top} is X -exact.*
- (iii) *The complex of presheaves $N^{\text{pre}} \xrightarrow{\alpha} M^{\text{pre}} \xrightarrow{\beta} L^{\text{pre}}$ on X_{top} is X -exact.*
- (iv) *$\text{kern } \beta \subseteq (\text{im } \alpha)^{\text{top-cl}}$, where top-cl denotes the closure operation of the topology (inside M).*

Then (i) $\Rightarrow$ (ii), (ii) $\Rightarrow$ (iii), (iii) and (iv) are equivalent. Moreover, if $M \rightarrow \Gamma(X_{\text{top}}, M_{\text{top}})$ is surjective and $L \rightarrow \Gamma(X_{\text{top}}, L_{\text{top}})$ is injective, then also (ii) is equivalent to (iii) and (iv).

Proof. (i) $\Rightarrow$ (ii) and (ii) $\Rightarrow$ (iii) were given in Proposition 1.2.4.

(iii) $\Rightarrow$ (iv). Let $s \in M$, $\beta(s) = 0$. By (iii) there exists a cover $R \rightarrow S$ and $t \in N \otimes_R S$ mapping to $s \otimes 1$. We may write $t = \sum_{j \in J} t_j \otimes h_j$, $t_j \in N$, $h_j \in S$. Then of course $\alpha(t_j) \in \text{im}(\alpha)$ and so $s \otimes 1 \in \text{im}((\text{im } \alpha) \otimes_R S \rightarrow M \otimes_R S)$, hence $s \in (\text{im } \alpha)^{\text{top-cl}}$.

(iv) $\Rightarrow$ (iii). Again let $s \in M$, $\beta(s) = 0$, so that $s \in (\text{im } \alpha)^{\text{top-cl}}$ by (iv). This means by Lemma 2.4.4 that there exists a cover $R \rightarrow S$ such that $s \otimes 1 \in \text{im}((\text{im } \alpha) \otimes_R S \rightarrow M \otimes_R S)$. So in particular there exists $\tilde{s} = \sum_{j \in J} s_j \otimes h_j$ mapping to $s \otimes 1$, where $s_j = \alpha(t_j)$, $t_j \in N$ and $h_j \in S$. But then also $s \otimes 1 = \alpha(\sum_{j \in J} t_j \otimes h_j)$, so locally s is in the image of the presheaf given by N .

Now suppose that (iii) holds and that $M \rightarrow \Gamma(X_{\text{top}}, M_{\text{top}})$ is surjective and $L \rightarrow \Gamma(X_{\text{top}}, L_{\text{top}})$ is injective. Then $s \in \Gamma(X_{\text{top}}, M_{\text{top}})$ mapping to 0 comes from $s \in M$, which also maps to 0 in L . So there exists a cover $R \rightarrow S$ and $t' \in N \otimes_R S$ mapping to $s \otimes 1$ in $M \otimes_R S$. Then also $t' \in \Gamma(S, N_{\text{top}})$ maps to the restriction of s in $\Gamma(S, M_{\text{top}})$. $\square$

Corollary 2.7.2. *Let $\mathcal{T}_X$ denote an affine single-handed Grothendieck topology on $X = \text{Spec } R$. Let $N \xrightarrow{\alpha} M \xrightarrow{\beta} L$ denote a complex of R -modules. Then the following are equivalent.*

- (i) *The complex of sheaves $N_{\text{top}} \xrightarrow{\alpha} M_{\text{top}} \xrightarrow{\beta} L_{\text{top}}$ on X_{top} is exact.*
- (ii) *The complex of presheaves $N^{\text{pre}} \xrightarrow{\alpha} M^{\text{pre}} \xrightarrow{\beta} L^{\text{pre}}$ on X_{top} is U -exact for every $U \rightarrow X$ in $\mathcal{T}_X$.*

- (iii) For every $R \rightarrow R'$ in $\mathcal{T}_X$ the complex of R' -modules $N \otimes_R R' \xrightarrow{\alpha} M \otimes_R R' \xrightarrow{\beta} L \otimes_R R'$ has the property that $\ker \beta \subseteq (\operatorname{im} \alpha)^{\operatorname{top-cl}}$, where $\operatorname{top-cl}$ denotes the closure operation induced by the topology (inside M).

Proof. Everything follows from Proposition 2.7.1 except (iii) $\Rightarrow$ (i). We have to show that the complex of sheaves is exact for every $U \rightarrow X$ in $\mathcal{T}_X$, and we may assume that $U = X$. Suppose that $s \in \Gamma(X, M_{\operatorname{top}})$ maps to 0 in $\Gamma(X, L_{\operatorname{top}})$, and assume that it is represented by $s \in M \otimes_R R'$, $R \rightarrow R'$ a cover, mapping to 0 in $L \otimes_R R'$. Then by (iii) we have $s \in (\operatorname{im} \alpha)^{\operatorname{top-cl}}$. This means that there exists by Lemma 2.4.4 a cover $R' \rightarrow R''$ such that $s \otimes 1 \in M \otimes_R R''$ is in the image of $N \otimes_R R''$. This means that $s \in \Gamma(X, M_{\operatorname{top}})$ comes locally from the left. $\square$

Corollary 2.7.3. Let $\mathcal{D}$ be a category of test rings and let R be a commutative ring. Let $X = \operatorname{Spec} R$ be endowed with the affine single-handed Grothendieck topology where the covers are given by $\mathcal{D}$ -pure homomorphisms of finite type. Suppose that $\mathcal{T}_X$ contains with every ring R' also the full category of finitely generated R' -algebras. Let $N \xrightarrow{\alpha} M \xrightarrow{\beta} L$ denote a complex of finitely generated R -modules. Then the following are equivalent.

- (i) The complex of sheaves $N_{\operatorname{top}} \xrightarrow{\alpha} M_{\operatorname{top}} \xrightarrow{\beta} L_{\operatorname{top}}$ on X_{top} is exact.
- (ii) For every $R \rightarrow R'$ in $\mathcal{T}_X$ the complex of R' -modules $N \otimes_R R' \xrightarrow{\alpha} M \otimes_R R' \xrightarrow{\beta} L \otimes_R R'$ has the property that $\ker \beta \subseteq (\operatorname{im} \alpha)^{\operatorname{top-cl}}$.
- (iii) For every ring homomorphism $\psi : R \rightarrow T$, $T \in \mathcal{D}$, we have that the complex $N \otimes_R T \xrightarrow{\alpha} M \otimes_R T \xrightarrow{\beta} L \otimes_R T$ is exact.

If the closure induced by $\mathcal{D}$ fulfills the first finiteness condition then this holds for arbitrary R -modules.

Proof. The equivalence of (i) and (ii) was given in Corollary 2.7.2. Suppose that (iii) holds. We may assume that $R = R'$. Suppose that $s \in M$ maps to 0 in L and consider $\operatorname{im}(\alpha) \subseteq M$. Let $\psi : R \rightarrow T$ be given, $T \in \mathcal{D}$. Then by assumption $s \otimes 1 \in \operatorname{im}(N \otimes_R T \rightarrow M \otimes_R T)$ and hence also $s \otimes 1 \in \operatorname{im}(\operatorname{im}(\alpha) \otimes_R T \rightarrow M \otimes_R T)$. But this implies by Lemma 2.6.3 that $s \in (\operatorname{im}(\alpha))^{\operatorname{top-cl}}$.

Suppose now that (ii) holds and let $\psi : R \rightarrow T$ be given, $T \in \mathcal{D}$. Suppose that $u \in M \otimes_R T$ is mapped to 0 in $L \otimes_R T$. There exists by Lemma 2.4.6 a finitely generated R -algebra R' with a factorization $R \rightarrow R' \rightarrow T$ and an element $v \in M \otimes_R R'$ mapping to u and also mapping to 0 in $L \otimes_R R'$. Hence by (ii) $v \in (\operatorname{im}(\alpha))^{\operatorname{top-cl}}$. Therefore by Lemma 2.6.3 we know that $u = v \otimes 1 \in \operatorname{im}(\operatorname{im}(\alpha) \otimes_{R'} T \rightarrow (M \otimes_R R') \otimes_{R'} T)$. Since $N \otimes_R R' \rightarrow \operatorname{im}(\alpha)$ is surjective, it follows also that $u \in \operatorname{im}((N \otimes_R R') \otimes_{R'} T = N \otimes_R T \rightarrow (M \otimes_R R') \otimes_{R'} T = M \otimes_R T)$. $\square$

Example 2.7.4. Let $X = \operatorname{Spec} R$ be an affine scheme endowed with a Grothendieck topology $\mathcal{T}_X$. Let $I = (f_1, \dots, f_n) \subseteq R$ denote an ideal with ideal closure $I^{\operatorname{top-cl}} \neq I$. Then the complex $R^n \xrightarrow{f_1, \dots, f_n} R \rightarrow R/I^{\operatorname{top-cl}}$ is not exact as a complex of R -modules, but the corresponding complex of presheaves is X -exact, because every element in $I^{\operatorname{top-cl}}$ comes from the left under some covering.

Remark 2.7.5. Complexes of R -modules with the property that the kernel is inside the closure of the image (as in Proposition 2.7.1(iv)) were studied by several people with

respect to different closure operations. Katz studies complexes acyclic up to integral closure [41] and Hochster and Huneke study phantom homology within the setting of tight closure [38, §10].

An exact sequence of R -modules leads in general not to an exact sequence of sheaves. There are however some cases when they do.

Proposition 2.7.6. *Let R denote a commutative ring, $0 \rightarrow N \rightarrow M \rightarrow L \rightarrow 0$ an exact sequence of R -modules and let an affine single-handed Grothendieck topology be given on $X = \operatorname{Spec} R$. Then the complex of sheaves $0 \rightarrow N_{\text{top}} \rightarrow M_{\text{top}} \rightarrow L_{\text{top}} \rightarrow 0$ is exact in the following cases.*

- (i) *The sequence splits.*
- (ii) *L is flat.*
- (iii) *N is pure in M .*

Proof. We only have to show that the corresponding complex of presheaves is exact. (i) is clear, since the presheafification respects direct sums. (ii),(iii). For $R \rightarrow R'$ we have the exact sequence of presheaves

$$\rightarrow \operatorname{Tor}_1^R(L, R') \rightarrow N \otimes_R R' \rightarrow M \otimes_R R' \rightarrow L \otimes_R R' \rightarrow 0.$$

If L is flat, then $\operatorname{Tor}_1^R(L, R') = 0$, and the presheaf complex is exact. If $N \rightarrow M$ is pure, then again $N \otimes_R R' \rightarrow M \otimes_R R'$ is injective. $\square$

2.8. Sheaf of syzygies and cohomological interpretation.

We want to relate the quotient $N^{\text{top-cl}}/N$ with the first cohomology of the syzygy sheaf in the topology. We consider finitely generated R -modules $N \subseteq M$ and a homomorphism $\sigma : R^n \rightarrow M$ such that the image of σ is N . Let a Grothendieck topology $\mathcal{T}_X$ on $X = \operatorname{Spec} R$ be given. This yields a surjective homomorphism of sheaves

$$\sigma : \mathcal{O}_{\text{top}}^n \longrightarrow N^{\text{top}} \subseteq M_{\text{top}}.$$

The kernel of this sheaf homomorphism is a sheaf on X_{top} which we denote by Syz . Note that this kernel sheaf is not the sheafification of the kernel of the given R -module homomorphisms.

Proposition 2.8.1. *Let R denote a commutative ring with a Grothendieck topology $\mathcal{T}_X$ on $X = \operatorname{Spec} R$. Let $N \subseteq M$ denote finitely generated R -modules and let $R^n \rightarrow M$ denote an R -module homomorphism with image N . Let Syz denote the kernel sheaf of the sheaf morphism $\mathcal{O}_{\text{top}}^n \rightarrow N^{\text{top}} \subseteq M_{\text{top}}$. Then the following holds.*

- (i) *We have a homomorphism $N^{\text{top-cl}}/N \rightarrow H^1(X_{\text{top}}, \operatorname{Syz})$. If $\Gamma(X_{\text{top}}, \mathcal{O}_{\text{top}}) = R$, $\Gamma(X_{\text{top}}, M_{\text{top}}) = M$ and $H^1(X_{\text{top}}, \mathcal{O}_{\text{top}}) = 0$, then this is an isomorphism.*
- (ii) *Suppose that $U \subseteq X$ is Zariski open, that $\mathcal{T}_X$ is a refinement of the Zariski topology and that the support of M/N is outside of U . Set $Y = X - U$. Then we have a*

commutative diagram

$$\begin{array}{ccccc}
N^{\text{top-cl}}/N & \longrightarrow & M/N & \longrightarrow & M/N^{\text{top-cl}} \\
\downarrow & & \downarrow & & \downarrow \\
\Gamma(X_{\text{top}}, N^{\text{top}})/N\Gamma(X_{\text{top}}, \mathcal{O}_{\text{top}}) & \longrightarrow & \Gamma(U_{\text{top}}, N^{\text{top}})/N\Gamma(U_{\text{top}}, \mathcal{O}_{\text{top}}) & \longrightarrow & H_Y^1(N^{\text{top}})/\text{im}(H_Y^1(\mathcal{O}_{\text{top}}^n)) \\
\downarrow \subseteq & & \downarrow \subseteq & & \downarrow \subseteq \\
H^1(X_{\text{top}}, \text{Syz}) & \longrightarrow & H^1(U_{\text{top}}, \text{Syz}) & \longrightarrow & H_{Y_{\text{top}}}^2(\text{Syz}).
\end{array}$$

Proof. We apply to the short exact sequence of sheaves on X_{top} ,

$$0 \longrightarrow \text{Syz} \longrightarrow \mathcal{O}_{\text{top}}^n \longrightarrow N^{\text{top}} \longrightarrow 0$$

several cohomological functors, to wit, $\Gamma(X_{\text{top}}, -)$, $\Gamma(U_{\text{top}}, -)$ and $\Gamma_{Y_{\text{top}}}(-)$, and get the commutative diagram

$$\begin{array}{ccccc}
\Gamma(X_{\text{top}}, \mathcal{O}_{\text{top}})^n & \longrightarrow & \Gamma(U_{\text{top}}, \mathcal{O}_{\text{top}})^n & \longrightarrow & H_Y^1(\mathcal{O}_{\text{top}})^n \\
\downarrow & & \downarrow & & \downarrow \\
\Gamma(X_{\text{top}}, N^{\text{top}}) & \longrightarrow & \Gamma(U_{\text{top}}, N^{\text{top}}) & \longrightarrow & H_Y^1(N^{\text{top}}) \\
\downarrow & & \downarrow & & \downarrow \\
H^1(X_{\text{top}}, \text{Syz}) & \longrightarrow & H^1(U_{\text{top}}, \text{Syz}) & \longrightarrow & H_Y^2(\text{Syz}) \\
\downarrow & & \downarrow & & \downarrow \\
H^1(X_{\text{top}}, \mathcal{O}_{\text{top}})^n & \longrightarrow & H^1(U_{\text{top}}, \mathcal{O}_{\text{top}})^n & \longrightarrow & H_Y^2(\mathcal{O}_{\text{top}})^n.
\end{array}$$

Note that the image of $\Gamma(X_{\text{top}}, \mathcal{O}_{\text{top}})^n$ in $\Gamma(X_{\text{top}}, N^{\text{top}})$ is just $N\Gamma(X_{\text{top}}, \mathcal{O}_{\text{top}})$ (same for the second column). From this everything in the lower half of the diagram in (ii) follows.

The first down arrow in the diagram in (ii) on the left comes from the sheafifying homomorphism $\theta : M \rightarrow \Gamma(X_{\text{top}}, M_{\text{top}})$, where by definition $N^{\text{top-cl}}$ is the submodule mapping to $\Gamma(X_{\text{top}}, N^{\text{top}}) \subseteq \Gamma(X_{\text{top}}, M_{\text{top}})$. This gives the first statement in (i). The other statement in (i) follows from the long exact sequence (in the first column), since $\Gamma(X_{\text{top}}, M_{\text{top}}) = M$ implies that $\Gamma(X_{\text{top}}, N^{\text{top}}) = N^{\text{top-cl}}$.

Suppose now that the support of M/N is outside U , or equivalently that $R^n \rightarrow M$ is surjective on U . Then we have $N^{\text{top}} = M_{\text{top}}$ on U_{top} . The mapping $M \rightarrow \Gamma(X_{\text{top}}, M_{\text{top}}) \xrightarrow{\rho} \Gamma(U_{\text{top}}, M_{\text{top}}) = \Gamma(U_{\text{top}}, N^{\text{top}})$ induces the first mapping in the second column and the upper left square commutes.

The induced mapping $M/N \rightarrow H_Y^1(N^{\text{top}})/\text{im}(H_Y^1(\mathcal{O}_{\text{top}})^n)$ sends $N^{\text{top-cl}} \mapsto 0$, so it factors through $M/N^{\text{top-cl}}$. $\square$

Example 2.8.2. We describe explicitly how a class $[s] \in N^{\text{top-cl}}/N$ gives a cohomology class $c = \delta([s]) \in H^1(X_{\text{top}}, \text{Syz})$ in the easiest case, that of a principal ideal $I = (g)$ in an affine single-handed Grothendieck topology on $X = \text{Spec } R$. For $U \in \mathcal{T}_X$ we have the sequence $0 \rightarrow \Gamma(U, \text{Syz}) \rightarrow \Gamma(U, \mathcal{O}_{\text{top}}) \xrightarrow{g} \Gamma(U, I^{\text{top}})$, so $\Gamma(U, \text{Syz}) = \{h \in \Gamma(U, \mathcal{O}_{\text{top}}) : hg = 0\} = \text{Ann}(g)$, which is zero if g is a non-zero divisor in $\Gamma(U, \mathcal{O}_{\text{top}})$. If $f \in R$ and

$f \in I^{\text{top-cl}}$, then there exists a cover $Y \rightarrow X$ and an element $q \in \Gamma(Y, \mathcal{O}_{\text{top}})$ such that $f = qg$. Therefore over $Y \times_X Y$ the element $p_2^*(q) - p_1^*(q)$ maps to 0 in $\Gamma(Y \times_X Y, I^{\text{top}})$ and comes from an element in $\Gamma(Y \times_X Y, \text{Syz})$. This element is a Čech representative of $\delta(f) \in H^1(X, \text{Syz})$.

The Čech cohomology of the syzygy sheaf (which is in our case an annihilator sheaf of one element) with respect to a cover $\check{H}^1(Y \rightarrow X, \text{Syz})$ is computed as the homology of the Čech complex $\Gamma(X, \text{Syz}) \rightarrow \Gamma(Y, \text{Syz}) \rightarrow \Gamma(Y \times_X Y, \text{Syz}) \rightarrow \Gamma(Y \times_X Y \times Y, \text{Syz})$. We compute an example in the finite topology in Example 5.4.3.

Remark 2.8.3. It is possible that two syzygy modules are isomorphic as R -modules and even isomorphic to the Zariski structure sheaf, but that they define different sheaves of syzygies in a Grothendieck topology. This is for example the case in the submersive topology [4] for $\text{Syz}(x, y)$ and $\text{Syz}(x^2, y^2)$ on $X = \text{Spec } K[x, y]$, which are both Zariski-isomorphic to $\mathcal{O}_X$. This is related to the fact that the containment $f \in \overline{(f_1, \dots, f_n)}$ in the integral closure is not a property of the Zariski cohomology class $\delta(f) \in H^1(D(f_1, \dots, f_n), \text{Syz}(f_1, \dots, f_n))$. For example, both data sets $(x, y); 1$ and $(x^2, y^2); xy$ yield the same Zariski cohomology class $1/xy \in H^1(D(x, y), \mathcal{O}_X)$, but $1 \notin \overline{(x, y)}$ and $xy \in \overline{(x^2, y^2)}$. This is also related to the fact that there exists no theory of test ideals for the integral closure.

3. THE SURJECTIVE TOPOLOGY AND THE RADICAL

In this part we describe our first main example, the radical of an ideal and its relationship to the surjective Grothendieck topology, where the coverings are given by affine spec-surjective morphisms.

3.1. The radical of an ideal and of a submodule.

The radical of an ideal $I \subseteq R$ is by definition

$$\sqrt{I} = \text{rad}(I) = \{f \in R : \exists n \in \mathbb{N} \text{ such that } f^n \in I\}.$$

The radical is also the intersection of all prime ideals which contain I [12, Corollary 2.12]. The containment $f \in \text{rad}(I)$ is thus equivalent with $\psi(f) \in IK$ for all ring homomorphisms $\psi : R \rightarrow K$ to fields.

Proposition 3.1.1. *Let $N \subseteq M$ denote R -modules such that M/N is finitely presented, let $s \in M$. Then the following are equivalent.*

- (i) *There exists a spec-surjective ring homomorphism $R \rightarrow A$ (of finite type) such that $s \otimes 1 \in \text{im}(N \otimes_R A \rightarrow M \otimes_R A)$.*
- (ii) *The forcing algebra B for (M, N, s) induces a surjective morphism $\text{Spec } B \rightarrow \text{Spec } R$.*
- (iii) *For all homomorphisms $R \rightarrow K$ to a field K we have $s \otimes 1 \in \text{im}(N \otimes_R K \rightarrow M \otimes_R K)$.*

Proof. (i) $\Rightarrow$ (ii). By the universal property of the forcing algebra (Proposition 2.1.4) we have a factorization $\text{Spec } A \rightarrow \text{Spec } B \rightarrow \text{Spec } R$, so also $\text{Spec } B \rightarrow \text{Spec } R$ is surjective (and B is of finite type). (ii) $\Rightarrow$ (i) is clear.

(ii) $\Leftrightarrow$ (iii). By the functoriality of forcing algebras (Remark 2.1.3) $B \otimes_R K$ is the forcing algebra for $(M \otimes_R K, N', s \otimes 1)$, N' being the image of $N \otimes_R K$ in $M \otimes_R K$. (ii) says that every fiber is non-empty, and this is equivalent to the property that $B \otimes_R K \neq 0$ for every

$R \rightarrow K$. This is equivalent to $K \rightarrow B \otimes_R K$ being a direct summand, hence this is by Lemma 2.1.5 equivalent to (iii). $\square$

Definition 3.1.2. For R -modules $N \subseteq M$ we define the radical of N in M to be the submodule

$$\text{rad}(N) = \{s \in M : s \otimes 1 \in \text{im}(N \otimes_R K \rightarrow M \otimes_R K) \text{ for all fields } R \rightarrow K\}.$$

Remark 3.1.3. This definition gives clearly a submodule. If N and M are such that M/N is finitely presented, then the containment $s \in \text{rad}(N)$ can be expressed by the equivalent properties of Proposition 3.1.1. The radical is by Proposition 2.3.4 an admissible closure operation with respect to every ring homomorphism. This follows also from the topological interpretation in Proposition 3.2.3 below.

Remark 3.1.4. Let M denote an R -module and consider the 0-submodule. Then the radical of 0 consists of all elements $s \in M$ such that $s \otimes 1 = 0$ in $M \otimes_R K$ for all fields $R \rightarrow K$.

Proposition 3.1.5. Suppose that R is a noetherian ring of finite dimension. Then the radical fulfills both finiteness properties from Definition 2.2.14.

Proof. Let $N \subseteq M$ be R -modules, $s \in M$ and $s \in \text{rad}(N)$. For the first finiteness property we may assume that $N = 0$. Let $\mathfrak{p}_i, i \in I$, denote the finitely many minimal prime ideals of R . For all i we have $s \otimes 1 = 0$ in $M \otimes_R Q(R/\mathfrak{p}_i)$. Then there exist by Lemma 2.4.6 finitely generated R -submodules $M_i \subseteq M$ such that $s \in M_i$ and $s \otimes 1 = 0$ in $M_i \otimes_R Q(R/\mathfrak{p}_i)$. The same holds for the finitely generated submodule $\tilde{M}$ generated by the M_i . This means that there exist $f_i \notin \mathfrak{p}_i$ such that $s \otimes 1 = 0$ in $\tilde{M} \otimes_R (R/\mathfrak{p}_i)_{f_i}$. Now if $R \rightarrow K$ is a homomorphism to a field K , then either f_i maps to a unit for some i or all f_i map to 0. In the first case the factorization $R \rightarrow R_{f_i} \rightarrow K$ shows that $s \otimes 1 = 0$ in $\tilde{M} \otimes_R K$. In the second case we look at the minimal primes of $R/(f_i, i \in I)$ as before, which has smaller dimension, so we can (increasing the submodule $\tilde{M}$) argue inductively.

The proof of the second finiteness condition is similar. $\square$

Definition 3.1.6. For an R -module M we denote the factor module by $M_1 = M/\text{rad}(0)$ and call it the R -reduction of M .

The R -reduction of an R -module M yields indeed the separated presheaf in the surjective topology, as we will see in a minute.

Example 3.1.7. Let R denote a principal ideal domain. Then every finitely generated R -module M is isomorphic to $M \cong R^k \times M(p_1) \times \dots \times M(p_m)$, where $p_1, \dots, p_m$ are non associated prime elements in R and where $M(p_i) = (R/p_i)^{u_{i1}} \times (R/p_i^2)^{u_{i2}} \times \dots \times (R/p_i^{n_i})^{u_{in_i}}$ is the p_i -primary component of M , see [62, Satz 61.8 and Satz 61.10]. The radical of 0 is then given by the direct product (Proposition 2.2.6(iv)) of the radicals of 0 in R^k (which is 0) and in $M(p_i)$. The radical of 0 in $(R/p)^{u_1} \times (R/p^2)^{u_2} \times \dots \times (R/p^n)^{u_n}$ is $0 \times (pR/p^2)^{u_2} \times \dots \times (pR/p^n)^{u_n}$ and the R -reduction is $(R/p)^{u_1+u_2+\dots+u_n}$.

Example 3.1.8. Let $R = K[x, y]$ (K a field) and let's consider the R -module $M = R^2/((x, y), (x^2y, x))$. We claim that the residue class s of $(x(1 - y^2), 0)$ is not 0 in M , but it is in the radical of 0. The determinant of the generators is $x^2(1 - y^2)$. If $R \rightarrow L$ is a homomorphism to a field, and if the determinant (in L) is $\neq 0$, then $M_L = 0$. If the determinant is 0, then also $s = 0$, so $s \in \text{rad}(0)$. Assume now that $(x(1 - y^2), 0) =$

$f(x, y) + g(x^2y, x)$ with $f, g \in R$. Then $f = hx$ and $g = -hy$ by the second column and the first gives $x(1 - y^2) = hx^2 - hx^2y^2$, which has no solution.

3.2. Surjective topology.

Definition 3.2.1. Let X be a scheme and let $\mathcal{T}_X$ be the full subcategory of schemes (or schemes of finite type) which are affine over X . The coverings in the (affine-) *surjective topology* on $\mathcal{T}_X$ are the affine surjections $V \rightarrow U$ of finite type (over X). We denote X endowed with this topology by X_{surj} .

Proposition 3.2.2. *The surjective topology is an affine single-handed Grothendieck topology, and it is a refinement of the single-handed Zariski-topology. Every scheme-morphism $Y \rightarrow X$ induces a site morphism $Y_{\text{surj}} \rightarrow X_{\text{surj}}$.*

Proof. This follows immediately from the definitions. The last statement follows since surjections are universally surjective. $\square$

Proposition 3.2.3. *Let R denote a ring and let $N \subseteq M$ denote R -modules such that M/N is finitely presented. Then the radical of N is the closure operation induced by the surjective topology. If R is noetherian of finite dimension, then this is true for arbitrary submodules $N \subseteq M$.*

Proof. The first statement follows from Proposition 3.1.1 and Lemma 2.4.4. The second statement follows from this, from Corollary 2.4.7 and from Proposition 3.1.5. $\square$

We are going to describe the sheafification in the surjective topology, in particular for the structure sheaf. We start with the separated presheaf.

Lemma 3.2.4. *Let R denote a commutative ring, M a finitely generated R -module yielding the presheaf $R' \mapsto M \otimes_R R'$ for $R \rightarrow R'$ of finite type. Then $(M \otimes_R R')_0 = \text{rad}(0)$ and the R' -reduction $(M \otimes_R R')_1 = M \otimes_R R' / (M \otimes_R R')_0$ is the associated separated presheaf.*

Proof. This follows from Proposition 3.1.1 and Section 1.5. $\square$

Corollary 3.2.5. *Let R denote a commutative ring endowed with the surjective topology. Then the separated presheaf $\mathcal{O}_1$ associated to the Zariski structure sheaf is the reduction, i.e. $\Gamma(R', \mathcal{O}_1) = R'_{\text{red}}$.*

Proof. This follows from Lemma 3.2.4. $\square$

Remark 3.2.6. It follows that an element $f \in S$, $R \rightarrow S$ spec-surjective, yields a global section in $\Gamma(R, \mathcal{O}_{\text{surj}})$ if and only if $\delta(f) = f \otimes 1 - 1 \otimes f \in S \otimes_R S$ is nilpotent. We will compute the ring of global sections in Section 3.8.

3.3. The Jacobson radical and the Jacobson topology.

In this section we describe briefly a variant of the radical and of the surjective topology, the Jacobson radical and the Jacobson topology. The *Jacobson radical* of a commutative ring R is by definition the intersection of all maximal ideals [12, Chapter 4]. Accordingly, we make the following definition.

Definition 3.3.1. The *Jacobson radical* of a submodule $N \subseteq M$ is

$$\text{Jac}(N) = \{s \in M : s \otimes 1 \in \text{im}(N \otimes_R \kappa(\mathfrak{m}) \rightarrow M \otimes_R \kappa(\mathfrak{m})) \text{ for all } \mathfrak{m} \text{ maximal}\}.$$

Remark 3.3.2. Note that this closure operation is not persistent with respect to all ring homomorphisms. It is however persistent with respect to homomorphisms between K -algebras of finite type over a field K , in which case it is the same as the radical, see Proposition 3.3.5. For $0 \subseteq M$ we have $\text{Jac}(0) = \bigcap_{\mathfrak{m} \text{ maximal}} \mathfrak{m}M$ (note that for the radical it is not true that $\text{rad}(0) = \bigcap \mathfrak{p}M$) and for N we have $\text{Jac}(N) = \bigcap_{\mathfrak{m} \text{ maximal}} (N + \mathfrak{m}M)$.

Definition 3.3.3. Let X be a scheme and let $\mathcal{T}_X$ be the subcategory of schemes of finite type which are affine over X . The *Jacobson topology* on X is the Grothendieck topology on $\mathcal{T}_X$, where $\varphi : V \rightarrow U$ is a covering if φ maps onto the closed points of U . We denote X endowed with this topology by X_{Jac} .

Remark 3.3.4. The Jacobson topology is a refinement of the surjective topology. For $X = \text{Spec } R$, a morphism $\text{Spec } S \rightarrow \text{Spec } R$ is a covering if and only if every maximal ideal of R lies in the image. For a local ring R , the morphism $\text{Spec } \kappa(\mathfrak{m}) \rightarrow \text{Spec } R$ is a covering, and more generally, if $I \subseteq \text{Jac}$, then $\text{Spec } R/I \rightarrow \text{Spec } R$ is a covering in the Jacobson topology.

A scheme morphism $\psi : Y \rightarrow X$ which sends closed points to closed points induces a site morphism between the Jacobson sites. This holds in particular for morphisms between schemes of finite type over a field [20, Corollaire 10.4.7]. The closure operation induced by the Jacobson topology is the Jacobson radical. Also, $\text{rad}(N) \subseteq \text{Jac}(N)$.

Recall that a commutative ring is called a *Jacobson ring* if every prime ideal is the intersection of maximal ideals [20, Définition 10.4.1].

Proposition 3.3.5. *For a noetherian Jacobson ring R and for finitely generated R -modules $N \subseteq M$ the identity $\text{Jac}(N) = \text{rad}(N)$ holds.*

Proof. Let $R \rightarrow S$ be of finite type and suppose that all maximal ideals of R are in the image of $\text{Spec } S \rightarrow \text{Spec } R$. We want to show that the morphism is surjective. For this we may assume that R is a domain and that $\mathfrak{p} = 0$. The closure of the image is $\text{Spec } R$ due to the Jacobson property. Hence by the Theorem of Chevalley ([20, Théorème 1.8.4] or [12, Corollary 14.7] the image contains a non-empty set and in particular it contains 0. This shows that a Jacobson cover is spec-surjective. The stated result follows then from Remark 3.3.4 and Proposition 3.2.3. $\square$

Definition 3.3.6. We say that a Grothendieck topology $\varphi : X_{\text{top}} \rightarrow X$ on a scheme X is *faithful* (or has the *Nakayama property*) if for every coherent $\mathcal{O}_X$ -module $\mathcal{F}$ the following holds: if $\varphi^*(\mathcal{F}) = \mathcal{F}_{\text{top}} = 0$, then $\mathcal{F} = 0$.

The following Lemma is a version of the Lemma of Nakayama [12, Corollary 4.8].

Lemma 3.3.7. *The Jacobson topology (and hence the surjective topology) on a noetherian scheme X is faithful.*

Proof. We may assume that $X = \text{Spec } R$ is affine. Let M be a finitely generated R -module and suppose that $M_{\text{Jac}} = 0$. This means that for every $s \in M$ there exists a Jacobson cover $\text{Spec } S \rightarrow \text{Spec } R$ such that $s \otimes 1 = 0$ in $M \otimes_R S$. Since M is finitely generated, there exists also such an S with $M \otimes_R S = 0$. Since $R \rightarrow S$ is a Jacobson cover, for every maximal ideal $\mathfrak{m} \subset R$ the homomorphism $R/\mathfrak{m} \rightarrow S/\mathfrak{m}S$ is faithful, and so $M \otimes_R R/\mathfrak{m} = M/\mathfrak{m}M = 0$ for every maximal ideal. But then $M_{\mathfrak{m}} = 0$ by the Lemma of Nakayama and so $M = 0$. $\square$

Corollary 3.3.8. *Let R denote a noetherian commutative ring and $N \subset M$ a submodule such that $M/N \neq 0$ is finitely generated. Then $\text{Jac}(N) \neq M$ and $\text{rad}(N) \neq M$.*

Proof. This follows from Lemma 3.3.7. $\square$

Example 3.3.9. Corollary 3.3.8 is not true for modules which are not finitely generated. If V is a discrete valuation domain, then the quotient module $Q(V)/V$ has the property that its tensoration both with the residue class field $V/\mathfrak{m}_V$ and with $Q(V)$ is zero, and hence the tensoration with every field. Hence the radical of 0 in this module is the whole module.

Remark 3.3.10. Let R be a principal ideal domain with infinitely many maximal ideals (like $\mathbb{Z}$). Then by Proposition 3.3.5 for a finitely generated R -module the radical is the same as the Jacobson radical. This is not true for not finitely generated modules, and the Jacobson radical does not fulfill the finiteness property 2.2.14(i). For example, for the quotient field $Q = Q(R)$ the radical of 0 is 0, since Q itself is a test ring, but the Jacobson radical of 0 is Q , since $Q \otimes_R R/\mathfrak{m} = 0$ for every maximal ideal $\mathfrak{m}$. Every finitely generated submodule $M' \subseteq Q$ is isomorphic to R^k , and there the Jacobson radical is 0.

Lemma 3.3.11. *If X is a scheme of finite type over an algebraically closed field K , and $N \subseteq M$ are coherent $\mathcal{O}_X$ -modules, then $s \in \text{rad}(N)$ if and only if this is true for all K -points of X .*

Proof. The surjectivity of the map from the forcing scheme to X can be tested on K -points alone. $\square$

Example 3.3.12. Let X be a scheme of finite type over $\mathbb{C}$ and consider the ringed site X_{arb} given by the point set $X(\mathbb{C})$ (with the complex topology) and the sheaf of all (not necessarily continuous) complex-valued functions on $X(\mathbb{C})$. The natural mapping $X_{\text{arb}} \rightarrow X_{\text{Zar}}$ is a morphism of ringed sites (in fact, of ringed spaces). We claim that the closure operation induced by this morphism (it is not a Grothendieck topology on X) equals the radical. This follows from the simple fact that $Y \rightarrow X$ is surjective if and only if $Y(\mathbb{C}) \rightarrow X(\mathbb{C})$ is surjective, and the later is equivalent to the existence of a (not necessarily continuous) section $X(\mathbb{C}) \rightarrow Y(\mathbb{C})$. Applying this fact to the forcing schemes $\psi : Y \rightarrow X$ gives the claim.

We relate briefly our notion of the radical of a submodule with another one found in the literature, see [47], [46], [54], [44] (there are also more abstract categorical definitions of a radical). A submodule $P \subseteq M$ of an R -module is called a *prime submodule* if whenever $fs \in P$ (where $f \in R$, $s \in M$), then $s \in P$ or $fM \subseteq P$. Maximal submodules $P \subseteq M$, which are characterized by $M/P \cong R/\mathfrak{m}$ for some maximal ideal $\mathfrak{m}$, are prime. For an arbitrary submodule $N \subseteq M$ one defines the (prime)-radical by $\widetilde{\text{rad}}(N) = \bigcap_{N \subseteq P \text{ prime}} P$.

Proposition 3.3.13. *Let R denote a noetherian Jacobson ring and let M denote a finitely generated R -module, let $N \subseteq M$ be a submodule. Then*

$$\text{rad}(N) = \text{Jac}(N) = \bigcap_{\mathfrak{m} \text{ maximal ideal}} (N + \mathfrak{m}M) = \bigcap_{N \subseteq P \text{ maximal}} P \supseteq \widetilde{\text{rad}}(N).$$

In particular, $\widetilde{\text{rad}}(N) \subseteq \text{rad}(N)$.

Proof. The first equation is Proposition 3.3.5, the second is the definition (Remark 3.3.2) and the forth inclusion is trivial, so we only have to look at the third equation. We may assume that $N = 0$. Assume first that $s \in \text{Jac}(N)$ and let $P \subseteq M$ be a maximal submodule. Then $M/P \cong R/\mathfrak{m}$ as R -modules for some maximal ideal $\mathfrak{m} \subset R$ and so M/P is annihilated by $\mathfrak{m}$. Then $\mathfrak{m}M \subseteq P$ and so $s \in P$.

Suppose now that $s \in P$ for all maximal submodules $P \subseteq M$, and let $\mathfrak{m}$ be a maximal ideal of R . Then $M/\mathfrak{m}M$ is a finitely generated $R/\mathfrak{m} = K$ -module, hence isomorphic to K^n (we may assume $n \geq 1$). The maximal submodules of K^n are the subspaces of dimension $n - 1$, and their intersection is 0. Pulling this back to M we see that $\mathfrak{m}M$ is the intersection of maximal submodules, hence $s \in \mathfrak{m}M$. $\square$

3.4. The surjective topology over a field.

We consider the case of a field K . Although an algebra $K \rightarrow A$ is spec-surjective if and only if it is faithfully flat (if and only if it is $\neq 0$), the surjective topology is not the same as the faithfully flat topology. This is because the compatibility condition in the surjective topology takes into account the reduction of $A \otimes_K A$. Recall that the *perfect closure* of a field K in characteristic zero is K itself and $K^\infty = \varinjlim_e {}^e K$ in positive characteristic, see also Part 4.

Proposition 3.4.1. *Let K denote a field endowed with the surjective topology. Then $\Gamma(K_{\text{surj}}, \mathcal{O}_{\text{surj}}) = K^\infty$, the perfect closure of K . If K is a perfect field, then $\Gamma(K_{\text{surj}}, \mathcal{O}_{\text{surj}}) = K$.*

Proof. Theorem 4.3.3 below shows that an element in the perfect closure K^∞ gives an element in $\Gamma(K_{\text{surj}}, \mathcal{O}_{\text{surj}})$. Therefore we may assume that K is perfect and we have to prove the second statement.

Let $K \rightarrow A$ denote an K -algebra $A \neq 0$ of finite type. We may assume that A is a domain, since any integral component yields a refined covering. Suppose that an element $f \in A$ is compatible, that is, $\delta(f) = f \otimes 1 - 1 \otimes f$ is nilpotent in $A \otimes_K A$. By [20, Proposition 4.6.1] this algebra is reduced, hence $\delta(f) = 0$ in $A \otimes_K A$. Therefore f is also compatible in the faithfully flat topology, hence $f \in K$ by [49, Proposition I.2.18]. $\square$

3.5. Absolute stalks in the surjective topology.

Let R be a ring and set $\text{rad}(R) = \prod_{x \in \text{Spec } R} \overline{\kappa(x)}$ (we call it the *radicalization* of R). The homomorphisms $R \rightarrow R_{\mathfrak{p}_x} \rightarrow \kappa(x) \rightarrow \overline{\kappa(x)}$ induce a homomorphism $R \rightarrow \text{rad}(R)$.

Proposition 3.5.1. *Let R denote a commutative ring, considered in the surjective topology. Then the fixation given by $R \rightarrow \text{rad}(R)$ defines an absolute filter and the radicalization is an absolute stalk.*

Proof. Let $R \rightarrow S$ in $\mathcal{T}_X$ be given with a fixation $S \rightarrow \text{rad}(R)$. Then $\text{Spec } S \rightarrow \text{Spec } R$ is surjective, hence a cover. Let $S \rightarrow T$ be a cover in the surjective topology. We have to show that there exists an S -algebra homomorphism $T \rightarrow \text{rad}(R)$, and this can be checked for every component $\overline{\kappa(x)}$ separately. For $x \in \text{Spec } R$ the geometric fiber ring $T_x := T \otimes_S \overline{\kappa(x)} \neq 0$ is of finite type over $\overline{\kappa(x)}$, hence there exists a homomorphism $T_x \rightarrow \overline{\kappa(x)}$. So this quasifilter is absolute. For $S \rightarrow \text{rad}(R)$ the image $\overline{S} = \text{im}(S) \subseteq \text{rad}(R)$ is also (spec-surjective and) of finite type over R , so $S \rightarrow \overline{S}$ belongs to $\mathcal{T}_X$. Hence by Proposition 1.8.6 this is a filter. Since $R \rightarrow R[U]$ is a spec-surjective covering, the corresponding homomorphism $\mathcal{O}_F = \varinjlim_{\lambda \in \Lambda} S_\lambda \rightarrow \text{rad}(R)$ is an isomorphism. $\square$

Example 3.5.2. Let V denote a discrete valuation domain. Then by Proposition 3.4.1 and Corollary 3.6.6 we have $\Gamma(V, \mathcal{O}_{\text{surj}}) = Q(V)^\infty \oplus (V/\mathfrak{m})^\infty =: K \oplus k$. An ideal $I = (\pi^n)$ extends in $\Gamma(V, \mathcal{O}_{\text{surj}})$ to the ideal $K \oplus k$ for $n = 0$ and to $K \oplus 0$ for $n \geq 1$. The zero-ideal in V defines the zero ideal. The ideal $0 \oplus k \subseteq \Gamma(V, \mathcal{O}_{\text{surj}})$ defines an ideal sheaf in the radical topology, but it is not the sheafification of an ideal in V .

Example 3.5.3. Let V denote a discrete valuation domain with maximal ideal $\mathfrak{m} = (\pi)$, consider $I = V\pi^2 \subset J = V\pi \subset V$. Then the radical of $V\pi^2$ inside V is $V\pi$. The inclusion $V\pi^2 \subset V\pi$ is as V -modules isomorphic to $V\pi \subset V$. Hence the radical of $V\pi^2$ inside $V\pi$ is just $V\pi^2$. So the radical of a submodule depends on the surrounding module.

This also shows that ideals $I \subseteq J$ with the same radical do not induce isomorphisms $I_{\text{surj}} \rightarrow J_{\text{surj}}$. This mapping is in our example just $\mathcal{O}_{\text{surj}} \rightarrow \mathcal{O}_{\text{surj}}$, $1 \mapsto \pi$. However the image ideal sheaves of I_{surj} and J_{surj} in $\mathcal{O}_{\text{surj}}$, that is, I^{surj} and J^{surj} , are the same. Also, the surjective homomorphism $R/I \rightarrow R/J$ induces an isomorphism $(\mathcal{O}/I)_{\text{surj}} \rightarrow (\mathcal{O}/J)_{\text{surj}}$.

3.6. Constructible partitions.

For a discrete valuation domain V with quotient field K and residue class field k the ring homomorphism $V \rightarrow K \oplus k$ defines a surjection

$$\text{Spec } K \uplus \text{Spec } k \rightarrow \text{Spec } V,$$

which is neither flat nor submersive. Since $(K \oplus k) \otimes_V (K \oplus k) = K \oplus k$, every element in $K \oplus k$ is compatible and defines a global section on $\text{Spec } V$ in the surjective topology. We treat here similar constructible partitions of a scheme and the global sections in the surjective topology which arise from them.

Recall that a subset in a scheme X is called *constructible* if it is the finite union of intersections of a closed and of an open subset (see [19, §0.9], [27, Exercise II. 3.18] or [12, Section 14.3]).

Definition 3.6.1. We call a finite disjoint union $X = \uplus_{i \in I} X_i$, where each X_i factors as $X_i \rightarrow W_i \rightarrow X$ with an open embedding $X_i \rightarrow W_i$ and a closed embedding $W_i \rightarrow X$ a *constructible partition* of X . If all X_i are affine, then we call this an *affine constructible partition* of X .

Remark 3.6.2. We can refine every constructible partition to an affine constructible partition. We can further refine so that each patch X_i is integral, i.e. irreducible and reduced. If X is a scheme of finite type over a field, then we can get also a regular constructible partition [20, Corollaire 6.12.5]. Note that every constructible partition $\uplus_{i \in I} X_i \rightarrow X$ is a bijection, and that for every $x \in X$ the induced homomorphism of residue class fields is an isomorphism.

We introduce the constructible topology, where essentially the constructible partitions are the coverings.

Definition 3.6.3. The *constructible topology* on a scheme X is given by the full subcategory $\mathcal{T}_X$ of schemes of finite type which are affine over X . A morphism $U \rightarrow V$ in $\mathcal{T}_X$ is a cover if there exists a constructible partition $\uplus_{i \in I} V_i \rightarrow V$ and a factorization $\uplus_{i \in I} V_i \rightarrow U \rightarrow V$. We denote a scheme X endowed with the constructible topology by X_{cons} .

Remark 3.6.4. The constructible topology is an affine single-handed Grothendieck topology. It is a refinement of the affine single-handed Zariski topology. Every scheme morphism $Y \rightarrow X$ induces a site-morphism $Y_{\text{cons}} \rightarrow X_{\text{cons}}$. The surjective topology is a refinement of the constructible topology, so there exists a site morphism $X_{\text{surj}} \rightarrow X_{\text{cons}}$. We will see below (Theorem 3.8.2, Corollary 3.8.5) that in characteristic 0 these two topologies yield the same modules of global sections.

Lemma 3.6.5. *Let X denote a scheme and let $X_i = \text{Spec } R_i$, $i \in I$, denote an affine constructible partition of X . Then $(\bigsqcup_{i \in I} X_i) \times_X (\bigsqcup_{i \in I} X_i) \cong (\bigsqcup_{i \in I} X_i)$, and the projections are isomorphisms.*

Proof. Note that $Z \times_X Z \cong Z$ for a closed immersion $Z \rightarrow X$ and $U \times_X U \cong U$ for an open subscheme $U \subseteq X$. Hence this is also true for a composition $U \rightarrow Z \rightarrow X$. Moreover, for disjoint immersions $X_i, X_j \rightarrow X$ we have $X_i \times_X X_j = \emptyset$. So

$$\left(\bigsqcup_{i \in I} X_i\right) \times_X \left(\bigsqcup_{i \in I} X_i\right) \cong \bigsqcup_{(i,j) \in I \times I} X_i \times_X X_j \cong \bigsqcup_{i \in I} X_i \times_X X_i \cong \bigsqcup_{i \in I} X_i.$$

□

Corollary 3.6.6. *Let X denote a scheme endowed with the constructible or the surjective topology. Let X_i , $i \in I$, denote a constructible partition of X , and let $\mathcal{F}$ denote a sheaf of abelian groups on X_{top} , where $\text{top} = \text{surj}$ or cons . Then the restriction homomorphism*

$$\Gamma(X_{\text{top}}, \mathcal{F}) \rightarrow \Gamma\left(\bigsqcup_{i \in I} X_{i \text{top}}, \mathcal{F}\right)$$

is an isomorphism.

Proof. This isomorphism follows from Lemma 3.6.5 and Lemma 1.6.7. □

Corollary 3.6.7. *Let X denote a scheme and let M be an $\mathcal{O}_X$ -module on X . Let $j_i : X_i \rightarrow X$, $i \in I$, denote a constructible partition of X . Then in the constructible or in the surjective topology we have an isomorphism*

$$\Gamma(X_{\text{top}}, M_{\text{top}}) \rightarrow \bigoplus_{i \in I} \Gamma(X_{i \text{top}}, (j_i^*(M))_{\text{top}}).$$

In particular, any $(s_i)_{i \in I} \in \bigoplus_{i \in I} \Gamma(X_{i \text{top}}, j_i^(M))$ yields a global section in $\Gamma(X_{\text{top}}, M_{\text{top}})$.*

Proof. Due to Corollary 3.6.6 we may assume that $X = \bigsqcup_{i \in I} X_i$ is a disjoint union of open subschemes. Then this follows from Lemma 1.5.10 □

Proposition 3.6.8. *Let $X = \text{Spec } R$ be noetherian of finite dimension. Then the closure operation for finitely generated modules induced by the constructible topology is the radical.*

Proof. Since the surjective topology is a refinement of the constructible topology and because of Proposition 3.2.3 we only have to show for $s \in \text{rad}(N)$, $N \subseteq M$ finitely generated modules, that s belongs locally to N in a constructible partition. We may assume that $N = 0$ and we may pass to a constructible partition where the patches are affine and integral. So we may assume that R is a noetherian domain of finite dimension. Let $s \in \text{rad}(0)$, so that $s \otimes 1 = 0$ in $M \otimes_R K$ for every field $R \rightarrow K$. Then in particular $s \otimes 1 = 0$ in $M \otimes_R Q(R)$ and so $s \otimes 1 = 0$ in $M \otimes_R R_g$ for some $0 \neq g \in R$. Considering the constructible partition $D(g) \sqcup V(g)$ we can go on inductively. □

Example 3.6.9. In the case of an ideal we can give a constructible partition as predicted by Proposition 3.6.8 immediately. Let $f \in \text{rad}(I)$, $I = (f_1, \dots, f_n)$ an ideal in a commutative ring R . An affine constructible partition is given by the reductions of

$$D(f_1) \uplus (D(f_2) \cap V(f_1)) \uplus \dots \uplus (D(f_n) \cap V(f_1, \dots, f_{n-1})) \uplus V(f_1, \dots, f_n).$$

The corresponding rings are $R_k = (R/(f_1, \dots, f_k)_{f_{k+1}})_{\text{red}}$, $k = 0, \dots, n$ (setting $f_{n+1} = 1$). In each of the R_k we clearly have $f \in IR_k$, since this extended ideal is the unit ideal for $k \leq n-1$ and for $k = n$ it is the zero-ideal, but also f becomes 0 in $R_n = (R/(f_1, \dots, f_n))_{\text{red}}$.

This constructible partition is perhaps even more canonical than the forcing algebra B of these data. By the universal property of the forcing algebra (Proposition 2.1.4) there exists a ring homomorphism $B \rightarrow \bigoplus_{k=0}^n R_k$. Since the global sections of this constructible partition are global sections in the surjective structure sheaf by Corollary 3.6.7, this also shows that $f \in \Gamma(R, \mathcal{O}_{\text{surj}})$, and not only that f belongs locally to this ideal sheaf.

3.7. Surjections and constructible partitions.

In this section we gather together several statements which relate surjections to constructible partitions and their finite extensions which will later on help us to compute modules of global sections and cohomology in the surjective topology.

Lemma 3.7.1. *Let X denote a noetherian scheme of finite dimension and let $\varphi : Y \rightarrow X$ be a spec-surjective morphism of finite type. Then there exists an affine constructible partition $\biguplus_{i \in I} X_i \rightarrow X$ such that there is an (X -covering) refinement $\biguplus_{i \in I} V_i \rightarrow \biguplus_{i \in I} X_i' \rightarrow Y$, where $X_i' = X_i \times_X Y$, $X_i = \text{Spec } R_i$ and $V_i = \text{Spec } S_i$, such that $R_i \rightarrow S_i$ is a finitely generated faithfully flat extension of domains. If X is a scheme over a field of characteristic 0, then one can also achieve that $S_i \otimes_{R_i} S_i$ is reduced.*

Proof. We do induction on the dimension of X and we construct step by step an affine integral constructible partition of X with the required properties. We pass first to an affine constructible partition of X such that the patches are affine and integral. We may also assume that Y is affine, by replacing Y by a disjoint union of an open affine cover. So we may assume that R is a domain, and that $\varphi : Y = \text{Spec } S \rightarrow X = \text{Spec } R$ is surjective. By the Theorem of Chevalley [12, Corollary 14.7] the images of the irreducible components of $\text{Spec } S$ are constructible. In particular there exists at least one component such that its image contains an open affine subset U of $\text{Spec } R$. We pass to the constructible affine partition $U \uplus Z$ (Z the closed component). Then, on the one hand, the morphism $\varphi^{-1}(U) \rightarrow U$ has a refinement by an affine integral scheme, and, on the other hand, the image of the surjection $\varphi^{-1}(Z) \rightarrow Z$ has lower dimension and can be treated as before.

So we may assume that $R \rightarrow S$ are both domains. By generic flatness ([20, Théorème 6.9.1] or [12, Theorem 14.4]) there exists $0 \neq g \in R$ such that $R_g \rightarrow S_g$ is faithfully flat. We pass to the corresponding constructible partition, the complement being again of lower dimension. So we get a partition and a refinement such that the mappings $U_i \rightarrow V_i$ are faithfully flat morphisms between domains.

Suppose now that R is a domain which contains a field of characteristic zero and let $R \subseteq S$ be a faithfully flat extension of finite type, S a domain. We consider the commutative

diagram

$$\begin{array}{ccccc}
 R & \longrightarrow & S & \xrightarrow{i_1} & S \otimes_R S \\
 \downarrow & & \downarrow & & \downarrow \\
 Q(R) & \longrightarrow & S_{R^*} & \xrightarrow{i_1} & (S \otimes_R S)_{R^*} = S_{R^*} \otimes_{Q(R)} S_{R^*}.
 \end{array}$$

Since the characteristic is zero, $Q(R)$ is a perfect field and so $S_{R^*} \otimes_{Q(R)} S_{R^*}$ is reduced by [20, Proposition 4.6.1]. If $\mathfrak{a} \subseteq S \otimes_R S$ is the nilradical, then it maps to 0 in $(S \otimes_R S)_{R^*}$, and so there exists $g \in R^*$ with $g\mathfrak{a} = 0$. Then $(S \otimes_R S)_g = S_g \otimes_{R_g} S_g$ is reduced and we look at the partition $X = D(g) \uplus V(g)$ and use the induction hypothesis on $V(g)$. $\square$

Lemma 3.7.2. *Let $Y \rightarrow X$ be a surjection of finite type between noetherian schemes of finite dimension. Then there exists an affine integral constructible partition $X_i \rightarrow X$ and finite surjective morphisms $X'_i \rightarrow X_i$, where X'_i are also affine and integral, and sections $X'_i \rightarrow Y$.*

Proof. The statement is stable under refining by constructible partitions, so we may assume that $X = \text{Spec } R$, where R is a noetherian domain with quotient field $Q(R)$. For a closed point $x \in Y_{Q(R)}$ the extension $Q(R) \subseteq k(x)$ is of finite type and hence finite. Let $R \subseteq R' \subset k(x)$ be finite such that $Q(R') = k(x)$. Then there exists an open subset $D(h) \subseteq \text{Spec } R'$ such that $i : x = \text{Spec}(k(x)) \rightarrow Y$ extends to $D(h) \rightarrow Y$. For this reduce to $Y = \text{Spec } A$ affine and find a common denominator h for the images $i^*(T_j) = q_j \in k(x)$, where the T_j are R' -algebra generators of A . There exists also $\emptyset \neq D(g) \subseteq X$ such that the preimage of $D(g)$ lies inside $D(h) \subseteq \text{Spec } R'$. Therefore for $D(g) \subseteq X$ we have a finite extension $X' = \text{Spec } R'|_{D(g)}$ and a section $X' \rightarrow Y$. We go on on $V(g) \subset X$ inductively. $\square$

Lemma 3.7.3. *Let $\varphi : Y \rightarrow X$ be a finite surjective morphism between noetherian schemes, where X is integral of finite dimension. Then there exists an affine integral constructible partition $X_i \rightarrow X$ such that $Y_i = X_i \times_X Y$ is a disjoint union of irreducible schemes.*

Proof. Let $Z_j, j \in J$, be the irreducible components of Y . The union of the intersections $Z_j \cap Z_{j'}$ ($j \neq j'$) has smaller dimension than $\dim(X)$, so its image in X is a proper closed subset $V \subset X$. For an open non-empty affine subset $U \subseteq X - V$ in its complement the components $U_j = \varphi^{-1}(U) \cap Z_j$ are disjoint. So we get inductively a constructible partition of X with the required properties. $\square$

3.8. Global sections in the surjective topology.

We characterize now the ring of global sections in the constructible topology and in the surjective topology. We first describe $\Gamma(X_{\text{cons}}, \mathcal{O}_{\text{cons}})$ as a subring of the radicalization of R .

Proposition 3.8.1. *Let R denote a noetherian ring, and let $X = \text{Spec } R$ be endowed with the constructive topology. Then*

$$\begin{aligned}
 \Gamma(X_{\text{cons}}, \mathcal{O}_{\text{cons}}) = \{ & f \in \prod_{x \in X} \overline{\kappa(x)} : \text{there exists an affine constructible partition} \\
 & R \rightarrow \bigoplus_{i \in I} R_i \text{ and } f_i \in R_i \text{ such that } f_i(x) = f(x) \text{ in } \overline{\kappa(x)} \}.
 \end{aligned}$$

Proof. We have natural homomorphisms

$$\Gamma(X_{\text{cons}}, \mathcal{O}_{\text{cons}}) \longrightarrow \Gamma(X_{\text{surj}}, \mathcal{O}_{\text{surj}}) \longrightarrow \prod_{x \in X} \overline{\kappa(x)}$$

by Proposition 3.5.1. Since the radicalization is the absolute stalk, the second homomorphism is injective by Lemma 1.9.5. The first homomorphism is induced by the site morphism $X_{\text{surj}} \rightarrow X_{\text{cons}}$. This homomorphism is also injective: if $f \in \Gamma(X_{\text{cons}}, \mathcal{O}_{\text{cons}})$ is represented by $f_i \in \Gamma(X_i, \mathcal{O}_i)$, $X = \bigsqcup_{i \in I} X_i$ an affine integral constructible partition and maps to 0 in $\Gamma(X_{\text{surj}}, \mathcal{O}_{\text{surj}})$, then there exists a surjection $Y \rightarrow X$ such that f becomes 0. But then also the pull-backs of f_i to $X_i \times_X Y$ are 0 for all $i \in I$. This implies, since the X_i are integral, that $f_i = 0$ and so $f = 0$ in $\Gamma(X_{\text{cons}}, \mathcal{O}_{\text{cons}})$.

The image of $\Gamma(X_{\text{cons}}, \mathcal{O}_{\text{cons}})$ inside the absolute stalk is the described subring: an element $f \in \Gamma(X_{\text{cons}}, \mathcal{O}_{\text{cons}})$ is represented by $f_i \in R_i$, where $X_i = \text{Spec } R_i$ is an affine constructible partition of X , and every such tuple yields a global element. $\square$

Theorem 3.8.2. *Let R denote a noetherian ring of finite dimension endowed with the surjective topology, $X = \text{Spec } R$. Suppose that R contains a field of characteristic zero. Then*

$$\Gamma(X_{\text{surj}}, \mathcal{O}_{\text{surj}}) = \Gamma(X_{\text{cons}}, \mathcal{O}_{\text{cons}}).$$

Proof. We have proven already in Proposition 3.8.1 that the natural homomorphism $\Gamma(X_{\text{cons}}, \mathcal{O}_{\text{cons}}) \rightarrow \Gamma(X_{\text{surj}}, \mathcal{O}_{\text{surj}})$ is injective. For the surjectivity suppose that $f \in \Gamma(X, \mathcal{O}_{\text{surj}})$ is represented by a compatible element $f \in S$, $R \rightarrow S$ being spec-surjective. We have to construct a constructible partition of X such that f comes from there. By Lemma 3.7.1 there exists an affine constructible partition $X = \bigsqcup_{i \in I} X_i$ and affine schemes $V_i \rightarrow X_i \times_X Y$ such that $\bigsqcup V_i \rightarrow Y \rightarrow X$ is a surjection, that $V_i \rightarrow X_i$ are faithfully flat morphisms of finite type between integral affine schemes and that $V_i \times_{X_i} V_i$ is reduced. We can pass to the refinement $\bigsqcup V_i$ and may assume that f is given as $f_i \in S_i$, $V_i = \text{Spec } S_i$, which are compatible over X_i . We can treat the f_i separately.

The compatibility of $f_i \in S_i$ means that $f_i \otimes 1 - 1 \otimes f_i$ is nilpotent by Corollary 3.2.5. Hence this difference is 0, since $S_i \otimes_{R_i} S_i$ is reduced. But this means that f_i is also compatible in the faithfully flat topology; hence $f_i \in R_i$ [49, Proposition I.2.18], where $X_i = \text{Spec } R_i$. $\square$

Remark 3.8.3. Theorem 3.8.2 and Proposition 3.8.1 show that every element $f \in \Gamma(X_{\text{surj}}, \mathcal{O}_{\text{surj}})$ has at every point $x \in X$ a well-defined value $f(x) \in \kappa(x)$, not only in $\overline{\kappa(x)}$. This is also clear considering $\text{Spec } \kappa(x) \rightarrow X$. This scheme morphism defines a site morphism $(\text{Spec } \kappa(x))_{\text{surj}} \rightarrow X_{\text{surj}}$ and therefore a homomorphism of the rings of global sections, $\Gamma(X, \mathcal{O}_{\text{surj}}) \rightarrow \Gamma(\text{Spec } \kappa(x), \mathcal{O}_{\text{surj}}) = \kappa(x)$ (in characteristic zero). This reasoning shows also that in positive characteristic every global function over X_{surj} has for every point x a well-defined value in $\kappa(x)^\infty$.

We extend some of the results in this section to coherent modules.

Lemma 3.8.4. *Let X be a noetherian scheme of finite dimension and let M be a coherent sheaf on X . Then there exists an affine constructible partition $\bigsqcup_{i \in I} X_i \rightarrow X$ such that $j_i^*(M)$ is free, where $j_i : X_i \rightarrow X$ is the natural embedding.*

Proof. We may assume that $X = \text{Spec } R$ is affine and integral. Then we have an isomorphism $M \otimes_R Q(R) \cong Q(R)^n$. Therefore $M_g \cong R_g^n$ for some $0 \neq g \in R$ and the statement is true on $U = D(g)$. Then we may go on on $V(g)$ inductively. $\square$

Corollary 3.8.5. *Let X be a noetherian scheme of finite dimension over a field of characteristic 0 and let M be a coherent sheaf on X . Then the global sections in $\Gamma(X_{\text{surj}}, M_{\text{surj}})$ are constructible sections, i.e. they are of the form $s = (s_i)_{i \in I}$, $s_i \in \Gamma(X_i, j_i^*(M))$, where $X = \bigsqcup_{i \in I} X_i$ is an affine constructible partition.*

Proof. We may assume by Lemma 3.8.4 that $X = \bigsqcup X_i$ is such that all $j_i^*(M)$ are free. Then $\Gamma(X_{\text{surj}}, M_{\text{surj}}) = \bigoplus_{i \in I} \Gamma(X_{i \text{ surj}}, M_{i \text{ surj}}) = \bigoplus_{i \in I} \Gamma(X_{i \text{ surj}}, \mathcal{O}_{\text{surj}}^{n_i})$ (by Corollary 3.6.7) and the result follows from Theorem 3.8.2. $\square$

3.9. Exactness in the surjective topology.

We characterize when a complex of modules is exact in the surjective topology. First examples were already given in Proposition 2.7.6.

Theorem 3.9.1. *Let R denote a commutative ring and let $N \xrightarrow{\alpha} M \xrightarrow{\beta} L$ denote a complex of finitely generated R -modules. Then the following are equivalent (suppose in the last statement that R is noetherian of finite dimension and contains a field of characteristic 0).*

- (i) *The complex of sheaves $N_{\text{surj}} \xrightarrow{\alpha} M_{\text{surj}} \xrightarrow{\beta} L_{\text{surj}}$ on $(\text{Spec } R)_{\text{surj}}$ is exact (where T_R contains all R -algebras or all R -algebras of finite type).*
- (ii) *For every ring homomorphism $R \rightarrow R'$ (of finite type), the complex $N \otimes_R R' \xrightarrow{\alpha} M \otimes_R R' \xrightarrow{\beta} L \otimes_R R'$ is such that the kernel lies inside the radical of the image.*
- (iii) *The complex $N \otimes_R \text{rad}(R) \xrightarrow{\alpha} M \otimes_R \text{rad}(R) \xrightarrow{\beta} L \otimes_R \text{rad}(R)$ of $\text{rad}(R)$ -modules is exact.*
- (iv) *For every homomorphism $R \rightarrow K$ to a field the complex $N \otimes_R K \xrightarrow{\alpha} M \otimes_R K \xrightarrow{\beta} L \otimes_R K$ of K -vector spaces is exact.*
- (v) *The complex $\Gamma(R', N_{\text{top}}) \xrightarrow{\alpha} \Gamma(R', M_{\text{top}}) \xrightarrow{\beta} \Gamma(R', L_{\text{top}})$ of global sections is exact for all ring homomorphism $R \rightarrow R'$ of finite type.*

Proof. The conditions (i), (ii) and (iv) are equivalent by Corollary 2.7.3. (i) $\Rightarrow$ (iii) follows from Corollary 1.8.5, Lemma 1.10.3 and Proposition 3.5.1. (iii) $\Rightarrow$ (iv). A homomorphism $R \rightarrow K$ factors as $R \rightarrow \kappa(x) \rightarrow K$, where $x \in \text{Spec } R$. By restricting the exactness of (iii) to the component given by x , it follows that $N \otimes_R \overline{\kappa(x)} \xrightarrow{\alpha} M \otimes_R \overline{\kappa(x)} \xrightarrow{\beta} L \otimes_R \overline{\kappa(x)}$ is exact. This is then also true if we replace $\overline{\kappa(x)}$ by $\kappa(x)$ and then by K .

(iv) $\Rightarrow$ (v). The condition in (iv) holds also for R' , and $(M \otimes_R R')_{\text{top}} = (M_{\text{top}})|_{X'}$ on X'_{surj} ($X' = \text{Spec } R'$) by Corollary 1.5.6, hence we may assume that $R = R'$. Let $s \in \Gamma(X_{\text{surj}}, M_{\text{top}})$ mapping to 0 in $\Gamma(X_{\text{surj}}, L_{\text{top}})$.

By Corollary 3.8.5 we may assume that s is represented by $s_i \in M \otimes_R R_i$, where $j_i : X_i = \text{Spec } R_i \rightarrow X$ is an affine integral constructible partition. We have to find a global section $t \in \Gamma(X_{\text{surj}}, N_{\text{surj}})$ which maps to s . Since $\Gamma(X_{\text{surj}}, N_{\text{surj}}) = \bigoplus_{i \in I} \Gamma(X_{i \text{ surj}}, j_i^*(N)_{\text{surj}})$ (Corollary 3.6.7) we may assume that $X = \bigsqcup_{i \in I} X_i$ is the disjoint union of open subschemes and we can treat the components separately; hence we skip the index i . Since s maps to 0, there exist spec-surjective homomorphisms $R \rightarrow S$ such that $\beta(s) \otimes 1 = 0$ in $L \otimes_R S$. Then by Proposition 3.1.1 for every field $R \rightarrow K$ we have $\beta(s) \otimes 1 = 0$ in $L \otimes_R K$, so in particular for the quotient field $Q(R)$.

By condition (iv) there exists an element $t \in N \otimes_R Q(R)$ mapping to s , and this is then also true for $t \in N \otimes_R R_g$, $0 \neq g \in R$. Therefore s comes from the left on the open set $D(g) \subseteq X$ and we pass to the constructible partition $D(g) \uplus V(g)$. The $R/(g)$ -complex $N \otimes_R (R/(g)) \xrightarrow{\alpha} M \otimes_R (R/(g)) \xrightarrow{\beta} L \otimes_R (R/(g))$ fulfills also property (iv), so we can construct inductively a constructible partition $X_j \rightarrow X$ such that s comes from an element $t_j \in N \otimes_R R_j$. Such a tuple is compatible and defines a global section in $\Gamma(X_{\text{surj}}, N_{\text{surj}})$ by Corollary 3.6.7.

(v) $\Rightarrow$ (i) is trivial, since a global solution is also a local solution. $\square$

Corollary 3.9.2. *Let R denote a commutative ring of finite dimension containing a field of characteristic 0, $X = \text{Spec } R$. Let $M \rightarrow L \rightarrow 0$ be an exact sequence of finitely generated R -modules. Then also $\Gamma(X_{\text{surj}}, M_{\text{surj}}) \rightarrow \Gamma(X_{\text{surj}}, L_{\text{surj}})$ is surjective.*

Proof. The complex of sheaves $M_{\text{surj}} \rightarrow L_{\text{surj}} \rightarrow 0$ is exact, hence by Theorem 3.9.1(v) also $\Gamma(X_{\text{surj}}, M_{\text{surj}}) \rightarrow \Gamma(X_{\text{surj}}, L_{\text{surj}}) \rightarrow 0$ is exact. $\square$

Example 3.9.3. Let R be a commutative ring with reduction R_{red} . Then the complex of R -modules $0 \rightarrow R \rightarrow R_{\text{red}}$ is exact only when R is reduced. But tensoring with $\kappa(\mathfrak{p})$ (and so with every field) yields always an exact complex, since $R_{\text{red}} \otimes_R R/\mathfrak{p} = R/\mathfrak{n}_R \otimes_R R/\mathfrak{p} = R/\mathfrak{q}$, where $\mathfrak{n}_R$ is the nilradical. So this complex is exact in the surjective topology.

Remark 3.9.4. Using the characterization in Theorem 3.9.1(ii) one can show that for a ring of finite type over a field K the exactness in the surjective topology is equivalent to the exactness after every ring homomorphism $R \rightarrow \overline{K}$.

Proposition 3.9.5. *Let R be a noetherian ring of finite dimension over a field of characteristic zero and let $N \subseteq M$ be finitely generated R -modules. Let $N_{\text{surj}} \rightarrow N^{\text{surj}} \subseteq M_{\text{surj}}$ be the surjection of sheaves in the surjective topology on $X = \text{Spec } R$. Then also $\Gamma(X_{\text{surj}}, N_{\text{surj}}) \rightarrow \Gamma(X_{\text{surj}}, N^{\text{surj}})$ is surjective.*

Proof. Let $s \in \Gamma(X_{\text{surj}}, N^{\text{surj}}) \subseteq \Gamma(X_{\text{surj}}, M_{\text{surj}})$. Recall that N^{surj} is the image sheaf of $N_{\text{surj}} \rightarrow M_{\text{surj}}$, so s comes locally from N_{surj} . Due to Corollary 3.8.5 we can represent s by $s_i \in \Gamma(X_i, j_i^*(M)) = M \otimes_R R_i$, (j_i^* is the Zariski pull-back), where $X_i \rightarrow X$ is an affine partition of X . As the global sections of N_{surj} also respect this partition we may assume that $s \in M$. Since $s \in \Gamma(X_{\text{surj}}, N^{\text{surj}})$, we know by Proposition 3.6.8 that there exists a constructible partition $X_i \rightarrow X$ and sections $t_i \in \Gamma(X_i, N_{\text{surj}})$ mapping to the restrictions of s . Again by Corollary 3.8.5 this represents a global element $t \in \Gamma(X, N_{\text{surj}})$ mapping to s . $\square$

For free modules the exactness in the surjective topology is quite a strong condition.

Lemma 3.9.6. *Let $R^a \xrightarrow{\alpha} R^b \xrightarrow{\beta} R^c$ be complex of free R -modules, so that α and β are given by matrices. Let $J_i(\alpha)$ denote the i th minor ideal, i.e. the ideal generated by all determinants of the $i \times i$ -submatrices of α . Then the following are equivalent.*

- (i) *We have $\sum_{i+j=b} J_i(\alpha) J_j(\beta) = (1)$.*
- (ii) *For every ring homomorphism $R \rightarrow R'$ the tensored complex is exact.*
- (iii) *The complex is exact in the surjective topology.*

Proof. (i) $\Rightarrow$ (ii). Since the minor condition is stable under arbitrary base change, we may assume that $R' = R$ and that R is local. For a local ring the condition means that there

exists an invertible submatrix of α of size i and an invertible submatrix of β of size j with $i + j = b$. After a base change in the first two modules we may assume that the first j columns in the matrix α are the first j unit vectors. It follows that the first j columns of β must be zero. Therefore the remaining $b - j = i$ columns of β contain an invertible $i \times i$ -submatrix. It follows that the first j unit vectors generate exactly the kernel of β , and this is also the image of α .

The implication (ii) $\Rightarrow$ (iii) is trivial. (iii) $\Rightarrow$ (i). If the minor condition is not fulfilled, then $\sum_{i+j=b} J_i(\alpha)J_j(\beta)$ is contained in a maximal ideal and so there exists a field $R \rightarrow K$ such that $\sum_{i+j=b} J_i(\alpha)J_j(\beta) = 0$. Let $i = \text{rk}(\alpha)$ over K . Then $\text{rk}(\beta) < b - i$ and $\dim(\text{kern } \beta) = b - \dim(\text{im } \beta) > b - (b - i) = i$, so the tensorized complex is not exact. $\square$

Example 3.9.7. Let $f \in R$ be a non-zero divisor and a non unit. Then the sequence $R \xrightarrow{0} R \xrightarrow{f} R$ is exact, but not exact in the surjective topology. For a maximal ideal $\mathfrak{m}$ containing f the tensoration with $K = R/\mathfrak{m}$ yields $0 \rightarrow K \xrightarrow{0} K$.

Remark 3.9.8. Lemma 3.9.6 says in particular that for given $\alpha : R^a \rightarrow R^b$ it can be checked in the surjective topology (i.e. after tensoration with fields alone) whether it is an isomorphism or not. This statement can be found for a homomorphism between flat modules over a noetherian ring in [28].

For a homomorphism α between free modules (or for a matrix) we call the maximum i such that $J_i(\alpha)$ contains a non zero divisor its rank (cf. [10, Proposition 1.4.11]).

Corollary 3.9.9. Let $G_\bullet = G_n \xrightarrow{\alpha_{n-1}} G_{n-1} \rightarrow \dots \rightarrow G_2 \xrightarrow{\alpha_1} G_1 \xrightarrow{\alpha_0} G_0 \rightarrow 0$ be a complex of free R -modules G_k of finite ranks b_k . Suppose that $r_k = \text{rk}(\alpha_k) = b_k - b_{k-1}$ (this is the standard condition on ranks). Then the following are equivalent.

- (i) The minors are $J_{r_k}(\alpha_k) = (1)$ for all $k = 0, \dots, n$.
- (ii) The complex $G_\bullet \otimes_R R'$ is exact for every $R \rightarrow R'$.
- (iii) The complex is exact in the surjective topology.

Proof. (i) $\Leftrightarrow$ (iii). Note that $r_k + r_{k-1} = b_k$ and that $J_i(\alpha_k) \neq (1)$ for $i > r_k$. Then the minor condition of Lemma 3.9.6 for the k th place reduces to $J_{r_k}(\alpha_k)J_{r_{k-1}}(\alpha_{k-1}) = (1)$. Hence this equivalence follows by induction on k .

(ii) $\Rightarrow$ (iii) is contained in Theorem 3.9.1, so assume that (i) holds. Since the minor condition respects base changes, we may assume that $R = R'$ and that R is local. Since $J_{r_0}(\alpha_0) = (1)$ and $r_0 = b_0$ the homomorphism α_1 is surjective. Then the kernel $\text{kern}(\alpha_0)$ is free of rank $b_1 - b_0$ and the homomorphism $R^{b_2} \rightarrow R^{b_1 - b_0} \cong \text{kern}(\alpha_0) \subseteq R^{b_1}$ fulfills also the minor condition. So we can go on inductively. $\square$

Remark 3.9.10. The Buchsbaum-Eisenbud criterion asserts that if in the situation of Corollary 3.9.9 (with also a 0 on the left) the so-called standard condition on depth is fulfilled, namely that $\text{depth}(J_{r_k}(\alpha_k)) \geq k + 1$, then the complex itself is exact. A variant of Hochster and Huneke [38, Theorem 10.3] says that if the height condition $\text{ht}(J_{r_k}(\alpha_k)) \geq k + 1$ is true, then the complex has phantom homology.

3.10. Cohomology in the surjective topology.

In this section we show that the first cohomology of coherent modules in the constructible and in the surjective topology is zero.

Proposition 3.10.1. *Let X denote a scheme endowed with the constructible or the surjective topology. Let $X' = \bigsqcup_{i \in I} X_i \rightarrow X$ denote a constructible partition. Let $\mathcal{P}$ denote a presheaf of abelian groups. Then the Čech cohomology groups for this covering are $\check{H}^0(X' \rightarrow X, \mathcal{P}) = \Gamma(X', \mathcal{P})$ and $\check{H}^k(X' \rightarrow X, \mathcal{P}) = 0$ for $k \geq 1$.*

Proof. This follows from Lemma 3.6.5 and Lemma 1.11.1. $\square$

The first cohomology for coherent sheaves vanishes also in the surjective topology, but the proof is more involved. We restrict to characteristic zero.

Theorem 3.10.2. *Let X be a noetherian scheme of finite dimension over a field of characteristic 0. Then $\check{H}^1(X_{\text{surj}}, \mathcal{O}_{\text{surj}}) = 0$.*

Proof. Let $Y \rightarrow X$ be a surjection of finite type. We have to show that a Čech cocycle represented by this covering is 0. To show this we may use refinements $Z \rightarrow Y$, where $Z \rightarrow X$ is also a covering. Let $X_i \rightarrow X$ be a constructible partition and set $Y_i = X_i \times_X Y \rightarrow X_i$. The first Čech cohomology (with respect to the given cover) is given as the homology of the first row in the following commutative diagram

$$\begin{array}{ccccc}
 \Gamma(Y, \mathcal{O}_{\text{surj}}) & \longrightarrow & \Gamma(Y \times_X Y, \mathcal{O}_{\text{surj}}) & \longrightarrow & \Gamma(Y \times_X Y \times_X Y, \mathcal{O}_{\text{surj}}) \\
 \downarrow & & \downarrow & & \downarrow \\
 \Gamma(\bigsqcup Y_i, \mathcal{O}_{\text{surj}}) & \longrightarrow & \Gamma(\bigsqcup (Y_i \times_{X_i} Y_i), \mathcal{O}_{\text{surj}}) & \longrightarrow & \Gamma(\bigsqcup (Y_i \times_{X_i} Y_i \times_{X_i} Y_i), \mathcal{O}_{\text{surj}}) \\
 \downarrow & & \downarrow & & \downarrow \\
 \bigoplus_i \Gamma(Y_i, \mathcal{O}_{\text{surj}}) & \longrightarrow & \bigoplus_i \Gamma(Y_i \times_{X_i} Y_i, \mathcal{O}_{\text{surj}}) & \longrightarrow & \bigoplus_i \Gamma(Y_i \times_{X_i} Y_i \times_{X_i} Y_i, \mathcal{O}_{\text{surj}}),
 \end{array}$$

where the down-arrows are isomorphisms by Corollaries 3.6.6 and 3.6.7. An element $s \in \Gamma((Y \times_X Y)_{\text{surj}}, \mathcal{O}_{\text{surj}})$ is a cocycle if and only if for every i its component $s_i \in \Gamma((Y_i \times_{X_i} Y_i)_{\text{surj}}, \mathcal{O}_{\text{surj}})$ is a cocycle. We have to show that s comes after a refinement $Z \rightarrow Y$ from the left, and for this we can look at the components separately. Therefore we will repeatedly simplify the situation by using a constructible partition of X .

By Lemma 3.7.2 and by the preceding consideration we may assume that $Y \rightarrow X$ is a finite surjective morphism between noetherian affine integral schemes. We may also assume by Lemma 3.7.1 that this morphism is flat. Suppose that $s \in \Gamma(Y \times_X Y, \mathcal{O}_{\text{surj}})$ maps to 0 on the right. This s is by Theorem 3.8.2 represented by $s_j \in \Gamma(Z_j, \mathcal{O})$, $j \in J$, where $\bigsqcup_{j \in J} Z_j \rightarrow Y \times_X Y$ is a constructible partition into affine integral schemes.

By Lemma 3.7.3 we find a constructible partition of X such that we may assume that all irreducible components V_i , $i \in I$, of $Y \times_X Y$ are disjoint. We may furthermore assume by refining that the partition Z_j of $Y \times_X Y$ (where the cocycle $s = s_j$ lives) is a subpartition of the disjoint irreducible components. So we write the cocycle as $s_{ij} \in \Gamma(Z_{ij}, \mathcal{O})$, where $Z_{ij} \rightarrow V_i$, $j \in J_i$, are constructible partitions. Let $W_i = Z_{ij_i}$ be the open subsets of the components V_i where the generic elements $s_{ij_i} \in \Gamma(W_i, \mathcal{O})$ live. Then the images of $V_i - W_i$ are closed and their union over $i \in I$ is a closed subset A of X of smaller dimension. We consider the constructible partition $X = U \sqcup A$. The generic tuple s_{ij_i} , $i \in I$, is represented by the element

$$(s_{ij_i})_{i \in I} \in \Gamma(\bigsqcup_{i \in I} V_i \cap (p^{-1}(U) \times_U p^{-1}(U)), \mathcal{O}) = \Gamma(p^{-1}(U) \times_U p^{-1}(U), \mathcal{O}).$$

Note that $V_i \cap (p^{-1}(U) \times_U p^{-1}(U)) \subseteq W_i$, so we can consider the restriction of s_{ij_i} . If we set $U = \operatorname{Spec} R$ (possibly after replacing U by a smaller affine non-empty subset) and $p^{-1}(U) = \operatorname{Spec} B$, then we have $\Gamma(p^{-1}(U) \times_U p^{-1}(U), \mathcal{O}) = B \otimes_R B$ and we may consider the generic part of the cocycle in this tensor product.

So we may assume now that R is a noetherian domain, $R \subseteq B$ is finite and flat and that we have an element $s \in B \otimes_R B$ which maps to 0 in $\Gamma((Y \times_X Y \times_X Y)_{\text{surj}}, \mathcal{O}_{\text{surj}})$. With a similar argument as in the proof of Lemma 3.7.1 we find an affine constructible partition of $\biguplus X_i \rightarrow X$ such that the $Y_i \times_{X_i} Y_i \times_{X_i} Y_i$ are reduced. But then the image of s is not only 0 in the surjective topology, but already in $B_i \otimes_{R_i} B_i \otimes_{R_i} B_i$. Then s comes from the left by the exactness of the Čech complex in the flat topology. $\square$

Corollary 3.10.3. *Let X be a noetherian scheme of finite dimension over a field of characteristic 0 endowed with the surjective topology. Let M be a coherent module on X with corresponding module M_{surj} in the surjective topology. Then $\check{H}^1(X, M_{\text{surj}}) = 0$ and $H^1(X, M_{\text{surj}}) = 0$.*

Proof. The vanishing of the first cohomology follows from the vanishing of the Čech cohomology. By Lemma 3.8.4 we may pass to a constructible partition $j_i : X_i \rightarrow X$ such that $j_i^*(M)$ is finitely generated and free. Then this follows from Theorem 3.10.2. $\square$

Corollary 3.10.4. *Let $X = \operatorname{Spec} R$ be an affine noetherian scheme of finite dimension over a field of characteristic 0. Let $0 \rightarrow N \rightarrow M \rightarrow L \rightarrow 0$ be a short exact sequence of finitely generated R -modules with corresponding short exact sequence of sheaves $0 \rightarrow \ker = N^{\text{surj}} \rightarrow M_{\text{surj}} \rightarrow L_{\text{surj}} \rightarrow 0$ on X_{surj} . Then $H^1(X_{\text{surj}}, \ker) = 0$. In the situation of Proposition 2.8.1 we get $H^1(X_{\text{surj}}, \operatorname{Syz}) = 0$.*

Proof. The exactness of $M_{\text{surj}} \rightarrow L_{\text{surj}} \rightarrow 0$ induces by Corollary 3.9.2 a global surjection $\Gamma(X_{\text{surj}}, M_{\text{surj}}) \rightarrow \Gamma(X_{\text{surj}}, L_{\text{surj}})$. Since $H^1(X_{\text{surj}}, M_{\text{surj}}) = 0$ by Corollary 3.10.3, it follows $H^1(X_{\text{surj}}, \ker) = 0$ by the long exact sequence of cohomology.

For the second statement let $R^n \rightarrow N \rightarrow 0$ be given. By Corollary 3.9.2 and Proposition 3.9.5 we get a surjection $\Gamma(X_{\text{surj}}, \mathcal{O}_{\text{surj}}^n) \rightarrow \Gamma(X_{\text{surj}}, N_{\text{surj}}) \rightarrow \Gamma(X_{\text{surj}}, N^{\text{surj}})$. Hence the result follows from $H^1(X_{\text{surj}}, \mathcal{O}_{\text{surj}}^n) = 0$. $\square$

4. FROBENIUS TOPOLOGY AND FROBENIUS CLOSURE

4.1. Frobenius morphism and Frobenius closure.

Let R denote a ring containing a field of positive characteristic p . Then the *Frobenius homomorphism* is given by $F : R \rightarrow R, f \mapsto f^p$. This is a ring homomorphism due to $(f + g)^p = f^p + g^p$ and so are its iterations $F^e : R \rightarrow R, f \mapsto f^{p^e}$, where $q = p^e$. The e -th *Frobenius power* of an ideal $I \subseteq R$ is the extended ideal

$$I^{[q]} = F^e(I) = \{f^q : f \in I\} = (f_1^q, \dots, f_n^q),$$

where $I = (f_1, \dots, f_n)$.

Definition 4.1.1. The *Frobenius closure* of an ideal $I \subseteq R$ in a commutative ring containing a field of positive characteristic p is given as

$$I^F = \{f \in R : \exists e \in \mathbb{N} \text{ such that } f^q \in I^{[q]}, q = p^e\}.$$

For a ring R of positive characteristic and an R -module M we denote ${}^e M = M \otimes_R {}^e R$, where ${}^e R$ is the ring R , but considered as an R -algebra via the e -th iteration of the

Frobenius homomorphism. For submodules $N \subseteq M$ we get eR -homomorphisms ${}^eN \rightarrow {}^eM$, and the Frobenius closure of a submodule $N \subseteq M$ is defined by

$$N^F = \{s \in M : \exists e \text{ such that } s \otimes 1 \in \text{im}({}^eN \rightarrow {}^eM)\}.$$

We consider the category of schemes of positive characteristic p , that is, the category of schemes X with a (unique) structure morphism $X \rightarrow \text{Spec } \mathbb{Z}/(p)$. The *absolute Frobenius morphism* is the identity on the underlying set X and on every open affine subset $\text{Spec } R = U \subseteq X$ it is the scheme morphism given by the Frobenius homomorphism. We denote the Frobenius by ${}^1X \rightarrow X$ and the e -th iteration by ${}^eX \rightarrow X$.

Definition 4.1.2. A (finite) morphism $V \rightarrow U$ of finite type of schemes of positive characteristic p is called a (finite) *Frobenius cover* if there exists $e \in \mathbb{N}$ and a factorization ${}^eU \rightarrow V \rightarrow U$ such that ${}^eU \rightarrow V$ is surjective and the composition is an iteration of an absolute Frobenius morphism.

Lemma 4.1.3. *The composition of two (finite) Frobenius covers and an arbitrary base change of a (finite) Frobenius cover is again a (finite) Frobenius cover.*

Proof. Let ${}^eU \rightarrow V \rightarrow U$ and ${}^{\tilde{e}}V \rightarrow W \rightarrow V$ be given. Then we also have ${}^{e+\tilde{e}}U \rightarrow {}^{\tilde{e}}V$ and so ${}^{e+\tilde{e}}U \rightarrow {}^{\tilde{e}}V \rightarrow W \rightarrow V \rightarrow U$. If $Y \rightarrow X$ is a base change, then the pull-back of ${}^eU \rightarrow V \rightarrow U$ and the universal property of the product give ${}^eY \times_{eX} {}^eU \rightarrow Y \times_X {}^eU \rightarrow Y \times_X V \rightarrow Y \times_X U$. $\square$

Lemma 4.1.4. *Let X denote a scheme in positive characteristic, $e \in \mathbb{N}$. Then the projections ${}^eX \times_X {}^eX \rightarrow {}^eX$ induce (identical) isomorphisms in the reductions, $({}^eX \times_X {}^eX)_{\text{red}} \cong ({}^eX)_{\text{red}}$. The same is true for a finite Frobenius cover $Y \rightarrow X$, and the diagonal $Y \rightarrow Y \times_X Y$ is also a finite Frobenius cover.*

Proof. We may assume that $X = \text{Spec } R$ is affine. We have the diagonal mapping ${}^eX \xrightarrow{\Delta} {}^eX \times_X {}^eX \xrightarrow{p_1} {}^eX$ resp. ${}^eR \xrightarrow{i_1} {}^eR \otimes_R {}^eR \xrightarrow{\Delta^*} {}^eR$, $f \mapsto f \otimes 1$, $a \otimes b \mapsto ab$. The composition is of course the identity. The other composition sends $a \otimes b \mapsto ab \otimes 1$. We have $(a \otimes b)^q = a^q \otimes b^q = a^q b^q \otimes 1$, because $b^q \in R$. Hence $a \otimes b - ab \otimes 1$ is nilpotent in ${}^eR \otimes_R {}^eR$, and so after reduction we have an isomorphism. The two projections yield the same isomorphism, because the inverse is Δ_{red} in both cases.

If S is finite over R with a spec-surjective homomorphism $S \rightarrow {}^eR$, then again we look at $S \otimes_R S \xrightarrow{\Delta^*} S \xrightarrow{i_1} S \otimes_R S$. The difference between $a \otimes b$ and $ab \otimes 1$ maps to a nilpotent element under the finite spec-surjective homomorphism $S \otimes_R S \rightarrow {}^eR \otimes_R {}^eR$. Since this homomorphism is spec-surjective, the difference itself must be nilpotent.

For the last statement we look at $S \otimes_R S \xrightarrow{\Delta^*} S \xrightarrow{i_1} S \otimes_R S \rightarrow {}^e(S \otimes_R S)$. The composition maps $a \otimes b \mapsto a^q b^q \otimes 1$, and for e big enough this equals $(a \otimes b)^q$, as the difference $a \otimes b - ab \otimes 1$ is nilpotent. $\square$

4.2. Frobenius topology.

Definition 4.2.1. The *Frobenius topology* on a scheme X of characteristic p is given by schemes over X where the morphisms $V \rightarrow U$ (including the structural morphisms $V \rightarrow X$) are finite Frobenius covers. We denote X endowed with the Frobenius topology by X_{frob} .

Proposition 4.2.2. *The Frobenius topology is an affine single-handed covering Grothendieck topology. Every scheme morphism $Y \rightarrow X$ of schemes over $\mathrm{Spec} \mathbb{Z}/(p)$ yields a site morphism $Y_{\mathrm{frob}} \rightarrow X_{\mathrm{frob}}$.*

Proof. This follows from Lemma 4.1.3. $\square$

Proposition 4.2.3. *The Frobenius closure is the closure operation induced by the Frobenius topology.*

Proof. This follows from the Definitions 4.1.1 and 4.2.1 and from Lemma 2.4.4. Just note that if $s \otimes 1 \in \mathrm{im}(N \otimes_R {}^e R \rightarrow M \otimes_R {}^e R)$, then there exists also a finite subalgebra $R \rightarrow S \rightarrow {}^e R$ such that $s \otimes 1 \in \mathrm{im}(N \otimes_R S \rightarrow M \otimes_R S)$. $\square$

Proposition 4.2.4. *Let R denote a ring containing a field of positive characteristic p . Then the following are equivalent.*

- (i) *R is F -pure, that is, the Frobenius homomorphism is a pure ring homomorphism.*
- (ii) *The Frobenius topology on $\mathrm{Spec} R$ is pure.*
- (iii) *The Frobenius closure is trivial on arbitrary submodules.*
- (iv) *If R is a local Gorenstein ring, then this is also equivalent with the property that the Frobenius closure is trivial on ideals.*

Proof. The equivalence (i) $\Leftrightarrow$ (ii) follows immediately from Definition 4.2.1 of the Frobenius topology and from Definition 2.5.1. The equivalence of (ii) and (iii) follows from Proposition 2.5.3. (iii) implies (iv). Under the Gorenstein hypothesis (iv) implies (iii) by [14, Corollary 1.5]. $\square$

F -pure rings have found a great deal of interest, including their classification in low dimension, see e.g. [37], [15], [14], [48], [25], [26].

4.3. Rings of global sections in the Frobenius topology.

The *perfect closure* of a ring R over a field of positive characteristic p is given as the colimit $R^\infty = \varinjlim_{e \in \mathbb{N}} {}^e R$, see [63]. For an R -module M we have $\varinjlim {}^e M = \varinjlim M \otimes_R {}^e R = M \otimes_R R^\infty$. We will see that this is also the module of global sections of M_{frob} in the Frobenius topology.

Proposition 4.3.1. *Let R^∞ be the perfect closure of R . The fixation given by R^∞ defines an absolute filter with absolute stalk R^∞ .*

Proof. If $R \rightarrow S$ is a Frobenius cover and $S \rightarrow R^\infty$ an R -algebra homomorphism, then this factors through some ${}^e R$. For a Frobenius cover $S \rightarrow T$ over R and a given fixation $S \rightarrow {}^e R \rightarrow R^\infty$ there exists by definition $T \rightarrow {}^e S \rightarrow {}^{e+e} R$. Hence there exists also an S -algebra homomorphism $T \rightarrow R^\infty$. Hence this quasifilter is absolute (Definition 1.9.1). The image of S under $\varphi : S \rightarrow {}^e R \rightarrow R^\infty$ gives also a Frobenius cover (over $\mathrm{Spec} R$ and $\mathrm{Spec} S$), since the kernel of φ is nilpotent as φ is spec-surjective. Hence by Proposition 1.8.6 it is a filter and R^∞ is the absolute stalk. $\square$

Remark 4.3.2. For a scheme X of positive characteristic such that the Frobenius morphism is finite (F -finite schemes), the Frobenius powers ${}^e X \rightarrow X$, $e \in (\mathbb{N}, \geq)$ belong to the category and they also form an absolute filter with absolute stalk R^∞ .

Theorem 4.3.3. *Let R denote a ring of positive characteristic p endowed with the Frobenius topology, $X_{\text{frob}} = (\text{Spec } R)_{\text{frob}}$. Let M denote an R -module with corresponding sheaf M_{frob} in the Frobenius topology. Then for every finite Frobenius cover (i.e. for every object in $\mathcal{T}_X$) $Y \rightarrow X$ we have $\Gamma(Y_{\text{frob}}, M_{\text{frob}}) = M \otimes_R R^\infty$. In particular, $\Gamma(X_{\text{frob}}, \mathcal{O}_{\text{frob}}) = R^\infty$.*

Proof. Let $Y = \text{Spec } S$. By Proposition 4.3.1 and Corollary 1.9.6 we have a natural homomorphism $\Gamma(Y, M_{\text{frob}}) \rightarrow \varinjlim_{\lambda \in \Lambda} \Gamma(U_\lambda, M_{\text{frob}}) = M \otimes_R R^\infty$, where Λ is the indexing category corresponding to the fixation given by the perfect closure of R . Let $f \in {}^e R$ be represented (e sufficiently large) by $f \in S'$ finite over S , $R \rightarrow S \rightarrow S' \rightarrow {}^e R$ finite. By Lemma 4.1.4 the element $f \otimes 1 - 1 \otimes f$ is nilpotent in $S' \otimes_S S'$. But nilpotent elements vanish in some Frobenius power, so f defines a global element in $\Gamma(Y, \mathcal{O}_{\text{frob}})$. It follows that $\Gamma(Y_{\text{frob}}, \mathcal{O}_{\text{frob}}) \rightarrow R^\infty$ is surjective and then also that $\Gamma(Y_{\text{frob}}, M_{\text{frob}}) \rightarrow M \otimes_R R^\infty$ is surjective. On the other hand, if $s \in \Gamma(Y_{\text{frob}}, M_{\text{frob}})$ maps to 0 in the stalk, then it is 0 in some $M \otimes_R {}^e R$ and also in a finite Frobenius cover $S \rightarrow S'$, so s itself must be 0. $\square$

Remark 4.3.4. Note that already for a (non-perfect) field K the ring of global sections in the Frobenius topology is not the field itself, as it is in the flat topology, though the Frobenius homomorphism on a field is flat. This rests on the fact that the Frobenius homomorphism on ${}^e K \otimes_K {}^e K$, which enters the computation of the global section ring in the Frobenius topology, is not flat.

4.4. Exactness in the Frobenius topology.

Proposition 4.4.1. *Let R denote a ring of positive characteristic and let $N \xrightarrow{\alpha} M \xrightarrow{\beta} L$ denote a complex of R -modules. Let $N_{\text{frob}} \xrightarrow{\alpha} M_{\text{frob}} \xrightarrow{\beta} L_{\text{frob}}$ denote the corresponding complex of sheaves on $(\text{Spec } R)_{\text{frob}}$. Then the following are equivalent.*

- (i) *The complex of sheaves $N_{\text{frob}} \xrightarrow{\alpha} M_{\text{frob}} \xrightarrow{\beta} L_{\text{frob}}$ is exact in the Frobenius topology.*
- (ii) *The complex $N \otimes_R R^\infty \xrightarrow{\alpha} M \otimes_R R^\infty \xrightarrow{\beta} L \otimes_R R^\infty$ is exact (this is the global complex as well as the corresponding complex in the absolute stalk given by R^∞).*
- (iii) *For every $e \in \mathbb{N}$ the complex ${}^e N \xrightarrow{\alpha} {}^e M \xrightarrow{\beta} {}^e L$ has the property that $\ker(\beta) \subseteq (\text{im } \alpha)^F$.*
- (iv) *For every $e \in \mathbb{N}$ and every $s \in M \otimes_R {}^e R$ mapping to 0 in $L \otimes_R {}^e R$ there exists $\tilde{e} \geq e$ and an element $t \in N \otimes_R {}^{\tilde{e}} R$ mapping to the image of s in $M \otimes_R {}^{\tilde{e}} R$.*

Proof. The equivalence (i) $\Leftrightarrow$ (ii) was given in Proposition 1.10.4. The equivalence (ii) $\Leftrightarrow$ (iii) (for which (iv) is an explicit version) follows directly and follows also in the F -finite case from Corollary 2.7.2. $\square$

Example 4.4.2. The sequence $0 \rightarrow R \rightarrow R_{\text{red}}$ is exact in the Frobenius topology.

Example 4.4.3. Let $R = K[x, y, v]/(x^3 + y^3 + v^3)$, where K is a field of positive characteristic $p \neq 2, 3$. It is known that $v^2 \in (x, y)^F$ if and only if $p = 2 \bmod 3$, and in this case already $(v^2)^p \in (x^p, y^p)$ [39, Example 2.2]. Look at the complex of R -modules

$$R^2 \xrightarrow{x, y} R \rightarrow R/(x, y, v^2).$$

This complex is of course not exact as R -modules. For $p = 2 \bmod 3$ it is however Frobenius-exact, since $(v^2)^p$ is in the image of x^p and y^p . Therefore $(x, y)R^\infty = (x, y, v^2)R^\infty$ and so it is exact. It is not Frobenius exact for $p = 1 \bmod 3$.

Example 4.4.4. Consider the domain $R = K[x, y, u]/(y^p - ux^p)$ with normalization $K[y/x, x]$ and the ideal $I = (x, y)$. The first Frobenius power of I is $I^{[p]} = (x^p, y^p) = (x^p)$, so it is the principal ideal generated by x^p . The ideal I is as an R -module isomorphic to $R^2/((y, -x), (ux^{p-1}, -y^{p-1})) \cong I$, where $e_1 \mapsto x$, $e_2 \mapsto y$. Therefore we get isomorphisms (say p is odd, $q = p^e$)

$$R^2/((y^q, -x^q), (u^q x^{q(p-1)}, -y^{q(p-1)})) \cong I \otimes_R^e R.$$

For $e = 1$ ($q = p$), the element $(u, -1)$ on the left corresponds to $x \otimes u - y \otimes 1$ in $I \otimes_R^1 R$, and the natural mapping $I \otimes_R^1 R \rightarrow I^{[p]}$ sends this to $ux^p - y^p = 0$. The element $x \otimes u - y \otimes 1$ is however not 0 in $I \otimes_R^1 R$ nor does it become 0 in $I \otimes_R^e R$ for some e . This is because $(u, -1) \in I \otimes_R^1 R$ is mapped to $(u^{q/p}, -1^{q/p})$ in $I \otimes_R^e R$, but this does not belong to $((y^q, -x^q), (u^q x^{q(p-1)}, -y^{q(p-1)}))$. Therefore $0 \rightarrow I \rightarrow R$ is exact, but not Frobenius-exact in the sense of Proposition 4.4.1.

4.5. Cohomology in the Frobenius topology.

Proposition 4.5.1. *Let R be a ring of positive characteristic, $X_{\text{frob}} = \text{Spec } R$ endowed with the Frobenius topology, and let M denote an R -module with corresponding Frobenius sheaf M_{frob} on X_{frob} . Then $\check{H}^k(X_{\text{frob}}, M_{\text{frob}}) = 0$ for $k \geq 1$ and in particular $H^1(X_{\text{frob}}, M_{\text{frob}}) = 0$.*

Proof. For every finite Frobenius cover $Y \rightarrow X$ we have $\Gamma(Y, M_{\text{frob}}) = M \otimes_R R^\infty$ by Theorem 4.3.3. In particular we have $\Gamma(Y \times_X \cdots \times_X Y, M_{\text{frob}}) = M \otimes_R R^\infty$, and all restriction morphisms are isomorphisms. Hence the Čech complex is

$$\Gamma(X_{\text{frob}}, M_{\text{frob}}) = R^\infty \otimes_R M \xrightarrow{\text{id}} R^\infty \otimes_R M \xrightarrow{0} R^\infty \otimes_R M \xrightarrow{\text{id}} R^\infty \otimes_R M \rightarrow \cdots.$$

Therefore $\check{H}^k(R_{\text{frob}}, M_{\text{frob}}) = 0$ for $k \geq 1$ and $H^1(R_{\text{frob}}, M_{\text{frob}}) = 0$ by Lemma 1.11.1. $\square$

Corollary 4.5.2. *Let R be a ring of positive characteristic. Let $0 \rightarrow N \rightarrow M \rightarrow L \rightarrow 0$ be a short exact sequence of R -modules with corresponding short exact sequence of sheaves $0 \rightarrow \text{kern} = N^{\text{frob}} \rightarrow M_{\text{frob}} \rightarrow L_{\text{frob}} \rightarrow 0$ on X_{frob} . Then $H^1(X_{\text{frob}}, N^{\text{frob}}) = 0$. In the situation of Proposition 2.8.1 we get $H^1(X_{\text{frob}}, \text{Syz}) = 0$.*

Proof. The exactness of $M_{\text{frob}} \rightarrow L_{\text{frob}} \rightarrow 0$ induces by Proposition 4.4.1 the global exactness $\Gamma(X_{\text{frob}}, M_{\text{frob}}) = M \otimes_R R^\infty \rightarrow \Gamma(X_{\text{frob}}, L_{\text{frob}}) = L \otimes_R R^\infty \rightarrow 0$. Since $H^1(X_{\text{frob}}, M_{\text{frob}})$ is 0 by Proposition 4.5.1, it follows $H^1(X_{\text{frob}}, N^{\text{frob}}) = 0$ by the long exact cohomology sequence.

For the second statement let $N \subseteq M$ be finitely generated R -modules and let $R^n \rightarrow N$ be a surjection. Then $\Gamma(X_{\text{frob}}, \mathcal{O}_{\text{frob}}^n) = (R^\infty)^n \rightarrow \Gamma(X_{\text{frob}}, N_{\text{frob}}) = N \otimes_R R^\infty$ is a surjection but also $N \otimes_R R^\infty \rightarrow \Gamma(X_{\text{frob}}, N^{\text{frob}}) \subseteq \Gamma(X_{\text{frob}}, M_{\text{frob}}) = M \otimes_R R^\infty$ is a surjection, as $\Gamma(X_{\text{frob}}, N^{\text{frob}})$ consists of the elements in $\Gamma(X_{\text{frob}}, M_{\text{frob}})$ which are in the Frobenius closure of N . Hence $\Gamma(X_{\text{frob}}, \mathcal{O}_{\text{frob}}^n) \rightarrow \Gamma(X_{\text{frob}}, N^{\text{frob}})$ is surjective and so again $H^1(X_{\text{frob}}, \text{Syz}) = 0$. $\square$

5. FINITE TOPOLOGY AND PLUS CLOSURE

5.1. Plus closure for ideals and for submodules.

The *plus* (or *finite*) *closure* of an ideal $I \subseteq R$ in a domain R is given by

$$I^+ = \{f \in R : \exists R \subseteq S \text{ finite domain such that } f \in IS\}.$$

The properties of this closure operation depend heavily on the characteristic. If R contains a field of characteristic zero and is normal, then $I^+ = I$ (this is due to the trace map, see [10, Remark 9.2.4]). If however R contains a field of positive characteristic, then we have $I^F \subseteq I^+ \subseteq I^*$, where I^F is the Frobenius closure (Section 4.1) and where I^* is the tight closure [38], [35]. Moreover, it is conjectured that $I^+ = I^*$ [34]. The plus closure may also be characterized with the help of the so-called *absolute integral closure* R^+ of R . This is the integral closure of R inside the algebraic closure $\overline{Q(R)}$ of its fraction field (first considered by M. Artin in [2]). The plus closure is then just $I^+ = R \cap IR^+$ [38, Theorem 1.7]. R^+ is a big Cohen-Macaulay algebra for excellent local domains of positive characteristic [38, Theorem 7.0], [40] or [36].

We first generalize this notion for rings which are not domains and for arbitrary submodules. For this we introduce finite covers.

Definition 5.1.1. A scheme morphism $Y \rightarrow X$ is called a *finite cover* if it is finite and surjective.

Definition 5.1.2. For R -submodules $N \subseteq M$ we define the *finite closure* of N in M by

$$N^+ = \{s \in M : \exists R \rightarrow S \text{ a finite cover such that } s \otimes 1 \in \text{im}(N \otimes_R S \rightarrow M \otimes_R S)\}.$$

This definition coincides for an ideal in a domain with the standard definition due to Proposition 5.1.3(v).

Proposition 5.1.3. *Finite covers have the following properties.*

- (i) *The composition of finite covers is a finite cover.*
- (ii) *The base change of a finite cover is a finite cover.*
- (iii) *The reduction $X_{\text{red}} \rightarrow X$ is a finite cover.*
- (iv) *If X_i are the integral components of X , then $\biguplus_{i \in I} X_i \rightarrow X$ is a finite cover.*
- (v) *If X is integral and $Y \rightarrow X$ is a finite cover, then there exists an integral component Y_0 of Y such that $Y_0 \rightarrow X$ is a finite cover. This means that every finite cover of an integral scheme can be refined by an integral finite cover.*

Proof. These are clear. For (v) let $y \in Y$ map to the generic point of X and let $y \in Y_0 \subseteq Y$ be an integral component. Then $Y_0 \rightarrow X$ is finite but also surjective by going-up [12, Proposition 4.15]. $\square$

Definition 5.1.4. Let X denote a scheme. The *finite topology* $\mathcal{T}_X$ on X is given by the schemes over X , where all morphisms are finite and where the finite covers are covers. The structural morphisms are declared to be covers. We denote X endowed with this topology by X_{fin} .

Proposition 5.1.5. *The finite topology is an affine single-handed Grothendieck topology. Every scheme morphism $Y \rightarrow X$ yields a corresponding site morphism $Y_{\text{fin}} \rightarrow X_{\text{fin}}$. The closure induced by the finite topology is the plus closure.*

Proof. This is clear from Proposition 5.1.3 and Lemma 2.4.4. $\square$

Remark 5.1.6. In the *qfh-topology* introduced by Voevodsky ([68], [64]) a cover is given by a topological epimorphism (or universal submersion) $\biguplus X_i \rightarrow X$ such that each $X_i \rightarrow X$ is a quasi-finite morphism [18, Définition 6.2.3]. So this quasifinite topology is a refinement of the finite topology. On the other hand, over a complete local domain it follows from [18, Corollaire 6.2.6] that a quasifinite cover contains a connected component

which is a finite cover of the X (this is not true for a quasifinite surjection without the submersive property).

5.2. The ring of global sections in the finite topology.

Definition 5.2.1. Let R denote a ring and let $R \rightarrow L$ be an R -algebra. Then we set

$$R^{\text{comp}/L} = \{f \in L : \text{there exists a finite } R\text{-algebra } T \subseteq L \text{ such that} \\ f \in T \text{ and } f \otimes 1 - 1 \otimes f \in T \otimes_R T \text{ is nilpotent}\}$$

and call it the *compatible integral closure* of R in L .

Compare [45, Section I] for this definition. By the *compatible normalization* of a domain R we mean the compatible integral closure of R in $L = Q(R)^\infty$, and denote it by R^{comp} . Recall that the *seminormalization* of a ring is the biggest ring inside the normalization which does not separate points and does not change the residue class fields, see [43, Definition 7.2.1]. The following characterization of the seminormalization is due to Manaresi, see [45, Theorem I.6]. We include a proof.

Lemma 5.2.2. *Let R denote a noetherian domain in characteristic zero. Then the seminormalization is the compatible normalization.*

Proof. The seminormalization $\text{Spec } R^{\text{semi}} \rightarrow \text{Spec } R$ is a homeomorphism and induces isomorphisms $\kappa(x) \rightarrow \kappa(x')$ for $x \in \text{Spec } R$. The same is true for $R \subseteq S \subseteq R^{\text{semi}}$. Let $f \in S \subseteq R^{\text{semi}}$ be given, $R \subseteq S$ finite, and consider $f \otimes 1 - 1 \otimes f \in S \otimes_R S \rightarrow R^{\text{semi}} \otimes_R R^{\text{semi}}$. For every ring homomorphism $R \rightarrow \kappa(x)$ there exists exactly one ring homomorphism $R^{\text{semi}} \rightarrow \kappa(x)$. Therefore by Lemma 1.6.5 we know that $f \otimes 1 - 1 \otimes f$ is nilpotent (in $S \otimes_R S$ and in $R^{\text{semi}} \otimes_R R^{\text{semi}}$). Hence $f \in R^{\text{comp}}$.

Now suppose that an element f in the normalization is compatible. We know by Lemma 1.6.6 that it has a uniquely determined value in $\kappa(x)$ at each point $x \in \text{Spec } R$. Therefore f belongs to the seminormalization. $\square$

Proposition 5.2.3. *Let R denote a noetherian domain, $X = \text{Spec } R$. Then the ring of global sections in the finite topology is the compatible normalization inside the perfect closure $Q(R)^\infty$, i.e. $\Gamma(X_{\text{fin}}, \mathcal{O}_{\text{fin}}) = R^{\text{comp}}$.*

Proof. The natural morphism $(\text{Spec } Q(R))_{\text{fin}} \rightarrow X_{\text{fin}}$ induces the ring homomorphism $\Gamma(X_{\text{fin}}, \mathcal{O}_{\text{fin}}) \rightarrow \Gamma(Q(R)_{\text{fin}}, \mathcal{O}_{\text{fin}})$. First note that $\Gamma(Q(R)_{\text{fin}}, \mathcal{O}_{\text{fin}}) = Q(R)^\infty$, by Theorem 4.3.3 and Proposition 3.4.1.

An element $f \in \Gamma(X, \mathcal{O}_{\text{fin}})$ is represented by $f \in S$, where $R \rightarrow S$ is a finite cover. The compatibility condition for the sheafification means exactly that $f \otimes 1 - 1 \otimes f \in S \otimes_R S$ is nilpotent. Hence the natural ring homomorphism maps into the compatible normalization.

Suppose that $f \in R^{\text{comp}} \subseteq Q(R)^\infty$ and that it is represented by $f \in S$, $R \rightarrow S$ a finite cover. Then $f \otimes 1 - 1 \otimes f \in S \otimes_R S$ is nilpotent. Since reductions are finite covers (Proposition 5.1.3(iii)), this means that $f \in S$ is compatible in the sense of sheafification and gives a global element in $\Gamma(Y, \mathcal{O}_{\text{fin}})$. Hence the homomorphism is surjective.

Suppose that $f \in \Gamma(X_{\text{fin}}, \mathcal{O}_{\text{fin}})$ maps to 0. We may assume that f is represented by $f \in S$, where S is a domain (Proposition 5.1.3(v)). Hence the image of f in $\Gamma(Q(R), \mathcal{O}_{\text{fin}})$ is represented by f considered in $Q(S)$. So $f = 0$. $\square$

Remark 5.2.4. The seminormalization is also the ring of global sections in the submersive topology resp. h -topology (see [68, Proposition 3.2.10] and [4]) The normalization cannot be the ring of global sections for a Grothendieck topology which is functorial for all scheme morphisms, since the normalization is not functorial for quotient rings, see the following example.

Example 5.2.5. Let $R \subset \mathbb{C}[[x]]$ denote the subring consisting of formal power series with constant term in $\mathbb{R}$. Then $\frac{x^2}{x} = i$ and $i^2 = -1 \in R$, hence i is both rational and integral over R and so $\mathbb{C}[[x]]$ is the normalization of R . The residue class field of R with respect to the maximal ideal $\mathfrak{m}_R = (x, ix)$ is the (normal) field $\mathbb{R}$, but there does not exist an R -algebra homomorphism between the normalizations $\mathbb{C}[[x]] \rightarrow \mathbb{R}$. In particular i is not compatible, and $\Gamma(R_{\text{fin}}, \mathcal{O}_{\text{fin}}) = R$.

5.3. Absolute stalks in the finite topology.

There are several relevant absolute filters in the finite topology to study. Already for an integral scheme it is not clear which one is the best choice.

Proposition 5.3.1. *Let R denote a noetherian ring with integral components R_i , $i \in I$. Set $R^+ := \prod_{i \in I} (R_i)^+$. Then $\prod_{i \in I} \overline{Q(R_i)}$ and also R^+ define an absolute filter in $X_{\text{fin}} = (\text{Spec } R)_{\text{fin}}$ with absolute stalk R^+ .*

Proof. By Proposition 5.1.3(v) we may assume that R is a domain, because we can treat the components separately. Every ring homomorphism $S \rightarrow \overline{Q(R)}$, S a finite cover, factors uniquely through R^+ , so we restrict to the fixation given by R^+ .

Note first that for every finite cover $R \rightarrow S$ there exists an R -algebra homomorphism $S \rightarrow R^+$. According to Proposition 5.1.3(v) there exists a homomorphism $S \rightarrow S_0$ to a domain which is a finite extension of R . Hence we may assume that S is a domain. Then $Q(R) \rightarrow Q(S)$ is a finite field extension and hence there exists an inclusion $Q(S) \subseteq \overline{Q(R)}$. This gives an inclusion $S \subseteq R^+$. So in the quasifilter given by the fixation with R^+ every finite cover occurs. It is a filter by Proposition 1.8.6, since the image of a finite cover S in R^+ is itself a finite (integral) cover over R .

To show that it is absolute let $S \rightarrow R^+$ be an R -algebra homomorphism and let $S \rightarrow T$ be in $\mathcal{T}_R$. Then let $\overline{S}$ be the image of S in R^+ and let $\overline{T}$ be a T -algebra which is a finite domain over $\overline{S}$. Then there exists an $\overline{S}$ -embedding of $\overline{T}$ into R^+ . This gives $T \rightarrow R^+$ over S . The stalk in this absolute filter is of course R^+ . $\square$

Example 5.3.2. It is also useful to consider the filter where the objects are only the domains S with a fixation $R \rightarrow S \rightarrow R^+$. This is a quasifilter since for two finite domains $R \subseteq S, S'$ there exists a common finite domain in R^+ containing both. It is again an absolute filter by similar arguments as in the proof of Proposition 5.3.1. The only morphisms in this filter are inclusion of domains. Moreover, it is a covering filter (Remark 1.9.2), since every finite R -homomorphism $S \rightarrow T$ is a cover when S is a domain (because $T \in \mathcal{T}_R$ is a finite cover over R).

Example 5.3.3. Consider the affine line $\mathbb{A}_K^1 = \text{Spec } R$, $R = K[x]$ in the finite topology and consider the fixation given by $R^+ \oplus K^+$ (where in the second component x is sent to 0). This is again an absolute filter with absolute stalk $R^+ \oplus K^+$. The sequence $0 \rightarrow R \xrightarrow{x} R$ is not exact in this absolute filter, since it is not exact in K^+ . Proposition 1.10.4(i)-(v) do not hold, but (vi) and (vii) hold, since for given Y the sequence is not $\mathcal{T}$ -exact in general,

but its restriction to an integral component of dimension one is. Note that these inclusions belong to the category, but are not always indicated by the filter (e.g. $\mathbb{A}^1 \subset \mathbb{A}^1 \uplus \{x\}$ is not indicated).

Proposition 5.3.4. *Let R denote a noetherian ring with integral components R_i , $i \in I$. Let $N \subseteq M$ be R -modules, $s \in M$. Then the following are equivalent.*

- (i) $s \in N^+$.
- (ii) $s \in N_i^+$, where N_i is the image of $N \otimes_R R_i$ inside $M_i = M \otimes_R R_i$.
- (iii) *There exists a finite and spec-surjective ring homomorphism $R \rightarrow S$ such that $s \otimes 1 \in \text{im}(N \otimes_R S \rightarrow M \otimes_R S)$.*
- (iv) $s \in N^{\text{top-cl}}$, where top-cl denotes the closure operation induced by the finite topology on $\text{Spec } R$.
- (v) $s \otimes 1 \in \text{im}(N \otimes_R R^+ \rightarrow M \otimes_R R^+)$, where R^+ is the absolute stalk.

Proof. (i) and (iii) are equivalent by Definition 5.1.2. The defining property (iii) might be checked on the components by Proposition 5.1.3, hence (i) and (ii) are equivalent. The equivalence of (iii) and (iv) was mentioned in Proposition 5.1.5. The equivalence of (iv) and (v) follows from Corollary 2.4.10 and Proposition 5.3.1. $\square$

Remark 5.3.5. The exactness of a complex in the finite topology is a strong condition, since $X \cup \{x\} \rightarrow X$ is a finite cover, where x is a closed point of X . If a sheaffied complex is exact on X_{fin} , then this implies that it is exact on every closed point! Hence the exactness in the finite topology for a scheme of finite type over a field implies the exactness in the surjective topology (Theorem 3.9.1 and Remark 3.9.4). The converse is not true, as $R^n \xrightarrow{f_1, \dots, f_n} R \rightarrow R/\text{rad}(I) \rightarrow 0$ shows, where $I = (f_1, \dots, f_n) \subseteq I^+ \subset \text{rad}(I)$.

We characterize now the exactness in the absolute filter given by R^+ .

Proposition 5.3.6. *Let R denote a domain endowed with the finite topology and let $N \xrightarrow{\alpha} M \xrightarrow{\beta} L$ denote a complex of R -modules. Let $N_{\text{fin}} \xrightarrow{\alpha} M_{\text{fin}} \xrightarrow{\beta} L_{\text{fin}}$ denote the corresponding complex of sheaves on $(\text{Spec } R)_{\text{fin}}$. Then the following are equivalent.*

- (i) *The complex of sheaves $N_{\text{fin}} \xrightarrow{\alpha} M_{\text{fin}} \xrightarrow{\beta} L_{\text{fin}}$ is exact in the absolute filter given by R^+ .*
- (ii) *The complex of sheaves $N_{\text{fin}} \xrightarrow{\alpha} M_{\text{fin}} \xrightarrow{\beta} L_{\text{fin}}$ is exact in the integral absolute filter given by R^+ where only integral rings occur (Example 5.3.2).*
- (iii) *The complex $N \otimes_R R^+ \xrightarrow{\alpha} M \otimes_R R^+ \xrightarrow{\beta} L \otimes_R R^+$ is exact.*
- (iv) *For every finite cover $R \rightarrow S \rightarrow R^+$ and every $s \in M \otimes_R S$ mapping to 0 in $L \otimes_R S$ there exists a finite S -algebra $T \rightarrow R^+$ (which might be chosen to be integral) such that there exists $t \in N \otimes_R T$ mapping to the restriction of s .*
- (v) *The complex of sheaves $N_{\text{fin}} \xrightarrow{\alpha} M_{\text{fin}} \xrightarrow{\beta} L_{\text{fin}}$ is $\mathcal{T}$ -exact in the finite topology on every integral object.*
- (vi) *For every finite cover $R \rightarrow S$ and every $s \in M \otimes_R S$ mapping to 0 in $L \otimes_R S$ there exists a morphism $S \rightarrow T$ in $\mathcal{T}_R$ and $t \in N \otimes_R T$ mapping to the restriction of s .*
- (vii) *For every finite cover $R \rightarrow S$ with S integral the complex $N \otimes_R S \xrightarrow{\alpha} M \otimes_R S \xrightarrow{\beta} L \otimes_R S$ has the property that $\text{kern}(\beta) \subseteq (\text{im } \alpha)^+$.*

Proof. This follows from Proposition 1.10.4 applied to the (integral) absolute filter given by R^+ . The equivalence of (v) and (vii) was proven in Proposition 2.7.1. $\square$

5.4. Cohomology in the finite topology.

The computation of cohomology in the finite topology, e.g. of the structure sheaf, provides us with several difficulties. So we give here only a result concerning the normalization of a curve and some examples.

Proposition 5.4.1. *Let $C = \operatorname{Spec} R$ be an affine integral curve over an algebraically closed field K and let $C^{\operatorname{nor}} \rightarrow C$ its normalization. Then $\check{H}^1(C^{\operatorname{nor}} \rightarrow C, \mathcal{O}_{\operatorname{fin}}) = 0$.*

Proof. Let $S \subset C$ be the set of singular points. For $P \in S$ let $P_1, \dots, P_{n_P}$ be the points in C^{nor} mapping to P . Then

$$C^{\operatorname{nor}} \times_C C^{\operatorname{nor}} = C^{\operatorname{nor}} \uplus \{P_{ij}, 1 \leq i, j \leq n_P, i \neq j : P \in S\}$$

(the points P_{ii} are the points $P_i \in C^{\operatorname{nor}}$), and similarly $C^{\operatorname{nor}} \times_C C^{\operatorname{nor}} \times_C C^{\operatorname{nor}} = C^{\operatorname{nor}} \uplus \{P_{ijk}\}$. The indices of the points show where to one has to project, e.g. $q_1(P_{ijk}) = p_{2,3}(P_{ijk}) = P_{jk}$. Let now a cocycle $h \in R^{\operatorname{nor}} \otimes_R R^{\operatorname{nor}}$ (this is seminormal) be given (Section 1.11). On the diagonal $\Delta = C^{\operatorname{nor}}$ the cocycle condition gives immediately $h|_{C^{\operatorname{nor}}} = 0$. On the isolated points the cocycle condition gives

$$(q_1^* - q_2^* + q_3^*)(h)(P_{ijk}) = h(P_{jk}) - h(P_{ik}) + h(P_{ij}) = 0.$$

For every $P \in S$ this is exactly the cocycle condition for the complex $K^I \rightarrow K^{I^2} \rightarrow K^{I^3}$, where $I = \{P_1, \dots, P_{n_P}\}$ (the Čech complex of the cover $I \rightarrow \{P\}$). Hence there exist values $h(P_1), \dots, h(P_{n_P})$ such that the differences of the pull-backs under the two projections are $h(P_{ij})$. Since C^{nor} is affine there exists a global function $g \in \Gamma(C^{\operatorname{nor}}, \mathcal{O}_{C^{\operatorname{nor}}})$ such that $g(P_i) = h(P_i)$ for all $P_i, P \in S, i = 1, \dots, n_P$. Then the derivation maps g to h , and so h is a coboundary. $\square$

Example 5.4.2. One of the difficulties in computing cohomology in the finite topology is that the ring of global sections is the seminormalization. This might bring in new cocycles which are not seen on the presheaf level. Let K be an algebraically closed field and consider the (flat) extension $R = K[x] \rightarrow K[y] = S, x \mapsto y^n$. Then $S \otimes_R S = K[y, z]/(y^n - z^n) = K[y, z]/(\prod_{\zeta_n} (y - \zeta_n z))$. This ring consists of n lines in the plane meeting in the origin (say the characteristic does not divide n). So $\operatorname{Spec} S \otimes_R S$ has embedding dimension 2 and is for $n \geq 3$ not seminormal, its seminormalization is the scheme of n axes with embedding dimension n [7, Section 3]. Hence $\Gamma(S \otimes_R S, \mathcal{O}_{\operatorname{fin}}) \neq S \otimes_R S$.

Example 5.4.3. Let $R = K[u, v]/(uv)$ with normalization $R^{\operatorname{nor}} = K[s] \oplus K[t], u = (s, 0)$ and $v = (0, t)$. Consider the principal ideal $I = (g) = (u + v)$ (cf. Example 2.8.2). Then in R^{nor} we have $f = u = (s, 0) = (1, 0)(s, t) = (1, 0)(u + v)$. So $f \in I^+$ (but $f \notin I$). Set $X = \operatorname{Spec} R$ and $Y = \operatorname{Spec} R^{\operatorname{nor}}$. We want to compute $\check{H}^1(Y \rightarrow X, \operatorname{Syz})$, where Syz is given as the kernel of $g : \mathcal{O}_{\operatorname{fin}} \rightarrow I^{\operatorname{fin}} \subseteq \mathcal{O}_{\operatorname{fin}}$ (so Syz is the annihilator of g), and $\delta(f) \in \check{H}^1(Y \rightarrow X, \operatorname{Syz})$ (the cohomology of the structure sheaf is 0 as in Proposition 5.4.1).

$Y \times_X Y$ consists of two lines L_1, L_2 and two points $P_{1,2}$ and $P_{2,1}$. $Y \times_X Y \times_X Y$ consists again of two lines L_1, L_2 and 8 points $P_{i,j,k}$, where $P_{1,1,1} \in L_1$ and $P_{2,2,2} \in L_2$ and where the other six points are isolated components.

The pull-back of the function $g = u + v$ to $Y \times_X Y$ and to $Y \times_X Y \times_X Y$ is the function which is s on L_1 and t on L_2 and has value 0 on the isolated points. So the annihilator of g consists of the functions which are zero on the two lines and have arbitrary values on the isolated points. Let $h \in \Gamma(Y \times_X Y, \text{Syz})$ have values $h(P_{1,2}) = h$ and $h(P_{2,1}) = b$ (and = 0 on the lines). The cocycle condition is $(p_{2,3}^* - p_{1,3}^* + p_{1,2}^*)(h) = 0$. This means evaluated at $P_{1,2,1}$ that $0 = h(P_{2,1}) - h(P_{1,1}) + h(P_{1,2}) = h - 0 + b$ ($P_{2,1,2}$ yields the same, the other points yield no condition). So the cocycle condition reduces to $b = -h$, and since there are no coboundaries at all, we infer that $\tilde{H}^1(Y \rightarrow X, \text{Syz})$ has dimension one.

We come to the computation of $\delta(f)$. In R^{nor} we have $f = qg$, $q = (1, 0)$, so we have to compute the difference between the pull-backs of q under the two projections. The pull back of q under both projections gives on L_1 the function 1 and on L_2 the function 0. The values of the pull backs under the projections at the points are $p_1^*(q)(P_{1,2}) = 1$, $p_1^*(q)(P_{2,1}) = 0$, $p_2^*(q)(P_{1,2}) = 0$, $p_2^*(q)(P_{2,1}) = 1$. Hence the difference is 0 on L_1 and L_2 and has value 1 at $P_{1,2}$ and -1 at $P_{2,1}$. So this difference gives an element in $\Gamma(Y \times_X Y, \text{Syz})$ which represents the cohomology class $\delta(f) \in H^1(X, \text{Syz})$.

6. TOPOLOGIES DEFINED BY ZARISKI FILTERS AND TORSION THEORIES

Some properties of a variety, or objects associated to it, depend only on the points of codimension one, i.e. they are unchanged if we remove a subset of codimension at least two, like e.g. the divisor class group. Other objects get simpler after removing small subsets, e.g. torsion-free sheaves are locally free in codimension one. A (hereditary) torsion theory for R -modules studies which part of a module can be neglected in a certain sense. The notions of a pure resp. parafactorial pair are defined by requiring that the restriction of an étale covering resp. an invertible sheaf to an open subset gives an equivalence of categories ([22, Exposés X.3 and XI.3], [20, Définition 21.13.1]). In birational geometry one may restrict to any non-empty open subset. So smaller open subsets may have the full or the essential information, hence they are covers in a certain sense. We formalize these observations in this part.

6.1. Zariski filters.

Definition 6.1.1. A *Zariski filter* on a scheme X is a *topological filter* F in the Zariski topology, i.e. a non-empty collection of open subsets $V \subseteq X$ such that $V \in F$ and $V \subseteq U$ implies $U \in F$ and such that $V, U \in F$ implies $V \cap U \in F$.

A Zariski filter on X defines for every morphism $\psi : Y \rightarrow X$ a Zariski filter $\psi^*(F)$ on Y by setting $\psi^*(F) = \{W \subseteq Y : W \supseteq \psi^{-1}(V) \text{ for some } V \in F\}$. In particular, for an open subset $U \subseteq X$ this is $F_U = F|_U = \{U \cap V : V \in F\}$ (we will also write $W \in F_U$ if we mean $U \cap W \in F_U$).

Definition 6.1.2. Let X denote a scheme and let F denote a Zariski filter on X . The Grothendieck topology X_{zftop} induced by F is given by all (Zariski-)open subsets of X , and $U_i \rightarrow U$, $i \in I$, is a covering if and only if $\bigcup_{i \in I} U_i \in F_U$, that is, $U \cap V \subseteq \bigcup_{i \in I} U_i$ for some $V \in F$.

Remark 6.1.3. Here we design the Grothendieck topology given by a Zariski filter to be a non-affine topology which is a refinement of the Zariski topology. The open sets $V \in F$ define the single covers of X . This topology is flat, but not faithfully flat. If X and Y

are schemes with Zariski filters F_X and F_Y , then a morphism $\psi : Y \rightarrow X$ induces a site morphism $Y_{\text{zftop}} \rightarrow X_{\text{zftop}}$ if and only if $\psi^*(F_X) \subseteq F_Y$.

Example 6.1.4. Consider a pair (X_0, X) consisting of a scheme X and a subset $X_0 \subseteq X$. All open neighborhoods $V \supseteq X_0$ form a filter denoted by $F(X_0)$. A morphism $\psi : Y \rightarrow X$ which maps $Y_0 \subseteq Y$ into X_0 induces a site morphism. The disjoint union $\bigsqcup_{x \in X_0} \text{Spec } \mathcal{O}_x$ (but also $\bigsqcup_{x \in X_0} \text{Spec } \kappa(x)$) gives an absolute cover (Definition 1.9.3) for this topology.

Example 6.1.5. For a natural number $k \in \mathbb{N}$ and a scheme X denote by $X^{(k)}$ the points of height at most k , that is,

$$X^{(k)} := \{x \in X : \dim \mathcal{O}_{X,x} \leq k\}.$$

Let F_X^k denote the filter consisting of all open subsets which contain $X^{(k)}$. A morphism $\psi : Y \rightarrow X$ with the property that $\psi(Y^{(k)}) \subseteq X^{(k)}$ (so that there are no contractions of points of height k to points of bigger height) induces a site morphism. For $k = 0$ see Example 6.1.10, for $k = 1$ see Section 6.4.

Proposition 6.1.6. *Let a Zariski filter F be given on a topologically noetherian scheme X given rise to the Grothendieck topology X_{zftop} . Let M be a quasicoherent $\mathcal{O}_X$ -module and let $N \subseteq M$ be a submodule. Then the following holds.*

- (i) *The module of global sections is $\Gamma(X_{\text{zftop}}, M_{\text{zftop}}) = \varinjlim_{V \in F} \Gamma(V, M)$.*
- (ii) *The assignment $U \mapsto \Gamma(U_{\text{zftop}}, M_{\text{zftop}})$ for $U \subseteq X$ open is a quasicoherent $\mathcal{O}_X$ -module.*
- (iii) *The sheafification $M \mapsto M_{\text{zftop}}$ is exact.*
- (iv) *The Zariski filter F is an absolute filter in the Grothendieck topology, and the absolute stalk of a module is the same as its module of global sections.*
- (v) *The closure operation induced by X_{zftop} is (on an open affine subset)*

$$N^{\text{zftop}-c} = \{s \in M : \rho_V^X(s) \in \Gamma(V, N) \text{ for some } V \in F\}.$$

If $F = F(X_0)$ for a subset $X_0 \subseteq X$, then

$$N^{\text{zftop}-c} = \{s \in M : s_x \in N_x \text{ for all } x \in X_0\}.$$

- (vi) *If M is coherent, then $M_{\text{zftop}} = 0$ if and only if $M|_V = 0$ for some $V \in F$, and this is true if and only if the support of M does not meet V .*
- (vii) *We have $H^k(X_{\text{zftop}}, M_{\text{zftop}}) = \varinjlim_{V \in F} H^k(V, M)$.*

Proof. (i). The (Zariski-)module $U \mapsto \Gamma(U, M)$ is the associated $\mathcal{O}_{\text{zftop}}$ -premodule in the sense of Proposition 1.5.5(iv), which we have to sheafify. We have

$$\Gamma(U, M_0) = \{s \in \Gamma(U, M) : \rho_{U \cap V}^U(s) = 0 \text{ for some } V \in F\}$$

and the associated separated presheaf is $\Gamma(U, M_1) = \Gamma(U, M)/\Gamma(U, M_0)$ (see Section 1.2). We claim that there are natural homomorphisms

$$\varinjlim_{V \in F} \Gamma(V, M) \longrightarrow \varinjlim_{V \in F} \Gamma(V, M_1) \longrightarrow \Gamma(X_{\text{zftop}}, M_{\text{zftop}}).$$

It is clear that the first mapping exists and that it is a bijection. For the existence of the second map, let $s \in \Gamma(V, M_1)$, $V \in F$. Since $V \times_X V = V \cap V = V$, the section s is compatible by Lemma 1.6.7, so we get a mapping $\Gamma(V, M_1) \rightarrow \Gamma(X_{\text{zftop}}, M_{\text{zftop}})$ and also from the colimit.

Every global section $s \in \Gamma(X_{\text{zftop}}, M_{\text{zftop}})$ is given by a compatible family $s_i \in \Gamma(U_i, M)$, $i \in I$, in a covering $U_i \rightarrow U$, so that the differences are 0 in $\Gamma(U_i \cap U_j, M_1)$. We may assume that I is finite, since U is quasicompact, and suppose that $s_i - s_j \in \Gamma(U_i \cap U_j, M)$ restricts to 0 in $U_i \cap U_j \cap V_{ij}$, $V_{ij} \in F$. Then $W := (\bigcup_{i \in I} U_i) \cap \bigcap_{(i,j) \in I \times I} V_{ij} \in F_U$ and s_i is represented by an element $s \in \Gamma(W, M)$. Hence the second mapping is surjective. It is injective as it is the sheafification homomorphism of a separated presheaf.

(ii). By (i) we have $\Gamma(U_{\text{zftop}}, M_{\text{zftop}}) = \varinjlim_{V \in F} \Gamma(U \cap V, M)$. For fixed V , the assignment $U \mapsto \Gamma(U \cap V, M)$ is a sheaf, as it is the push forward of $M|_V$ under $V \subseteq X$. Therefore also the colimit $U \mapsto \varinjlim_{V \in F} \Gamma(U \cap V, M)$ is a sheaf [27, Exercise II.1.11]. For quasicoherece we may assume that X is affine and that $f \in \Gamma(X, \mathcal{O}_X)$. Then we have $\varinjlim_{V \in F} \Gamma(D(f) \cap V, M) = \varinjlim_{V \in F} \Gamma(V, M)_f = (\varinjlim_{V \in F} \Gamma(V, M))_f$ (see [27, Lemma II.5.14] for the first equality).

(iii). The sheafification is $\mathcal{G} \mapsto \varinjlim_{V \in F} (i_*(\mathcal{G}|_V))$, $(i : V \rightarrow X)$ so the left exactness of the sheafification follows from the left exactness of push forward and from the exactness of taking the colimit. The right exactness follows from Lemma 1.3.6.

(iv). The filter F is a small cofiltered category (Definition 1.2.1) with an arrow $U \rightarrow V$ corresponding to an inclusion $U \subseteq V$. The inclusions $U \rightarrow V$ and $U' \rightarrow V$ come together in the intersection $U \cap U' \rightarrow U, U'$. The identical mapping gives then a filter in the sense of Definition 1.7.1. It is clearly an absolute filter in the sense of Definition 1.9.1 (but not irreducible, as the topology is not single-handed). Then we have $\varinjlim_{V \in F} \Gamma(V_{\text{zftop}}, M_{\text{zftop}}) = \varinjlim_{V \in F} (\varinjlim_{W \in F} \Gamma(V \cap W, M)) = \varinjlim_{V \in F} \Gamma(V, M) = \Gamma(X_{\text{zftop}}, M_{\text{zftop}})$.

(v) follows from Lemma 1.4.5 or directly. For the second part suppose first that $\rho_V^X(s) \in \Gamma(V, N)$ for some $V \in F$. Then $X_0 \subseteq V$ and so $s_x = \rho_x(\rho_V^X(s)) \in N_x$ for all $x \in X_0$. On the other hand, if $s_x \in N_x$, then there exists an open neighborhood of $x \in V_x$ such that $\rho_{V_x}^X(s) \in \Gamma(V_x, N)$. Then $V = \bigcup_{x \in X_0} V_x$ is an open neighborhood of X_0 and $\rho_V^X(s) \in \Gamma(V, N)$, since it is locally true.

(vi). We may reduce to X affine by (ii). Let $s_1, \dots, s_n$ be generators of M . Since $M_{\text{zftop}} = 0$, there exists by (v) for every i an $V_i \in F$ such that $\rho_{V_i}^X(s_i) = 0$ in $\Gamma(V_i, M)$, so all generators become 0 in $\Gamma(V_1 \cap \dots \cap V_n, M)$. On the other hand, if $M|_V = 0$, $V \in F$, then the module is 0 in a covering, hence it is 0.

(vii). The right hand side has all the properties of a universal δ -functor for X_{zftop} [27, Corollary III.1.4]. $\square$

Remark 6.1.7. The statements (i), (iii) and (vii) in Proposition 6.1.6 hold for arbitrary sheaves of abelian groups on X in the sense that $\Gamma(X_{\text{zftop}}, \varphi^{-1}(\mathcal{G})^\#) = \varinjlim_{V \in F} \Gamma(V, \mathcal{G})$.

Example 6.1.8. Let $X = \text{Spec } R$ denote an affine scheme and let T denote a multiplicatively closed subset in R defining the topological filter F_T consisting of open subsets $V \supseteq D(f)$ for some $f \in T$. Then for an R -module M the module of global sections is

$$\Gamma(X_{\text{zftop}}, M_{\text{zftop}}) = \varinjlim_{V \in F_T} \Gamma(V, M) = \varinjlim_{f \in T} M_f = M_T.$$

The closure of $N \subseteq M$ is $N^c = \{s \in M : s \in N_T \text{ in } M_T\} = \{s \in M : \exists f \in T \text{ such that } fs \in N\}$. In the case of such an algebraically defined filter one can also define an affine Grothendieck topology with the same closure operation by declaring $R \rightarrow R_f$ to be a cover if f belongs to the saturated multiplicative system given by T (that is, f divides some element in T).

The affine scheme $\operatorname{Spec} R_T$ gives an absolute cover for this topology. If $X_0 = \{\mathfrak{p} \in \operatorname{Spec} R_T : \mathfrak{p} \cap T = \emptyset\}$, then also $F(X_0) = F_T$. On the other hand, if X_0 is a finite collection of prime ideals $\mathfrak{p}_1, \dots, \mathfrak{p}_n$, then the filter of open neighborhoods $F(X_0)$ is given algebraically by $T = R - \bigcup_{i=1}^n \mathfrak{p}_i$.

Example 6.1.9. Let $X = \operatorname{Spec} R$ be affine and let $T \subseteq R$ be the multiplicative system of all non-zero divisors. The open subsets $\{V \subseteq X : V \supseteq D(f) \text{ for some } f \in T\}$ form a Zariski filter, and we denote the corresponding Grothendieck topology by X_{nzd} . The induced closure operation on submodules is the (non-zero divisor-) *torsion*,

$$N^{\text{nzd}-c} = \operatorname{tors}(N) = \{s \in M : \bar{s} = 0 \text{ in } (M/N)_T\}.$$

The quotient module $M/\operatorname{tors}(0)$ is torsion-free. The module of global sections is then $\Gamma(R_{\text{nzd}}, M_{\text{nzd}}) = M_T$, and $M/\operatorname{tors}(0) \subseteq M_T$ is a torsion-free submodule. M_{nzd} is trivial if and only if M is a torsion module. If R is noetherian, then $T = R - \bigcup_{i=1}^n \mathfrak{p}_i$, where $\mathfrak{p}_i$ runs through the associated primes of R [12, Theorem 3.1]. Therefore $N^{\text{nzd}-c} = \{s \in M : s \in N_{\mathfrak{p}} \text{ for all associated primes } \mathfrak{p}\}$ (Proposition 6.1.6(v)). Here one can also restrict to the maximal members among $\operatorname{Ass}(R)$. For an ideal $I \neq 0$ in a domain R always $I^{\text{nzd}-c} = R$.

For a scheme X the same definitions apply. For an $\mathcal{O}_X$ -submodule $\mathcal{G} \subseteq \mathcal{F}$ the closure $\mathcal{G}^{\text{nzd}-c}$ defined by the property that $\mathcal{F}/\mathcal{G}^{\text{nzd}-c}$ is torsion-free is sometimes called the saturation of $\mathcal{G}$.

Example 6.1.10. For a scheme X let F_{dense} be the topological filter of all open dense subsets V in X . This means for X irreducible that V is non-empty, and in general that V meets every irreducible component. This is the special case $k = 0$ of Example 6.1.5, and we call this topology the *dense topology* and denote it by X_{dense} .

Let $X = \operatorname{Spec} R$ be affine, and let $N \subseteq M$ be R -modules. The closure operation induced by the dense topology is the “dense” torsion of a submodule, in the sense that

$$N^{\text{dense}-c} = \{s \in M : \bar{s} = 0 \text{ in } (M/N)_{\mathfrak{p}} \text{ for all minimal prime ideals } \mathfrak{p}\}.$$

For R noetherian, the module of global sections is $\Gamma(R_{\text{dense}}, M_{\text{dense}}) = \bigoplus_{\operatorname{ht} \mathfrak{p}=0} M_{\mathfrak{p}}$.

6.2. Submodules with support condition.

The easiest topological filters are the ones given by just one open subset $U \subseteq X$, so that $F(U) = \{V : V \supseteq U\}$. The closure operations coming from these filters are directly related to support conditions and to local cohomology. Recall the following definition.

Definition 6.2.1. Let R be a ring, J an ideal and M an R -module. Then

$$\Gamma_J(M) = \{s \in M : J^k s = 0 \text{ for some } k \in \mathbb{N}\}$$

is called the *module of sections in M with support in J* .

If X is a scheme with an open subset $U \subseteq X$ and closed complement $Z = X - U$, and if $\mathcal{F}$ is a sheaf on X , then

$$\Gamma_Z(X, \mathcal{F}) = \{s \in \Gamma(X, \mathcal{F}) : \operatorname{supp}(s) \subseteq Z\}$$

is called the *module of sections with support in Z* [22, Exposé I].

Lemma 6.2.2. Let X be a scheme and let $U \subseteq X$ be an open subset. Let $\mathcal{F}$ be an $\mathcal{O}_X$ -module. Then $\Gamma_{X-U}(X, \mathcal{F})$ is the closure of 0 in $\Gamma(X, \mathcal{F})$ in the Grothendieck topology induced by U (that is, induced by the filter $F(U)$). In particular, if R is a noetherian ring with an ideal J , then $\Gamma_J(M)$ is the closure of 0 in the topology induced by $D(J)$.

Proof. In general, $\text{supp}(s) = \{x \in X : s_x \neq 0\} \subseteq Z$ if and only if $s = 0$ in $\Gamma(U, \mathcal{F})$. So the statement follows from Proposition 6.1.6(v). In the case of a noetherian ring, if $s \in M$ restricts to 0 on $U = D(J)$, then, letting $J = (g_1, \dots, g_m)$, we also have $g_i^{k_i} s = 0$ and so $J^k s = 0$ for $k = k_1 + \dots + k_m$. On the other hand, if $J^k s = 0$, then $s|_U = 0$. $\square$

Recall that $\Gamma_J(-)$ is left exact (this follows also from Proposition 6.1.6(iii) and Lemma 2.4.9) and induces a right derived functor, called local cohomology (see [22], [9]). This is true more generally for a Zariski filter F (but not for an arbitrary closure operation, see Example 6.2.4). In this case we also write

$$\Gamma_F(M) = \{s \in \Gamma(X, M) : s|_V = 0 \text{ for some } V \in F\} = \varinjlim_{V \in F} \Gamma_{X-V}(M).$$

The long exact cohomology sequences [22, Corollaire I.2.9] for $V \in F$ induce a long exact cohomology sequence in the colimit, namely

$$\begin{aligned} 0 \longrightarrow \Gamma_F(M) \longrightarrow \Gamma(X, M) \longrightarrow \varinjlim_{V \in F} H^0(V, M) = H^0(X_{\text{zftop}}, M_{\text{zftop}}) \longrightarrow \\ H_F^1(M) \longrightarrow H^1(X, M) \longrightarrow \varinjlim_{V \in F} H^1(V, M) = H^1(X_{\text{zftop}}, M_{\text{zftop}}) \longrightarrow . \end{aligned}$$

Example 6.2.3. For a commutative ring R with an ideal J the closure of an R -submodule $N \subseteq M$ is given as $N^c = \{s \in M : J^k s \subseteq N \text{ for some } k\}$. If $R = K[x_1, \dots, x_d]$ is a polynomial ring, and $J = \mathfrak{m} = (x_1, \dots, x_d)$, then for an ideal $I \subseteq R$ this closure yields $I^c = \{f \in R : \mathfrak{m}^k f \subseteq I \text{ for some } k\}$, which is also called the *saturation* of the ideal. A homogeneous ideal I and its saturation I^{sat} define the same projective variety $\text{Proj } R/I = \text{Proj } R/I^{\text{sat}}$.

Example 6.2.4. The closure of 0 is not left exact in general. Let K be a field and let $R = K[\varepsilon]/(\varepsilon^2)$, and suppose we are in the surjective Grothendieck topology (Part 3). Then the short exact sequence of R -modules $0 \rightarrow K \cong (\varepsilon) \rightarrow R \rightarrow R/(\varepsilon) = K \rightarrow 0$ yields the non-exact sequence of closures of 0 (radicals), $0 \rightarrow 0 \rightarrow (\varepsilon) \rightarrow 0 \rightarrow 0$. So the left-exactness of $\Gamma_J(-)$ is a particularity of the functor “sections with support in”.

6.3. Torsion theories, Zariski filters and Grothendieck topologies.

In this section we relate Grothendieck topologies induced by a Zariski filter to hereditary torsion theories and show that they are essentially the same. Torsion theories for abelian categories have gained a lot of attention, in particular for the category of left-modules for non-commutative rings. We recall briefly the basics of torsion theory until we are able to translate them into our setting. Introductions to torsion theories can be found in [3], [11], [16] or [65, Chapter 3].

A *torsion theory* on an abelian category $\mathcal{C}$ (think of the category of R -modules) consists in two classes $\mathcal{T}$ and $\mathcal{F}$ of objects (the “torsion class” and the “torsion-free class”) of $\mathcal{C}$ such that $\mathcal{T} \cap \mathcal{F} = \{0\}$, $\mathcal{T}$ is closed under quotients, $\mathcal{F}$ is closed under subobjects, and for every object $M \in \mathcal{C}$ there exists a short exact sequence $0 \rightarrow T \rightarrow M \rightarrow F \rightarrow 0$ with $T \in \mathcal{T}$ and $F \in \mathcal{F}$, see [11]. There exists then also such a short exact sequence where T is maximal ([11, Proposition 2.4] or [65, Proposition 3.1]), called the *torsion submodule* of M .

We restrict to the case of R -modules and we denote the torsion submodule of M by T_M . The class $\mathcal{T}$ is characterized by the property $T_M = M$ [65, Proposition 3.2]. A torsion

theory is immediately a closure operation for the 0-submodule and it becomes a closure operation for submodules $N \subseteq M$ by pulling back the torsion submodule from M/N [65, Chapter 3.4]. A torsion theory is called *hereditary* if $\mathcal{T}$ is also closed under subobjects.

Theorem 6.3.1. *Let R be a noetherian commutative ring. Then the following structures are equivalent.*

- (i) *A closure operation on $R\text{-Mod}$ coming from a Grothendieck topology on $\text{Spec } R$ (i.e. an admissible closure operation) such that the functor $M \mapsto 0_M^c$ is left exact.*
- (ii) *A hereditary torsion theory on $R\text{-Mod}$.*
- (iii) *A Gabriel filter GF of ideals. This is a set of ideals fulling (a) if $J \in GF$ and $g \in R$, then $(J : g) \in GF$ and (b) if $J \in GF$ and if $(I : g) \in GF$ for all $g \in J$, then also $I \in GF$ [65, Definition 3.4].*
- (iv) *A Zariski filter of open subsets in $\text{Spec } R$.*

Proof. We describe the constructions leading from one structure to another. From (i) to (ii). One declares $\mathcal{T}$ to be the class of R -modules M with $0^c = M$ and $\mathcal{F}$ to be the R -modules with $0^c = 0$. We go through the axioms of a torsion theory. $\mathcal{T} \cap \mathcal{F} = 0$ is clear. Let $\varphi : T \rightarrow S \rightarrow 0$ be a surjection and suppose that T is $\mathcal{T}$ -torsion. Then $0_S^c \supseteq \varphi(0_T^c) = \varphi(T) = S$. If $0 \rightarrow G \rightarrow F$ is an injection and $0_F^c = 0$, then by the module persistence (Proposition 2.2.6(i)) of the closure operation also $0_G^c = 0$, so $G \in \mathcal{F}$.

So far we have not used the left-exactness. For M we consider the short exact sequence $0 \rightarrow 0^c \rightarrow M \xrightarrow{\varphi} M/0^c = F \rightarrow 0$. By left exactness we have also the intersection property $N \cap 0_M^c = 0_N^c$ for $N \subseteq M$. For if $t \in N$ maps to 0_M^c , then it maps to 0 in $0_{M/N}^c$ and by left exactness it must belong to 0_N^c . From this intersection property it follows that $0_{0_M^c}^c = 0_M^c$, so this module belongs to $\mathcal{T}$. By the independence of presentation (Proposition 2.2.6(ii)) we have $\varphi^{-1}(0_F^c) = (\varphi^{-1}(0))^c = (0^c)^c = 0^c$, hence $0_F^c = 0$ and $F \in \mathcal{F}$. By the intersection property it also follows for $0 \rightarrow S \rightarrow T$ and $T \in \mathcal{T}$ that $S \in \mathcal{T}$, so $(\mathcal{T}, \mathcal{F})$ is a hereditary torsion theory.

The equivalence of (ii) and (iii) is a well known fact from torsion theory (see [65, Proposition 3.6]). In a hereditary torsion theory for R -modules one has $T_M = \{s \in M : R/\text{Ann}(s) \in \mathcal{T}\}$, hence a hereditary torsion theory is characterized by the set of ideals $J \subseteq R$ such that R/J belongs to $\mathcal{T}$. These ideals form the *Gabriel filter* $GF = GF(\mathcal{T}, \mathcal{F})$ of the torsion theory. A Gabriel filter GF defines a hereditary torsion theory by setting $T_M = \{s \in M : \text{Ann}(s) \in GF\}$.

From (iii) to (iv). A Gabriel filter is closed under bigger ideals and contains with J and I also JI . In particular, it contains arbitrary powers of I . Therefore, in the case of a commutative noetherian ring, a Gabriel filter contains with an ideal also all ideals with the same radical. Hence it corresponds to a Zariski filter.

From (iv) to (i) follows from Proposition 6.1.6(iii) and Lemma 2.4.9.

We have to show that the constructions are inverse to each other. A left exact closure operation c yields the Zariski filter $D(J)$, $J^c = R$ and the corresponding Grothendieck topology, and we have to show that $c = \text{zftop} - c$. We can restrict to the closure of 0. If $s \in 0^c$, then $s \in T_M$, so $\text{Ann}(s) \in GF$ and so $D(\text{Ann}(s)) \in \mathcal{F}$. Since $s|_{D(\text{Ann}(s))} = 0$, we have $s \in 0^{\text{zftop}-c}$. If $s \in 0^{\text{zftop}-c}$, then $s|_{D(J)}$ for some ideal J with $J^c = R$. Then $\text{supp}(s) \cap D(J) = \emptyset$ and so $D(J) \subseteq D(\text{Ann}(s))$, hence $\text{Ann}(s) \in GF$. Therefore $\text{Ann}(s)^c = R$ and $R/\text{Ann}(s)$ is a torsion module. We have $R/\text{Ann}(s) \rightarrow M$, $1 \mapsto s$, and $1 \in 0^c$ implies by persistence that $s \in 0^c$. The other directions are similar. $\square$

Example 6.3.2. We give an example of a non-hereditary torsion theory [65, Example 3.1(8)]. Consider the torsion theory for R -modules which is cogenerated by R . This means that an R -module M is torsion if and only if $\text{Hom}_R(M, R) = 0$. If R is a domain, but not a field, then $\text{Hom}_R(Q(R), R) = 0$, so the quotient field is torsion, but the submodule $R \subset Q(R)$ is not-torsion, so the hereditary property does not hold. This torsion theory does not fulfill the first finiteness condition 2.2.14, so it can not be induced by a quasicompact Grothendieck topology by Corollary 2.4.7.

6.4. The divisorial topology, reflexive hull and symbolic powers.

In this section we deal with the following special case of Example 6.1.5.

Definition 6.4.1. Let X be a scheme and let $X^{(1)}$ be its points of height at most one. The Grothendieck topology on X , where the open sets are the Zariski open sets and where $U_i \rightarrow U$, $i \in I$, is a covering if $U^{(1)} \subseteq \bigcup_{i \in I} U_i$, is called the *divisorial topology*, denoted by X_{div} .

We will relate the divisorial topology with symbolic powers of ideals and with the reflexive hull of a module. Recall that the dual of an R -module is $M^\vee = \text{Hom}_R(M, R)$ and that the *bidual* of a module M is $M^{\vee\vee}$. We denote the natural homomorphism $M \rightarrow M^{\vee\vee}$ given by $s \mapsto (\varphi \mapsto \varphi(s))$ by θ . M is called *torsionless* if θ is injective and *reflexive* if θ is bijective; see [10, 1.4] for these definitions. For a domain R a finite torsion-free (with respect to non-zero divisors, see Example 6.1.9) module M is also torsionless.

Lemma 6.4.2. *Let R be a noetherian normal domain and let M be a finite torsion-free R -module. Then $\Gamma(R_{\text{div}}, M_{\text{div}}) = M^{\vee\vee}$, the reflexive hull of M .*

Proof. We have a natural R -module homomorphism

$$M \longrightarrow \varinjlim_{X^{(1)} \subseteq V} \Gamma(V, M) = \Gamma(R_{\text{div}}, M_{\text{div}}).$$

We first claim that the colimit on the right is reflexive, using [10, Proposition 1.4.1(b)]. For prime ideals $\mathfrak{p}$ with $\text{depth } R_{\mathfrak{p}} \leq 1$ also the height is ≤ 1 (as R is normal) and so $R_{\mathfrak{p}}$ is a field or a discrete valuation domain. Hence $M_{\mathfrak{p}}$ is free, since it is torsion-free, and so it is reflexive.

Let now $\mathfrak{p}$ be a prime ideal with $\text{depth } R_{\mathfrak{p}} \geq 2$ and let $t_1, t_2 \in R$ be a regular sequence in $R_{\mathfrak{p}}$. We have to show that they are a regular sequence for M as well. Let s_1, s_2 be two sections of the colimit and suppose that $t_1 s_1 + t_2 s_2 = 0$. We may assume that $s_1, s_2 \in \Gamma(V, M)$, that $V \subseteq D(t_1, t_2)$ and that the equation holds over V . Then $\frac{s_1}{t_2} = -\frac{s_2}{t_1} =: s$ is a global element of $\Gamma(V, M)$ and hence $s_1 = t_2 s$.

The reflexivity implies that we get a homomorphism $M^{\vee\vee} \rightarrow \Gamma(R_{\text{div}}, M_{\text{div}})$. There exists an open subset $W \supseteq X^{(1)}$ where $M|_W$ is locally free, since $M_{\mathfrak{p}}$ is free for every prime ideal of height one. Hence the colimit equals $\Gamma(W, M)$ and it is a finite R -module. A homomorphism between two finite reflexive R -modules is by [61, Lemma 24.2] a bijection if this is true after localization at every prime ideal of height one. For such a prime ideal we have $(\varinjlim_{X^{(1)} \subseteq V} \Gamma(V, M))_{\mathfrak{p}} = \varinjlim_{\mathfrak{p} \in V} \Gamma(V, M) = M_{\mathfrak{p}} = (M^{\vee\vee})_{\mathfrak{p}}$. $\square$

We come to the closure operation induced by the divisorial topology.

Proposition 6.4.3. *Let R be a normal noetherian domain. The closure operation for finite module $N \subseteq M$ induced by the divisorial topology is the ('internal') reflexive hull, namely*

$$N^{\text{div}-c} = \{s \in M : \theta(s) \in N^{\vee\vee} \text{ (inside } M^{\vee\vee})\}.$$

For submodules of a finite reflexive module, the closure is the reflexive hull $N^{\vee\vee}$. For ideals $I \subseteq R$ we have

$$\begin{aligned} I^{\text{div}-c} &= R \cap \bigcap_{I \subseteq \mathfrak{p} \text{ prime of height one}} IR_{\mathfrak{p}} \\ &= \{f \in R : \nu_{\mathfrak{p}}(f) \geq \min\{\nu_{\mathfrak{p}}(g) : g \in I\} \text{ for all } \mathfrak{p} \text{ of height one}\}, \end{aligned}$$

where $\nu_{\mathfrak{p}}$ denotes the corresponding discrete valuation. In particular, for a prime ideal $\mathfrak{p}$ of height one we have $(\mathfrak{p}^n)^{\text{div}-c} = \mathfrak{p}^{(n)}$, the n th symbolic power of $\mathfrak{p}$. For ideals $I \subseteq R$ of height ≥ 2 we have $I^{\text{div}-c} = R$.

Proof. The inclusion $N \subseteq M$ yields by left exactness (Proposition 6.1.6(iii)) an inclusion $N_{\text{div}} \subseteq M_{\text{div}}$ of sheaves in the divisorial topology. The modules of global sections are by Lemma 6.4.2 $\Gamma(R_{\text{div}}, N_{\text{div}}) = N^{\vee\vee} \subseteq \Gamma(R_{\text{div}}, M_{\text{div}}) = M^{\vee\vee}$. For M reflexive we have the inclusions $N \subseteq N^{\vee\vee} \subseteq M^{\vee\vee} = M$.

By Proposition 6.1.6(v) we have $I^{\text{div}-c} = R \cap \bigcap_{I \subseteq \mathfrak{p} \text{ prime of height one}} IR_{\mathfrak{p}}$, and we can skip the prime ideals which do not contain I . The extended ideal $IR_{\mathfrak{p}}$ for a prime ideal $\mathfrak{p}$ of height one equals $(\mathfrak{p}R_{\mathfrak{p}})^{\nu}$ for $\nu = \min\{\nu_{\mathfrak{p}}(g) : g \in I\}$, which gives the second equality. Also, $\mathfrak{p}^n \subseteq \mathfrak{p}$ is the only minimal prime ideal, and $\mathfrak{p}^{(n)} = R \cap \mathfrak{p}^n R_{\mathfrak{p}}$ is by definition the n -th symbolic power of $\mathfrak{p}$ [12, Section 3.9]. Finally, if I has height ≥ 2 , meaning that every minimal prime ideal of I has height ≥ 2 , then the index set is empty. $\square$

Proposition 6.4.4. *Let X be a normal noetherian integral scheme endowed with the divisorial topology, let $Q = Q(X)$ be its function field. Then the following holds.*

- (i) *The coherent submodules of Q in the divisorial topology are all invertible and are in one to one correspondence with Weil divisors. The coherent ideal sheaves correspond to effective Weil divisors.*
- (ii) *We have $H^1(X_{\text{div}}, \mathcal{O}_{\text{div}}^{\times}) = \text{Cl } X$, the divisor class group of X (compare [22, Proposition XI.3.7.1]).*

Proof. (i). By coherent we mean here just coherent as a module in the Zariski topology. Let $\mathcal{L} \subseteq Q$ be a coherent submodule. For $\mathfrak{p} \in X$ of height one the localization $\mathcal{L}_{\mathfrak{p}} \subseteq Q$ is free of rank one, and by coherence there exists an open affine neighborhood $\mathfrak{p} \in U$ such that $\mathcal{L}|_U$ is free (in the Zariski topology and in the divisorial topology). These neighborhoods do cover X in the divisorial topology (not necessarily in the Zariski topology), hence $\mathcal{L}$ is invertible. In general, a Weil divisor corresponds to an invertible submodule of Q on some open subset $V \supseteq X^{(1)}$. So the result follows.

(ii). The exact sequence $0 \rightarrow \mathcal{O}^{\times} \rightarrow Q^{\times} \rightarrow Q^{\times}/\mathcal{O}^{\times} \rightarrow 0$ of sheaves of abelian groups yields the corresponding exact sequences on X_{div} (Proposition 6.1.6(iii) and Remark 6.1.7) and the global sequence

$$0 \rightarrow \Gamma(X, \mathcal{O}^{\times}) \rightarrow Q^{\times} \rightarrow \varinjlim_{X^{(1)} \subseteq V} \Gamma(V, Q^{\times}/\mathcal{O}^{\times}) \rightarrow \varinjlim_{X^{(1)} \subseteq V} H^1(V, \mathcal{O}^{\times}) = H^1(X_{\text{div}}, \mathcal{O}_{\text{div}}^{\times}) \rightarrow 0.$$

A section $s \in \Gamma(V, Q^{\times}/\mathcal{O}^{\times})$ is given by a covering $V = \bigcup_{i \in I} V_i$ and $q_i \in Q^{\times}$ with $q_i q_j^{-1} \in \Gamma(V_i \cap V_j, \mathcal{O}^{\times})$. It defines clearly a submodule which is invertible in codimension one. On

the other hand, a Weil divisor corresponds to such a submodule $\mathcal{L} \subseteq Q$, and $\mathcal{L}$ is for every $x \in X^{(1)}$ generated by $q_x \in Q$, and this is true in a neighborhood of x . The union of these neighborhoods together is then an open set $V \supseteq X^{(1)}$. This gives an identification $\text{Div} X = \varinjlim_{X^{(1)} \subseteq V} \Gamma(V, Q^\times / \mathcal{O}^\times)$. So the written down cohomology sequences is just the defining sequence of the divisor class group [27, Section II.6]. $\square$

7. FURTHER EXAMPLES

7.1. Completion.

Also the completion of a ring with respect to an ideal may be considered as given a Grothendieck topology, but quite in a different flavor. In particular the coverings there are not coverings in the set-theoretical sense, and it is not a single-handed Grothendieck topology nor a quasicompact one.

Let R denote a commutative ring and $\mathfrak{a} \subseteq R$ an ideal. We consider the category over $X = \text{Spec } R$ consisting of X and of all $X_k := \text{Spec } R/\mathfrak{a}^k$, $k \geq 0$, and all X -morphisms. The coverings in this topology are given by families which contain the identity ($X_k \rightarrow X_k$ or $X \rightarrow X$) and by families which contain $X_k \rightarrow X$, $k \in I \subseteq \mathbb{N}$, where I is infinite. We denote this Grothendieck topology by $X_{\mathfrak{a}\text{-top}}$ (similar definitions are possible for ideal families $\mathfrak{a}_k$, $k \in \mathbb{N}$, with $\mathfrak{a}_0 = R$ and $\mathfrak{a}_k \cdot \mathfrak{a}_\ell \subseteq \mathfrak{a}_{k+\ell}$; one can also formulate scheme versions). An R -module M defines the presheaf M^{pre} by

$$\Gamma(X_k, M^{\text{pre}}) := M \otimes_R R/\mathfrak{a}^k = M/\mathfrak{a}^k M.$$

Proposition 7.1.1. *Let $X = \text{Spec } R$ be an affine scheme endowed with the topology $X_{\mathfrak{a}\text{-top}}$ for an ideal $\mathfrak{a} \subseteq R$. Let M denote an R -module. Then the following holds.*

- (i) *The presheaf M^{pre} is a sheaf with module of global sections*

$$\Gamma(X_{\mathfrak{a}\text{-top}}, M^{\text{pre}}) = \hat{M} = \varprojlim_{k \in \mathbb{N}} M/\mathfrak{a}^k M,$$

the $\mathfrak{a}$ -adic completion of M . In particular, the ring of global sections in this topology is just the completion $\hat{R}$ of R with respect to the ideal $\mathfrak{a}$.

- (ii) *The global sheafification homomorphism $\theta : R \rightarrow \Gamma(R_{\mathfrak{a}\text{-top}}, \mathcal{O}) = \hat{R}$ has $\mathfrak{b} = \bigcap_{k \in \mathbb{N}} \mathfrak{a}^k$ as its kernel, which is zero for $\mathfrak{a} \neq R$ in a noetherian ring.*
- (iii) *The mapping $\overline{\mathbb{N}} = \mathbb{N} \cup \{\infty\} \rightarrow \mathcal{T}_X$, $k \mapsto X_k$ ($X_\infty = X$), is a non-absolute irreducible filter. The stalk of a module is 0.*
- (iv) *The mapping $\{P\} \rightarrow \mathcal{T}_X$, $P \mapsto X$, is an absolute filter which is not irreducible. Its stalk is the same as the module of global sections.*
- (v) *The closure operation induced by this topology is just $I \rightarrow I^{\text{top-cl}} = I + \mathfrak{b}$, so it is the identical operation in the noetherian case.*
- (vi) *A ring homomorphism $\psi : R \rightarrow R'$ where $\psi(\mathfrak{a}) \subseteq \mathfrak{a}'$ induces a site morphism $X'_{\mathfrak{a}'\text{-top}} \rightarrow X_{\mathfrak{a}\text{-top}}$ and the corresponding homomorphism of rings of global sections is just $\hat{R} \rightarrow \hat{R}'$.*

Proof. (i) An element in $\hat{M}$ is given by a sequence $s_k \in M/\mathfrak{a}^k M$ with the compatibility condition $\varphi_{k\ell}(s_k) = s_\ell$ for all $k \geq \ell$. This is exactly the compatibility condition in the topology. (ii) is clear.

(iii) For $k \leq \ell$ we have $\mathfrak{a}^k \supseteq \mathfrak{a}^\ell$, hence $R/\mathfrak{a}^\ell \rightarrow R/\mathfrak{a}^k$ and so $X_k \rightarrow X_\ell$. As $X_k \rightarrow X$ are not covers, this filter is not absolute. Since $\emptyset = \text{Spec } R/\mathfrak{a}^0$ is the initial object of the filter, all stalks are 0.

(iv) The only single cover of X is the identity, so the filter is clearly absolute. The covering $X_k \rightarrow X$, $k \in \mathbb{N}$, is such that the identity $X \rightarrow X$ factors through no X_k , hence this filter is not irreducible (Definition 1.8.1). (v) and (vi) are clear. $\square$

7.2. Proper topology.

Proper topologies seem to have first been considered by S.-I. Kimura [42, Definition 3.1] in order to give a cohomological characterization of Alexander schemes. Here we treat the proper topology in its relation to the integral closure of ideals.

Definition 7.2.1. Let X be a scheme. Let $\mathcal{T}_X$ denote the category of all schemes over X and define a covering to be a proper surjective morphism $V \rightarrow U$ (we do not insist that the structure morphisms are proper). We denote X endowed with this Grothendieck topology by X_{prop} .

Remark 7.2.2. Recall that a proper morphism is separated, of finite type and universally closed [27, Section II.4]. Proper morphisms are stable under base change and under composition, hence we get indeed a Grothendieck topology. Note that the proper topology is not an affine topology and that it is not a refinement of the (single-handed) Zariski topology. Every scheme morphism induces a site morphism in the proper topology. The proper topology is a refinement of the finite topology. A proper and surjective homomorphism is universally submersive.

Important examples of proper morphisms are given by blow-ups of ideals, see [18, Définition 8.1.3] or [27, Propositions II.7.10, II.7.13]. Recall that the blow-up of an ideal $H \subseteq R$ is given by $X_H = \text{Proj } B_H$, $B_H = \bigoplus_{k \in \mathbb{N}} H^k$. We characterize first which blow-ups yield surjective morphisms.

Lemma 7.2.3. *A blow-up $X_H \rightarrow X = \text{Spec } R$ of an ideal $H \subseteq R$ in a noetherian ring is surjective if and only if H is not contained in any minimal prime ideal of R .*

Proof. Let $X_i = \text{Spec } R_i$ denote the integral components of $\text{Spec } R$. If H is not contained in the minimal prime $\mathfrak{q}$, then the extended ideal in $R/\mathfrak{q}$ is not the 0-ideal. For a domain the blow-up of a non-zero ideal defines a surjective morphism, since the image is closed and the morphism is an isomorphism on a non-empty open set [27, Proposition II.7.13]. Since the blow-up of the component factors through $X_H \times_X X_i$ [27, Corollary II.7.15], it follows that the morphism is surjective on every component, hence surjective.

On the other hand, if $HR_i = 0$, then over the open set $U_i = X_i - \bigcup_{j \neq i} X_j$ we have $X_H \times_X U_i = (U_i)_{H|U_i} = \emptyset$. $\square$

The *extended ideal* of $I \subseteq R$ under a scheme morphism $Y \rightarrow \text{Spec } R$ is on every affine subset $V = \text{Spec } S \subseteq Y$ given by IS . This gives an ideal sheaf on Y . The following lemma is motivated by Ratliff's Δ -closure, see Section 7.3.

Lemma 7.2.4. *Let R denote a noetherian ring and suppose that $fH \subseteq IH$, where $f \in R$ and where I and H are ideals. Then f belongs to the extended ideal of I under the blow-up $X_H \rightarrow X$.*

Proof. Set $I = (f_1, \dots, f_n)$ and $H = (h_1, \dots, h_m)$. The condition $fH \subseteq IH$ means explicitly that for every h_i we may write $fh_i = \sum_{1 \leq r \leq n, 1 \leq s \leq m} a_{rsi} f_r h_s$, $a_{rsi} \in R$. This translates to

$$f = \sum_{r=1}^n \left(\sum_{s=1}^m \frac{a_{rsi} h_s}{h_i} \right) f_r \quad \text{in } (B_H)_{h_i},$$

so $f \in I((B_H)_{h_i})_0$ for $i = 1, \dots, m$. This means that f belongs to the extended ideal of I on $D_+(h_i) \subseteq X_H$. These open sets cover X_H , therefore f belongs also globally to the extended ideal on X_H . $\square$

Proposition 7.2.5. *Let $X = \operatorname{Spec} R$ denote an affine scheme endowed with the proper topology. Then the closure induced by this topology is the integral closure.*

Proof. Assume first that f belongs to the extended ideal $I\mathcal{O}_Y$ for a proper surjective morphism $Y \rightarrow X$. Such a morphism is universally submersive [4]. Under the base change $\gamma : \operatorname{Spec} V \rightarrow X$, where V is a discrete valuation domain, the image $\gamma^*(f)$ is also in the extended ideal of $I\mathcal{O}_V$ in Y_V . Thus we may assume that $R = V$ is a discrete valuation domain with local parameter π , that $f = \pi^n$ and $I = (\pi^m)$. Then by [4] it follows that there exists a discrete valuation domain W such that $\operatorname{Spec} W \rightarrow Y \rightarrow V$ is surjective, which shows that $f \in \bar{I} = I$.

Suppose now that $f \in \bar{I}$. Let $X_j = \operatorname{Spec} R_j$, $j \in J$, denote the integral components of X . The morphism $\bigsqcup X_j \rightarrow X$ is proper and surjective. We will construct $Y \rightarrow X$ proper and surjective on each component X_j separately according to whether the extended ideal $I\mathcal{O}_{X_j}$ is 0 or not. In the first case we also have $f_j = 0$, and we do not need to modify X_j .

So suppose that I extends to a non-zero ideal in R_j , or that R is a domain and $I \neq 0$. We use the theory of reductions, see [38, Chapter 5]. If $f \in \bar{I}$, then I is a reduction of $L = (I, f)$, that is, $L^{r+1} = IL^r$ for some r . So in particular $fL^r \subseteq IL^r$. Hence f belongs to the extended ideal of the blow-up along the ideal L^r by Lemma 7.2.4. Since $L \neq 0$, this blow-up is a surjection. $\square$

Remark 7.2.6. This result is related to the known fact that f belongs to the integral closure of an ideal I in a domain if and only if it belongs to the extended ideal in the normalization of the blow-up of I , see [13, Proposition 1.4].

7.3. The Δ -closure of Ratliff.

In [55], Ratliff introduced the concept of the Δ -closure of an ideal (see also [57] and [58]). Here Δ is a fixed multiplicatively closed set of ideals in a commutative ring R and the closure operation for an ideal $I \subseteq R$ is given by

$$I_\Delta = \bigcup_{H \in \Delta} (IH : H).$$

Thus an element $f \in R$ belongs to I_Δ if and only if there exists an ideal $H \in \Delta$ such that $fH \subseteq IH$. Under mild conditions the Δ -closure lies inside the integral closure (see [55, Theorem 3.2] or Proposition 7.3.6 below).

We will translate this closure operation into our language and we will show how some of Ratliff's basic results follow immediately. It will turn out that the most natural Grothendieck topology to induce the Δ -closure is given by certain non-affine proper mappings. The basic relationship is given by the following proposition (compare with [29, Fact 2.1] for the case of the Ratliff-Rush closure).

Proposition 7.3.1. *Let R denote a commutative noetherian ring, let Δ be a multiplicatively closed collection of ideals in R . Let $X_H = \operatorname{Proj} B_H, B_H = \bigoplus_{k \in \mathbb{N}} H^k$, $H \in \Delta$, denote the corresponding blow-ups. Then an element $f \in R$ belongs to Ratliff's Δ -closure of an ideal $I \subseteq R$, i.e. $f \in I_\Delta$, if and only if f belongs for some $X_H \rightarrow X = \operatorname{Spec} R$, $H \in \Delta$, to the extended ideal of I .*

Proof. Set $I = (f_1, \dots, f_n)$. Suppose first that $f \in I_\Delta$, so that there exists an ideal $H \in \Delta$ with $fH \subseteq IH$. Then f belongs to the extended ideal under the blow-up $X_H \rightarrow X$ by Lemma 7.2.4.

For the other direction suppose that f belongs to the extended ideal of I under the blow-up $X_H \rightarrow X$, $H = (h_1, \dots, h_m)$. This means that we have locally $f \in I$ on the affine covering sets $D_+(h_i)$, that is, $f \in I((B_H)_{h_i})_0$. This means explicitly that $f = \sum_{j=1}^n \frac{b_{ij}}{h_i^{t_{ij}}} f_j$, where $b_{ij} \in H^{t_{ij}}$. Hence there exists also a t such that $fh_i^t \in IH^t$ for all i . It follows that $fH^{mt} \subseteq IH^{mt}$, since for a monomial $h_1^{\nu_1} \dots h_m^{\nu_m} \in H^{mt}$ we have $\nu_i \geq t$ for some i and hence

$$fh_1^{\nu_1} \dots h_m^{\nu_m} = fh_i^t h_1^{\nu_1} \dots h_i^{\nu_i - t} \dots h_m^{\nu_m} \in IH^t \cdot H^{mt-t} \subseteq IH^{mt}.$$

Since Δ is multiplicatively closed we have $H^{mt} \in \Delta$ and we get $f \in I_\Delta$. $\square$

Definition 7.3.2. Let X denote a scheme and let Δ denote a multiplicatively closed set of ideal sheaves. For $\psi : Y \rightarrow X$ denote the set of extended ideal sheaves $H\mathcal{O}_Y$, $H \in \Delta$, by Δ_Y . Then the Δ -topology on X is given by the full category of schemes of finite type over X , and we declare $Z \rightarrow Y$ to be a cover if there exists a factorization $Y_H \rightarrow Z \rightarrow Y$, where $H \in \Delta_Y$ and where $Y_H \rightarrow Y$ is the blow-up of $H\mathcal{O}_Y$.

Remark 7.3.3. If $Y_H \rightarrow Z \rightarrow Y$ is a covering and $U \rightarrow Y$ an X -morphism, then we get a factorization $U_H \rightarrow Y_H \times_Y U \rightarrow Z \times_Y U \rightarrow U$ by the universal property of blow-up [27, Proposition 7.14]. Hence the pull-back of a covering is a covering. If $Z \rightarrow Y$ and $W \rightarrow Z$ are coverings with factorizations $Y_H \rightarrow Z \rightarrow Y$ and $Z_G \rightarrow W \rightarrow Z$, then we get $Y_{HG} \cong (Y_H)_G \rightarrow Z_G$, which shows that the composition of coverings is a covering. So we have indeed a Grothendieck topology.

Corollary 7.3.4. Let R denote a noetherian commutative ring and let Δ denote a multiplicatively closed set of ideals. Then the Δ -closure equals the closure induced by the Δ -topology on $X = \text{Spec } R$.

Proof. This follows immediately from Lemma 2.4.4, Proposition 7.3.1 and Definition 7.3.2. $\square$

Proposition 7.3.5. Let $\varphi : Y_{\Delta'} \rightarrow X_\Delta$ be a scheme morphism such that $H\mathcal{O}_Y \in \Delta'$ for all ideal sheaves $H \in \Delta$. Then φ induces a site morphism in the Δ -topology. In particular, the persistence property holds for the Δ -closure for these morphisms (as stated in [55, Theorem 5.1]).

Proof. If $V \rightarrow X$ is a covering with factorization $X_H \rightarrow V \rightarrow X$, then the factorization $Y_H \rightarrow X_H \times_X V \rightarrow U \times_X V \rightarrow V$ shows that the pull-back is also a covering. The second statement follows from Theorem 2.4.8. $\square$

Proposition 7.3.6. Let R denote a noetherian ring and let Δ denote the set of all ideals which are not contained in any minimal prime ideal. Then for every ideal $I \in \Delta$ we have $I_\Delta = \overline{I}$, the integral closure of I .

Proof. Let $f \in I_\Delta$, and suppose that f belongs to the extended ideal of I under $X_H \rightarrow X$, $H \in \Delta$. This blow-up is surjective by Lemma 7.2.3, hence f belongs to the proper closure of I and hence to the integral closure by Proposition 7.2.5.

On the other hand, if $f \in \overline{I}$, then f belongs to the extended ideal of I under the blow-up of a power of (I, f) as in the proof of Proposition 7.2.5. $\square$

Ratliff also associates to a commutative ring endowed with Δ another ring R^Δ , the Δ -closure of the ring [55, Definition 6.2]. In the case where Δ does not contain the zero-ideal and R is noetherian this is (see [55, Theorem 6.2])

$$R^\Delta = \{c/b : b \text{ non-zero divisor and } c \in (b)_\Delta\} \subseteq Q(R),$$

where $Q(R)$ denotes the total quotient ring. We relate this construction to absolute stalks and rings of global sections in the Δ -topology.

We endow a given multiplicatively closed set Δ of ideals with the structure of a cofiltered set by declaring that an arrow $H \rightarrow G$ exists if G extends to an invertible ideal in X_H . In this case there is a unique morphism $X_H \rightarrow X_G$ by [27, Proposition II.7.14].

Lemma 7.3.7. *Let R denote a commutative ring, and let Δ denote a multiplicatively closed set of ideals. Then Δ is a cofiltered set and $\Delta \rightarrow \mathcal{T}_X$, $H \mapsto X_H$, is an absolute filter. If $R \rightarrow S$ is a ring homomorphism such that for every $H \in \Delta$ the extended ideal HS is invertible, then there exists a ring homomorphism $\varinjlim_{H \in \Delta} \Gamma(X_H, \mathcal{O}) \rightarrow S$.*

Proof. Two arrows $H \rightarrow E$, $G \rightarrow E$ have the common refinement HG , because in X_{HG} both H and G become invertible. Since there is at most one arrow, Δ is cofiltered in the sense of Definition 1.2.1.

By the definition of a cover in the Δ -topology every morphism $X_H \rightarrow X$ indexed by this filter is a cover. If $Y \rightarrow X_H$ is a cover, then there exists an ideal $G \in \Delta$ and a factorization $(X_H)_G \rightarrow Y \rightarrow X_H$, and so this filter is absolute (Definition 1.9.1). The corresponding absolute stalk of the structure sheaf is hence $\varinjlim_{H \in \Delta} \Gamma(X_H, \mathcal{O})$.

If $R \rightarrow S$ is given and the extended ideal HS is invertible, then there exists a unique morphism $\text{Spec } S \rightarrow X_H$ and hence $\Gamma(X_H, \mathcal{O}) \rightarrow S$, which extends to a ring homomorphism $\varinjlim_{H \in \Delta} \Gamma(X_H, \mathcal{O}) \rightarrow S$. $\square$

Proposition 7.3.8. *Let $X = \text{Spec } R$ denote an affine scheme and let Δ denote a multiplicatively closed set of ideals in R such that every $H \in \Delta$ contains a non-zero divisor. Then we have identities $\Gamma(X_\Delta, \mathcal{O}_\Delta) = \varinjlim_{H \in \Delta} \Gamma(X_H, \mathcal{O}) = R^\Delta$.*

Proof. We have natural homomorphisms $\Gamma(X_\Delta, \mathcal{O}_\Delta) \rightarrow \varinjlim_{H \in \Delta} \Gamma(X_H, \mathcal{O}) \rightarrow Q(R)$, where the second homomorphism exists due to Lemma 7.3.7 applied to $S = Q(R)$. The first homomorphism exists by Lemma 1.8.4 and is injective by Lemma 1.9.5. So let $f \in \Gamma(X_H, \mathcal{O})$, $H \in \Delta$, be given. The diagonal $X_H \rightarrow X_H \times_X X_H$ is a cover in the Δ -topology, since we have a factorization $(X_H \times_X X_H)_H \rightarrow X_H \rightarrow X_H \times_X X_H$. Hence f is compatible (Lemma 1.6.7) and $f \in \Gamma(X_\Delta, \mathcal{O}_\Delta)$.

Let now $c/b \in Q(R)$ be such that $c \in (b)_\Delta$, say $cH \subseteq bH$, $H = (h_1, \dots, h_m)$. This means that for every h_i there exists $g_i \in H$ such that $ch_i = bg_i$. For i, j it follows that $bg_i h_j = ch_i h_j = bg_j h_i$ and so $g_i h_j = g_j h_i$, since b is a non zero divisor. Therefore $g_i/h_i = g_j/h_j$ in $\Gamma(D_+(h_i h_j), \mathcal{O})$ and so we get a global element in $\Gamma(X_H, \mathcal{O})$. This element maps to c/b under the natural homomorphism.

On the other hand, suppose that $c/b \in Q(R)$ is the image of $q \in \Gamma(X_H, \mathcal{O})$, $H \in \Delta$, $H = (h_1, \dots, h_m)$. Let $h \in H$ be a non zero divisor, and let $q = g/h^r \in \Gamma(D_+(h), \mathcal{O})$. This maps to $g/h^r \in Q(R)$ and so $c/b = g/h^r$. So we may assume that $b \in H$. We have $c/b = g_i/h_i$ in $\Gamma(D_+(bh_i), \mathcal{O})$, and so $ch_i^t \in bH^t$ for some t and all i . Then the ideal H^{tm} shows that $c \in (b)_\Delta$ (as in the proof of Proposition 7.3.1). $\square$

REFERENCES

- [1] M. Artin. *Grothendieck topologies*. Harvard University Math. Dep., 1961.
- [2] M. Artin. On the joins of Hensel rings. *Advances in Math.*, 7:282–296, 1971.
- [3] P. E. Bland. *Topics in torsion theory*, volume 103 of *Math. Research*. Wiley-VCH Verlag, Berlin, 1998.
- [4] M. Blickle and H. Brenner. Integral closure and the submersive Grothendieck topology. *In preparation*, 2006.
- [5] M. Blickle and H. Brenner. Tight topology and tight closure. *In preparation*, 2006.
- [6] H. Brenner. How to rescue solid closure. *J. Algebra*, 265:579–605, 2003.
- [7] H. Brenner. Continuous solutions to algebraic forcing equations. *Submitted*, 2006.
- [8] H. Brenner. Test rings for weak subintegral closure. *In preparation*, 2006.
- [9] M. P. Brodmann and M. P. Sharp. *Local Cohomology*. Cambridge University Press, 1998.
- [10] W. Bruns and J. Herzog. *Cohen-Macaulay Rings, Revised edition*. Cambridge University Press, 1998.
- [11] S. E. Dickson. A torsion theory for abelian categories. *Trans. Amer. Math. Soc.*, 121:223–235, 1966.
- [12] D. Eisenbud. *Commutative Algebra with a View Toward Algebraic Geometry*. Springer-Verlag, 1995.
- [13] S. Faridi. The blowup closure of a set of ideal with applications to TI closure. *J. Algebra*, 249(2):472–511, 2002.
- [14] R. Fedder. F -purity and rational singularity in graded complete intersection rings. *Trans. Amer. Math. Soc.*, 301(1):47–62, 1987.
- [15] R. Fedder and K. Watanabe. A characterization of F -regularity in terms of F -purity. In M. Hochster, C. Huneke, and J. Sally, editors, *Commutative Algebra*, pages 227–245, 1989.
- [16] J. S. Golan. *Torsion theories*, volume 29 of *Pitman Monographs and Surveys in Pure and Applied Mathematics*. John Wiley and Sons, New York, 1986.
- [17] T. G. Goodwillie and S. Lichtenbaum. A cohomological bound for the h -topology. *Amer. J. Math.*, 123:425–443, 2001.
- [18] A. Grothendieck and J. Dieudonné. *Eléments de géométrie algébrique II*, volume 8. Inst. Hautes Études Sci. Publ. Math., 1961.
- [19] A. Grothendieck and J. Dieudonné. *Eléments de géométrie algébrique III*, volume 11, 17. Inst. Hautes Études Sci. Publ. Math., 1961–1963.
- [20] A. Grothendieck and J. Dieudonné. *Eléments de géométrie algébrique IV*, volume 20,24,28,32. Inst. Hautes Études Sci. Publ. Math., 1964–1967.
- [21] A. Grothendieck and J. Dieudonné. *Eléments de géométrie algébrique I*. Springer, 1971.
- [22] A. Grothendieck et al. *Cohomologie locale des faisceaux cohérents et Théorèmes de Lefschetz locaux et globaux* (=SGA2). North Holland, 1968.
- [23] A. Grothendieck et al. *Revêtements Étales et Groupe Fondamental* (=SGA1). Springer, 1970.
- [24] A. Grothendieck et al. *Théorie des topos* (=SGA4) I,II. Springer, 1970.
- [25] N. Hara. Classification of two-dimensional F -regular and F -pure singularities. *Adv. Math.*, 133:33–53, 1998.
- [26] N. Hara and K. I. Watanabe. F -regular and F -pure rings vs. log terminal and log canonical singularities. *J. Algebraic Geom.*, 11(2):363–392, 2002.
- [27] R. Hartshorne. *Algebraic Geometry*. Springer, 1977.
- [28] M. Hashimoto. *Auslander-Buchweitz Approximations of Equivariant Modules*. 2000.
- [29] W. Heinzer, D. Lantz, and K. Shah. The Ratliff-Rush ideals in a Noetherian ring. *Communications in Algebra*, 20(2):591–622, 1992.
- [30] R. C. Heitmann. Plus closure in mixed characteristic. *J. Algebra*, 193(2):688–708, 1997.
- [31] R. C. Heitmann. Extensions of plus closure. *J. Algebra*, 238(2):801–826, 2001.
- [32] R. C. Heitmann. The direct summand conjecture in dimension three. *Ann. Math. (2)*, 156(2):695–712, 2002.
- [33] M. Hochster. Solid closure. *Contemp. Math.*, 159:103–172, 1994.
- [34] M. Hochster. Tight closure in equal characteristic, big Cohen-Macaulay algebras, and solid closure. *Contemp. Math.*, 159:173–196, 1994.
- [35] M. Hochster and C. Huneke. Tight closure, invariant theory, and the Briançon-Skoda theorem. *J. Amer. Math. Soc.*, 3:31–116, 1990.
- [36] M. Hochster and C. Huneke. Infinite integral extensions and big Cohen-Macaulay algebras. *Ann. Math.*, 135:53–89, 1992.

- [37] M. Hochster and J. Roberts. The purity of Frobenius and local cohomology. *Advances in Math.*, 21:117–172, 1976.
- [38] C. Huneke. *Tight Closure and Its Applications*, volume 88 of *CBMS Lecture Notes in Mathematics*. AMS, Providence, 1996.
- [39] C. Huneke. Tight closure, parameter ideals, and geometry. In *Six Lectures on Commutative Algebra*, pages 187–329. Birkhäuser, 1998.
- [40] C. Huneke and G. Lyubeznik. Absolute integral closure in positive characteristic. *ArXiv*.
- [41] D. Katz. Complexes azyclique up to integral closure. *Math. Proc. Camb. Philos. Soc.*, 116(3):401–414, 1994.
- [42] S.-I. Kimura. A cohomological characterization of Alexander schemes. *Invent. Math.*, 137, 1999.
- [43] J. Kollar. *Rational Curves on Projective Varieties*. Springer, 1995.
- [44] M. Maani Shirazi and H. Sharif. Hilbert modules. *Int. J. Pure Appl. Math.*, 20(1):1–7, 2005.
- [45] M. Manaresi. Some properties of weakly normal varieties. *Nagoya Math. J.*, 77, 1980.
- [46] A. Marcelo, F. Marcelo, and C. Rodríguez. Effectively computation of some radicals of submodules of free modules. In *Computer algebra in scientific computing - CASC'99 (Munich)*, pages 287–292, 1999.
- [47] R. L. McCasland and M. E. Moore. On radicals of submodules of finitely generated modules. *Canad. Math. Bull.*, 29(1):37–39, 1986.
- [48] V. B. Mehta and V. Srinivas. Normal F -pure surface singularities. *J. Algebra*, 143(1):130–143, 1991.
- [49] J. S. Milne. *Étale Cohomology*. Princeton University Press, 1980.
- [50] Y. A. Nisnevich. The completely decomposed topology on schemes and associated descent spectral sequences in algebraic K -theory. In *Algebraic K-theory: connections with geometry and topology*, volume 279 of *NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci.*, pages 241–342, 1989.
- [51] D. G. Northcott and D. Rees. Reductions of ideals in local rings. *Proc. Cambridge Philos. Soc.*, 50:145–158, 1954.
- [52] S. Oda and K.I. Yoshida. On the $(2, 3)$ -closure of ideals. *Math. J. Okayama Univ.*, 34:1–12, 1992.
- [53] G. Picavet. Submersion et descente. *J. Algebra*, 103:527–591, 1986.
- [54] D. Pusat-Yilmaz and P. F. Smith. Modules which satisfy the radical formula. *Acta Math. Hungar.*, 95(1-2), 2002.
- [55] L. J. Ratliff, JR. Δ -closure of ideals and rings. *Trans. Amer. Math. Soc.*, 313(1):221–247, 1989.
- [56] L. J. Ratliff, JR. and D. E. Rush. Two notes on reductions of ideals. *Indiana Univ. Math. J.*, 27:929–934, 1978.
- [57] L. J. Ratliff, JR. and D. E. Rush. Δ -reductions of modules. *Comm. Alg.*, 21(8):2667–2685, 1993.
- [58] L. J. Ratliff, JR. and D. E. Rush. Asymptotic primes of delta closures of ideals. *Comm. Alg.*, 30(3):1513–1531, 2002.
- [59] L. Reid and M. A. Vitulli. The weak subintegral closure of a monomial ideal. *Comm. Algebra*, 27(11):5649–5667, 1999.
- [60] P. Roberts. A computation of local cohomology. *Contemp. Math.*, 159:351–356, 1994.
- [61] G. Scheja and U. Storch. *Lokale Verzweigungstheorie*, volume 5. Schriftenreihe des mathematischen Institutes der Universität Freiburg, 1974.
- [62] G. Scheja and U. Storch. *Lehrbuch der Algebra, Teil 2*. Teubner, 1988.
- [63] R. Y. Sharp and N. Nossem. Ideals in a perfect closure, linear growth of primary decompositions, and tight closure. *Trans. Amer. Math. Soc.*, 356(9):3687–3720, 2004.
- [64] A. Suslin and V. Voevodsky. Singular homology of abstract varieties. *Invent. Math.*, 123:61–94, 1996.
- [65] L. Vas. *Torsion theories for group Von Neumann algebras*. PhD thesis, University of Maryland at College Park, 2002.
- [66] A. Vistoli. Notes on Grothendieck topologies, fibered categories and descent theory. In *B. Fantechi, L. Göttsche, L. Illusie, S. Kleiman, N. Nitsure, A. Vistoli, Fundamental Algebraic Geometry: Grothendieck's FGA explained*. Math. Surveys and Monographs 123.
- [67] M. A. Vitulli and J.V. Leahy. The weak subintegral closure of an ideal. *J. Pure and Applied Algebra*, 141:185–200, 1999.
- [68] V. Voevodsky. Homology of schemes. *Selecta Math. (N.S.)*, 2:111–153, 1996.
- [69] M. E. Walker. The primitive topology of a scheme. *J. Algebra*, 201(2):656–685, 1998.

DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF SHEFFIELD, HICKS BUILDING, HOUNSFIELD
ROAD, SHEFFIELD S3 7RH, UNITED KINGDOM
E-mail address: `H.Brenner@sheffield.ac.uk`